**Karolinska
Institutet**

Karolinska Institutet

<http://openarchive.ki.se>

This is a Peer Reviewed Accepted version of the following article, accepted for publication in Environmental science & technology.

2017-07-07

Detection of benz[j]aceanthrylene in urban air and evaluation of its genotoxic potential

Lim, Hwanmi; Mattsson, Åse; Jarvis, Ian W H; Bergvall, Christoffer; Bottai, Matteo; Morales, Daniel A; Kummrow, Fabio; Umbuzeiro, Gisela A; Stenius, Ulla; Westerholm, Roger; Dreij, Kristian

Environ Sci Technol. 2015 Mar 3;49(5):3101-9.

<http://doi.org/10.1021/es505458g>

<http://hdl.handle.net/10616/45982>

If not otherwise stated by the Publisher's Terms and conditions, the manuscript is deposited under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

**Karolinska
Institutet**

This is the peer reviewed version of the following article:
Environ Sci Technol. 2015 Mar 3;49(5):3101-9, which has
been published in final form at
<http://dx.doi.org/10.1021/es505458g>

**Detection of benz[j]aceanthrylene in urban air and
evaluation of its genotoxic potential.**

**Lim, H; Mattsson, Å; Jarvis, Ian WH; Bergvall,
Christoffer; Bottai, Matteo; Morales, DA; Kummrow, F;
Umbuzeiro, GA; Stenius, Ulla; Westerholm, Roger; Dreij,
Kristian**

Access to the published version may require subscription.
Published with permission from: **American Chemical Society**

Detection of benz[*j*]aceanthrylene in urban air and evaluation of its genotoxic potential

Hwanmi Lim^{1#}, Åse Mattsson^{2#}, Ian WH Jarvis², Christoffer Bergvall¹, Matteo Bottai³, Daniel A Morales⁴, Fábio Kummrow⁴, Gisela A Umbuzeiro⁴, Ulla Stenius², Roger Westerholm¹ and Kristian Dreij^{2*}

¹ Department of Analytical Chemistry, Stockholm University, Svante Arrhenius väg 16 SE-106 91, Stockholm, Sweden.

² Unit of Biochemical Toxicology, Institute of Environmental Medicine, Karolinska Institutet, Box 210, SE-171 77 Stockholm, Sweden.

³ Unit of Biostatistics, Institute of Environmental Medicine, Karolinska Institutet, Box 210, SE-171 77 Stockholm, Sweden.

⁴ Faculty of Technology, State University of Campinas (Unicamp), Limeira, SP, Brazil.

[#] These authors contributed equally

^{*} Corresponding author. E-mail: Kristian.Dreij@ki.se; Phone: +46 8 524 875 66

ABSTRACT

Benz[*j*]aceanthrylene (B[*j*]A) is a cyclopenta-fused polycyclic aromatic hydrocarbon with strong mutagenic and carcinogenic effects. We have identified B[*j*]A in air particulate matter in samples collected in Stockholm, Sweden and in Limeira, Brazil using LC-GC/MS analysis. Determined concentrations ranged between 1.57-12.7 and 19.6-30.2 pg/m³ in Stockholm and Limeira, respectively, which was 11-30 times less than benzo[*a*]pyrene (B[*a*]P) concentrations. Activation of the DNA damage response was evaluated after exposure to B[*j*]A in HepG2 cells in comparison to B[*a*]P. We found that significantly lower concentrations of B[*j*]A was needed for an effect on cell viability compared to B[*a*]P and equimolar exposure resulted in significant more DNA damage with B[*j*]A. Additionally, levels of γ H2AX, pChk1, p53, pp53 and p21 proteins were higher in response to B[*j*]A than B[*a*]P. Based on dose response induction of pChk1 and γ H2AX, B[*j*]A potency was 12.5 and 33.3 higher than B[*a*]P, respectively. Although B[*j*]A levels in air were low, including B[*j*]A in the estimation of excess lifetime cancer risk increased the risk up to 2-fold depending on which potency factor for B[*j*]A was applied. Together our results show that B[*j*]A could be an important contributor to the cancer risk of air PM.

INTRODUCTION

Particulate matter (PM) from urban air contains a mixture of different chemicals that can interact and cause adverse effects to human health. Outdoor air pollution and its PM component have been classified as carcinogenic to humans by IARC.¹ One group of chemicals that are found in air PM is the polycyclic aromatic hydrocarbons (PAHs), ubiquitous environmental contaminants that are formed during incomplete combustion of organic matter. PAHs in the environment are of concern because of their carcinogenic activity and several individual and mixtures of PAHs have been classified as possible or probable carcinogens to humans.² To date, benzo[*a*]pyrene (B[*a*]P) is the only individual PAH classified as carcinogenic to humans by IARC.²

Health risk assessment of PAH mixtures is often carried out using B[*a*]P for comparison as its toxicological profile has been extensively characterized. One approach to perform risk assessment is based on additivity of toxic equivalency factors (TEFs) or relative potency factors (RPFs) where the carcinogenic potential is expressed relative to B[*a*]P.³ A second method is to use B[*a*]P as a surrogate marker for all PAHs. This has been applied to air⁴ and the European Commission Air Quality Standards has put a target value of PAHs in air expressed as 1 ng B[*a*]P/m³ over an exposure period of one year (Directive 2004/107/EC). However, as has recently been discussed, these risk assessment approaches have limited application to studying mixture effects and are likely to misestimate the actual risk to human health.⁵ Most PAHs have not been assigned TEF/RPF values or their contribution to mixtures has been overlooked due to a lack of sensitivity in detection methods. This is particularly applicable to PAHs which are found at low levels in the environment but have demonstrated high carcinogenic potentials. An example of this is dibenzo[*def,p*]chrysene (DBC, also

known as dibenzo[*a,l*]pyrene), the most potent PAH known to date⁶⁻⁷ yet not included among the US EPA priority PAHs nor routinely used as an indicator of carcinogenic PAHs.^{3,8-9} Recent studies have suggested however that the quantification of DBC in urban air has become more common as a result of improved detection methods.¹⁰⁻¹⁴ For the reasons stated above it is important to analyze and determine the levels of PAHs with high carcinogenic potential in environmental samples to include them in the evaluation of mixture exposure risks.

One poorly studied PAH that has shown high carcinogenic potential is benz[*j*]aceanthrylene (B[*j*]A).¹⁵⁻¹⁶ B[*j*]A is a cyclopenta-fused and bay-region containing PAH. As with other PAHs B[*j*]A requires metabolic activation in order to exert its biological activity and studies have identified two major routes of activation: epoxidation of the cyclopenta-ring, resulting in the B[*j*]A-1,2-epoxide, and diol-epoxidation of the bay-region, resulting in the B[*j*]A-9,10-diol-7,8-epoxide.^{15,17-18} The former is the major metabolite found as a 1,2-diol in human and rat liver tissues and cells.¹⁹⁻²⁰ B[*j*]A is a potent bacterial and mammalian cell mutagen and is more tumorigenic than B[*a*]P in SENCAR mice and in A/J mice lung.¹⁵⁻¹⁶ Trace amounts of B[*j*]A have been found in the emissions from coal-fired residential furnaces and hard-coal combustion²¹⁻²² but not in wood smoke PM.²³ Substituted B[*j*]As, such as 3-methyl- and 1,2-dihydro-3-methyl-B[*j*]A, have been determined in coal tar pitch, cigarette smoke and air PM.²⁴⁻²⁷ To date, very few studies have analyzed the levels of B[*j*]A in air PM²⁸ and to the authors' knowledge no epidemiological studies have investigated the carcinogenicity of B[*j*]A.

In the present study we have quantified the concentration of B[*j*]A in urban air PM collected in Stockholm, Sweden and Limeira, Brazil. Our analytical setup allowed for detection and

quantification of very low levels of B[j]A in air PM samples from the two city locations. We have also investigated the effects of B[j]A on cell viability, DNA damage and activation of DNA damage signaling in human-derived hepatocellular carcinoma (HepG2) cells to evaluate its potency relative to B[a]P. Taken together our results show that although levels of B[j]A are very low in the air PM samples, it is significantly more potent than B[a]P and thus is likely to be an important contributor to the cancer risk of air PM.

MATERIAL AND METHODS

Chemicals and reagents

Hexane, acetone and toluene (HPLC grade) were obtained from Rathburn Ltd. (Walkerburn Scotland) and dodecane (anhydrous, $\geq 99\%$) from Sigma-Aldrich (St. Louis, MO, USA). Synthesized B[j]A (purity determined to be at least 95% by NMR) was kindly provided by Dr. Avram Gold and Dr. Zhenfa Zhang, University of North Carolina, Chapel Hill, NC, USA. B[a]P-D₁₂ (98.7%) was supplied by Chiron AS (Trondheim, Norway) and benzo[b]fluoranthene (100%) and benzo[k]fluoranthene (98.3%) were obtained from Chem Service (West Chester, PA, USA). B[a]P (97.6%), benzo[e]pyrene (99.7%) and perylene (99.5%) were from Sigma-Aldrich (St. Louis, MO, USA). Benzo[a]fluoranthene was obtained from National Institute of Standards and Technology (Gaithersburg, MD, USA), and benzo[j]fluoranthene (100%) was from Larodan Fine Chemicals AB (Limhamn, Sweden). All cell culture reagents were supplied by Gibco (Life Technologies, Stockholm, Sweden). Antibodies used for Western blotting were phospho-Chk1 (Ser317), phospho-H2AX (Ser139) and phospho-p53 (Ser15) from Cell Signaling Technology (Beverly, MA, USA), and p53 (DO-1), Cdk2 (M2) and secondary anti-mouse and anti-rabbit antibodies from Santa Cruz

(Santa Cruz, CA, USA). For immunocytochemistry, anti-phospho H2AX (Ser139, clone JBW301) was from EMD Millipore Corp. (Billerica, MA, USA) and secondary Alexa Fluor 594 goat anti-mouse from Molecular Probes (Life Technologies, Stockholm, Sweden).

Air sampling

Two collection sites for air PM were used: Stockholm University, Stockholm, Sweden and the Faculty of Technology, UNICAMP, Limeira, Brazil. The Stockholm sampling has been described previously.¹⁰ Briefly, three air PM samples (STO1, STO2 and STO3) were collected on fluorocarbon coated glass fiber filters ($\varnothing = 235$ mm, Fiberfilm Filters, Pallflex, Pall Corporation, Putnam, CT, USA) at roof top level using an in-house pump device for three or seven days at an average flow rate of 70.6 m³/hr. The filters were desiccated for at least 24 h before and after the sampling. In Limeira, two air PM samples (LMR1 and LMR2) were collected on glass fiber filters (254 × 233 mm, 0.33 mm pore size, Energética Ind. Com. LTDA, Rio de Janeiro, RJ, Brazil) at the street level. The sampling was performed with a high-volume sampler (Energética Ind. Com. LTDA, Rio de Janeiro, RJ, Brazil) for 24 h at an average flow rate of 67.8 m³/hr. Until extraction filters from both Stockholm and Limeira were wrapped in aluminum and stored at -20 °C.

Sample preparation and LC-GC/MS analysis

A pressurized liquid extraction system (ASE 200 Accelerated Solvent Extraction System, Dionex Co., Sunnyvale, CA, USA) was used for the filter extraction. Samples were extracted with toluene for 5 consecutive cycles of 30 min. Then, 0.6 mL of the extract (corresponding to 56.1 m³, 214.6 m³, 144.5 m³, 34.5 m³ and 26.3 m³ for STO1, STO2, STO3, LMR1 and

LMR2, respectively) was spiked with 60 µL of 442 pg/µL B[a]P-D₁₂ and applied to an SPE column (silica, 100 mg, IST Isolute, Biotage, Cardiff, UK). A PAH enriched fraction was obtained by elution with 2 mL of hexane. The final hexane eluate was evaporated until 100 µL under a gentle nitrogen stream and transferred to a 300 µL micro vial for LC-GC/MS analysis. All sample preparations were performed in triplicate. The extraction and SPE clean-up procedures²⁹ and the LC-GC/MS method³⁰ are described in detail elsewhere. Blank samples were prepared in the same manner using blank filters.

Cell culture and exposure to PAHs

Human-derived hepatocellular carcinoma cells (HepG2) were obtained from American Type Culture Collection (Rockville, MD, USA) at passage 78 and used within 20 passages for all experiments. The rationale for using this cell line in these studies is the capacity to metabolize PAHs³¹ and previously demonstrated response to low levels of PAHs.³²⁻³⁴ Cells were cultured in Minimal Essential Medium supplemented with 10% fetal bovine serum, 1 mM sodium pyruvate, 0.1 mM non-essential amino acids, 100 units/ml *penicillin* and 0.1 mg/ml *streptomycin*, and maintained at 37°C in 5% CO₂. Cells were exposed to solvent control (0.1% DMSO), B[j]A or B[a]P for up to 48 h.

MTT assay

Cell viability was assessed using MTT assay as described previously.³⁵ Briefly, HepG2 cells (0.3 x 10⁵) were plated in 24-well plates (Corning Inc., Corning, NY, USA) and allowed to grow for 24 h. Cells were exposed to solvent control, B[j]A or B[a]P for 48 h in 1.5 ml medium and thereafter incubated with 0.5 mg/ml MTT reagent in 250 µl HBSS for 4 h.

Subsequently, formazan crystals were dissolved in 500 μ l DMSO (15 min, shaking) and plates were spectrophotometrically analyzed at 570 nm with a reference wavelength at 690 nm (Wallac Victor³ V 1420 multilabel counter, Perkin Elmer, Waltham, MA, USA). Results are presented as percent of solvent control and EC₅₀ was established using non-linear regression in GraphPad Prism 6.0 (GraphPad Software, Inc., La Jolla, CA, USA).

Comet assay

The alkaline comet assay was performed as described previously.³⁶ In brief, HepG2 cells (0.3×10^5) were plated in 24-well plates (Corning Inc., Corning, NY, USA) and grown for 24 h prior to exposure to solvent control, B[j]A or B[a]P in 1.5 ml for 48 h. Cells were trypsinized and mixed with 0.75% w/v low melting point agarose and smeared on slides pre-coated with 0.3% agarose. Slides were incubated in cold lysis buffer (1% Triton X-100, 2.5 M NaCl, 10 mM Tris, 0.1 M EDTA, pH 10) for 1 h on ice followed by incubation in cold alkaline solution (0.3 M NaOH, 1 mM EDTA, pH >13) for 40 min on ice. Electrophoresis was run in the alkaline solution at 29 V (1.15 V/cm) for 30 min and thereafter slides were neutralized in 0.4 M Tris-HCl, pH 7.5, dried overnight and fixed in methanol for 5 min. After ethidium bromide staining, at least 100 cells were scored per treatment using a Leica DMLB fluorescent microscope and Comet Assay IV (Perceptive Instruments Ltd., Haverhill, UK).

Immunocytochemistry

HepG2 cells (0.2×10^4) were plated in 12-well plates containing 13 mm glass cover slips and grown for 24 h before exposure to solvent control, B[j]A or B[a]P in 2.0 ml for 48 h. Cells were washed and fixed in 4% paraformaldehyde for 20 min at room temperature and then

permeabilized with 0.2% Triton X-100 for 10 min followed by blocking in 2% bovine serum albumin in TBS-Tween 0.1% with 5% normal goat serum. Incubation with primary antibody was overnight at 4°C followed by washing and incubation with secondary antibody for 1 h in the dark. Cell nuclei were counterstained with 4',6-diamidino-2-phenylindole (DAPI) prior to mounting with Vectashield H-1000 mounting medium (Vector Labs, Burlingame, CA, USA). Images were captured using a 63x oil immersion objective on an LSM 510 Meta confocal laser scanning microscope (Zeiss, Göttingen, Germany). Cell foci were counted using CellProfiler cell image analysis software version 2.1.0 (MIT and Broad Institute, Cambridge, MA, USA).³⁷

For determining statistical effects in the immunocytochemistry experiments, we estimated the expected mean count in each exposure group with a linear regression model for each experiment separately. The experimental groups were introduced by means of indicator variables and count data analyzed marginally with respect to the experiment's day. We used the sandwich robust estimator for the standard errors³⁸ which is robust to misspecification of the modeling assumptions (e.g. homoscedasticity). The analyses were performed in Stata version 13 (StatCorp, Collage Station, TX, USA).

Western Blotting

Western blotting was performed as described previously³² with minor modifications. HepG2 cells (4.0×10^5) were plated in 6-well plates (Corning Inc., Corning, NY, USA) and cultured for 72 h. Cells were exposed to solvent control, B[j]A or B[a]P for 48 h, washed with ice-cold PBS and scraped into IPB-7 buffer containing protease inhibitors. Protein concentration was measured and samples subjected to standard SDS-PAGE. Subsequently, proteins were

transferred to PVDF membrane (Bio-Rad, Hercules, CA) and protein levels were detected using specific antibodies and visualized using enhanced chemiluminescence (Amersham GE Healthcare, Bio-Sciences AB, Uppsala, Sweden). Densitometric analysis was performed using ImageJ software version 1.48f (National Institute of Health, USA).

Cancer risk assessment

In order to estimate the excess lifetime cancer risk for inhalation exposure to PAHs in air PM, B[a]P equivalency concentrations ($B[a]P_{eq}$) were determined using current RPF scales (Table S1).³⁹ The contribution to the cancer risk from B[j]A was estimated using published RPFs^{15-16,40-41} and potency factors based on H2AX and Chk1 activation. The cancer risk was determined by multiplying PAH concentration with a unit risk (per 100 000 people) for B[a]P set by WHO⁴ to 8.7×10^{-5} ng/m³ (based on epidemiological study of coke oven workers) using the equation below.

$$\text{Cancer risk} = \sum([\text{PAH}] \times \text{RPF}_{\text{PAH}}) \times \text{unit risk}$$

Statistics

All data presented are means $\pm$ standard error (SE). With the exception of the immunocytochemistry analysis (described above), One-Way ANOVA with Bonferroni's t-test correction was used to determine statistical significance ($p < 0.05$).

RESULTS

Detection limit and linearity

A series of six B[j]A standard solutions were injected on the LC-GC/MS system in triplicate and a calibration curve of the area ratio (B[j]A/B[a]P-D₁₂) against the amount of B[j]A was plotted in the range of 2–493 pg B[j]A spiked with 10 µL of 442 pg/µL B[a]P-D₁₂. A GC/MS chromatogram of a calibration standard is shown in Fig. S1. The coefficient of determination (R^2) was 0.9975 and the limit of detection (at a signal to noise ratio (S/N) of 4.2) and limit of quantification (at a S/N of 14.9) were determined to be 2.5 and 12 pg, respectively.

Determination of B[j]A in air PM

Representative GC/MS chromatograms of air PM samples from Stockholm and Limeira are shown in Fig. 1A and 1B, respectively. The B[j]A peak was well-separated from the other seven molecular weight 252 Da isomers detected, whilst partially co-eluting with an unknown peak displaying a characteristic PAH mass spectrum (mother ion m/z 268). Together with mass spectra, the peak identification was confirmed by comparing the retention time of B[j]A from the original and spiked samples in the GC/MS chromatograms as shown in Fig. 1C and 1D. No detectable amount of B[j]A was found in blank samples generated by extracting the different filter types used for sampling air PM in Stockholm and Limeira (Fig. S2). Sample identities and collection details including levels of B[a]P and B[j]A in the air PM from Stockholm and Limeira are summarized in Table 1. Determined B[j]A concentrations were between 11 to 30 times less than those of B[a]P. The concentrations of all PAHs determined in the air PM samples are shown in Table S2. The levels of PAHs in STO1 and LMR2 have previously been published.⁴²

B[j]A exerts stronger effects on cell viability than B[a]P consistent with increased DNA damage

In vitro studies have shown that B[j]A is a genotoxic agent and a more potent inducer of apoptosis than B[a]P.^{17,43} In agreement, we found that B[j]A was more toxic to the cells than B[a]P in the MTT assay (Fig. 2A). The estimated EC₅₀ value for cell viability was 0.39±0.20 and 1.45±0.09 µM for B[j]A and B[a]P, respectively. A statistically significant decrease in cell viability compared to control levels was observed after exposure to 2.0 µM B[a]P or 0.3 µM B[j]A (p<0.001). There was also a significant difference between cells exposed to equimolar concentrations of B[j]A and B[a]P at 0.3 and 1.0 µM (p<0.001).

Increased potency between different PAHs has previously been shown to be intrinsically linked to their ability to form DNA damage and the persistence of this damage/resistance to repair.⁴⁴⁻⁴⁶ We therefore hypothesized that in a comparable manner, the increased toxicity of B[j]A over B[a]P correlates with differences in DNA damage levels. A single exposure time of 48 h was chosen for these analyses as we have previously used this time point to study the effects of individual and mixtures of PAHs on DNA damage.³²⁻³³ Significantly increased levels of DNA damage were observed in the Comet assay after exposure to 1.0 µM B[a]P (p<0.001) or 0.3 µM B[j]A (p<0.001) when compared to control levels (Fig. 2B). No significant changes in DNA damage levels were observed after exposure to 0.3 µM B[a]P (Fig. 2B) or 0.1 µM B[j]A (data not shown). Comparing equimolar concentrations showed that B[j]A induced significantly more DNA damage than B[a]P (p<0.001). These results were further confirmed by immunostaining for formation of phosphorylated H2AX (Ser139) foci (γH2AX). Visual comparison between treatments showed more foci formation in response to B[j]A>B[a]P>DMSO control (Fig. 2C). Both the B[a]P group and the B[j]A group had

higher foci rate compared to control group, 5.7 and 17.2 times more, respectively ($p < 0.001$) (Fig. 2D). The number of foci in B[j]A exposed cells were also significantly higher from that in the equimolar B[a]P group ($p < 0.001$). Taken together these data show that B[j]A is significantly more potent than B[a]P in reducing cell viability and that this most likely results from increased damage to DNA.

B[j]A induces a stronger activation of DNA damage signaling than B[a]P

In agreement with the Comet assay and γ H2AX foci results, our data revealed B[j]A to be a more potent inducer of DNA damage signaling than B[a]P. At 1 μ M, B[j]A induced phosphorylation of Chk1 Ser317 (pChk1), γ H2AX and p53 Ser15 (pp53) to a higher extent than B[a]P (Fig. 3A). Total p53 level was also more elevated in response to B[j]A compared to B[a]P as well as the protein level of the cell cycle regulator p21. To further study the difference in potencies between B[j]A and B[a]P we applied a dose response analysis for the induction of pChk1, pp53 and γ H2AX. As can be seen, B[j]A induced phosphorylation of all proteins at lower concentration compared to B[a]P (Fig. 3B and 3C). To allow for a quantitative comparison of potency, densitometric analysis of pChk1 and γ H2AX levels (Fig. 3C) was used to estimate the concentrations of B[j]A and B[a]P required to induce a particular fold induction of the proteins (Table S3). Due to no visible bands of pp53 in control and at lower concentrations of B[j]A and B[a]P proper densitometry analysis and comparison of potencies could not be accurately performed. For comparison, we included previous data from DBC exposures.³² DBC induces pChk1 at lower concentrations compared to B[j]A whereas γ H2AX was induced similarly by DBC and B[j]A (Fig. 3C). The results showed that levels of pChk1 increased at lower levels for B[j]A compared to γ H2AX, in

agreement with our previous data.³² On average B[j]A was 12.5- and 33.3-fold more potent inducer of pChk1 and γ H2AX levels, respectively, compared to B[a]P.

Contribution of B[j]A to air PM cancer risk

If only the B[a]P concentration in air PM was considered the estimated excess lifetime cancer cases per 100 000 people was 0.15-2.5 (ex. STO1: $0.289 \text{ ng B[a]P/m}^3 \times 8.7 \times 10^{-5} = 2.5 \times 10^{-5}$) and 4.9-7.8 for Stockholm and Limeira, respectively (Table 2). Next we calculated the excess lifetime cancer cases based on all the determined PAHs in the air PM samples (except B[j]A) with an assigned RPF value. Compared to the excess lifetime cancer cases based on levels of B[a]P alone, the cases increased to 0.77-11.3 and 16.2-22.7 per 100 000 people for Stockholm and Limeira, respectively (Table 2). The additional contribution to the cancer risk by including B[j]A in the B[a]P_{eq} levels was estimated using two different published RPFs for B[j]A of 10 and 60^{15-16,40-41} and our potency factor of 30 based on H2AX activation (Table 2). The published RPF of 10 was similar to our potency factor based on activation of Chk1. The results showed that inclusion of B[j]A increased the estimated cancer risk of the air PM. Depending on which potency factor that was applied, inclusion of B[j]A resulted in an up to doubling of expected excess lifetime cancer cases (Table 2). This provides convincing evidence that B[j]A contributes to the carcinogenic potency of urban air PM and warrants further investigation.

DISCUSSION

Concentrations of B[j]A in urban air PM

337 The present study determined the atmospheric B[j]A concentration in air PM collected at two
338 different locations, Stockholm, Sweden and Limeira, Brazil. The air PM from Limeira
339 contained higher levels of PAHs compared to the samples from Stockholm. Besides the
340 probable difference in PAH levels due to emission profiles in the respective areas, the
341 samples from Limeira were collected at street level while in Stockholm sampling was
342 performed on a roof top. PAHs at the Limeira site can be mostly attributed to heavy traffic
343 and biomass burning whereas traffic is a main source at the Stockholm site.⁴⁷ The PAH levels
344 in Stockholm air PM displayed large seasonal changes during the different sampling periods:
345 the B[a]P concentration was 17.3 pg/m³ in September and increased to 49.5 and 289 pg/m³ in
346 December and January, respectively. The B[j]A concentration followed the same trend: 1.57,
347 3.14 and 12.7 pg/m³, as well as the other measured PAHs and this is in line with recently
348 reported PAH levels from the Stockholm atmosphere showing a high annual variability with
349 higher concentrations during the colder parts of the year.⁴⁸ Higher PAH levels during the
350 winter season in cities around the world has been attributed to meteorological conditions such
351 as inversion and lower mixing layer, less efficient atmospheric reactions and increase in
352 emissions from domestic heating.⁴⁹ The two air PM samples from Limeira were both
353 collected in July and had higher concentration of B[a]P, B[j]A and of most of the other
354 investigated PAHs with a few exceptions compared to the Stockholm samples. A previous
355 study of air PM PAHs in Saitama City, Japan, reported measurements of 37 PAHs including
356 B[a]P, B[j]A and DBC and to our knowledge it is the only other study reporting measurement
357 of B[j]A in urban air.²⁸ The concentrations of these PAHs in total air PM was 270, 130 and
358 70 pg/m³ for B[a]P, B[j]A and DBC, respectively. The levels of B[a]P in Saitama (sampling
359 period November-December 2013) was similar to STO1, exceeded STO2 and STO3 and was
360 less than LMR1 and LMR2. For B[j]A and DBC the levels were much higher in Saitama
361 compared to all our investigated samples. Further studies to investigate the generation and

deposition of PAHs in the different locales might explain the observed differences in B[j]A and DBC levels.

Stronger activation of DNA damage and DNA damage signaling after exposure to B[j]A

In HepG2 cells, B[j]A was significantly more toxic than B[a]P. This is in agreement with a previous study performed in mouse Hepa1c1c7 hepatoma cells showing that B[j]A is a stronger inducer of apoptosis compared to B[a]P.⁴³ As mentioned before, previous studies have correlated increased potency of PAHs with the ability to form DNA adducts and avoidance of repair. The Comet assay revealed that B[j]A was significantly more genotoxic than B[a]P at equimolar concentrations and this is in line with earlier studies showing that B[j]A induces more DNA adducts in rat and human liver microsomes compared to B[a]P.¹⁹⁻²⁰ Comparing the cytotoxic and genotoxic potencies of the two compounds showed that the cytotoxic doses of B[j]A were also genotoxic, this was not true for 1 μ M B[a]P which was genotoxic but not cytotoxic. The increase in genotoxicity was further confirmed by immunostaining for phosphorylated H2AX (γ H2AX). H2AX is rapidly phosphorylated (at Ser139) at sites of DNA damage, both in response to strand breaks and stable DNA adducts, forming γ H2AX foci that can be detected by immunocytochemistry.⁵⁰⁻⁵¹ Furthermore, phosphorylation of H2AX in the early stages of damage detection is attributed to the propagation of DNA damage signaling and hence we also analyzed levels of γ H2AX by Western blotting alongside other major DNA damage signaling proteins including Chk1, p21 and p53. We found B[j]A to be a stronger inducer of all investigated signaling proteins.

From the dose response curves of pChk1 and γ H2AX we calculated the PAH concentrations needed to induce a fold change in activated protein levels as described earlier.³² These

proteins have previously been shown to be sensitive markers for studying genotoxic potencies of both individual and mixtures of PAHs.^{32-33,52} We calculated the difference in potency of B[j]A compared to B[a]P to be 12.5- and 33.3-fold for pChk1 and γ H2AX, respectively. Previous studies have reported RPFs of 10 and 60 for B[j]A. The RPF of 10 is based on Ames test where an approximate 10-fold lower concentration of B[j]A compared to B[a]P induce optimal activity in S9 dependent test.⁴⁰⁻⁴¹ Similarly in SENCAR mice treated topically with B[j]A tumor initiation was 12 times higher compared to B[a]P.¹⁶ The RPF of 60 is based on B[j]A inducing 16-60 times more lung tumors in A/J mice subjected to a single intraperitoneal injection relative to B[a]P.¹⁵ These tumor data are in the same range as the fold induction data generated in the present study from the phosphorylation of Chk1 and H2AX. It has previously been shown for DBC that fold induction data generated from DNA damage signaling *in vitro* is in agreement with animal experiments regarding carcinogenic potency suggesting that activation of the DNA damage response could serve as a marker for *in vitro* testing of PAH potency.^{5,32} The higher sensitivity of pChk1 and γ H2AX compared to Comet assay data and the close agreement with published RPFs motivated us to include our potency factors based on activation of DNA damage signaling in the cancer risk estimation as further discussed below.

Risk assessment of air PM and contribution of B[j]A to lifetime cancer risk

PAHs have been suggested to be the group of airborne contaminants that contribute most to human health risk. In urban sites 83-94% of the health risk was related to particle associated compounds and of these PAHs was responsible for 99% of the risk.⁵³ An investigation of excess lifetime cancer risk in different age categories in Cordoba, Argentina, showed that already in the age category 1-6 years one child out of one million would get cancer as a result

of airborne PAH exposure.⁵⁴ In the present study the excess lifetime cancer cases was estimated based on B[a]P_{eq} levels and the WHO unit risk value 8.7×10^{-5} ng B[a]P /m³.⁴ We showed that the air samples from Limeira were more polluted by PAHs than the Stockholm samples which was also reflected in a higher number of expected excess lifetime cancer cases, 17.9-38.4 and 0.91-18.0 cases per 100 000 people, respectively. Furthermore, our results showed that B[j]A greatly contributes to the cancer risk of air PM despite the low concentrations that we detected. Although B[j]A levels in air were low, including B[j]A in the estimation of excess lifetime cancer risk increased the risk up to 2-fold depending on which potency factor for B[j]A was applied. The importance of including highly potent PAHs, even though found in low levels, in the cancer risk assessment of air PM is further confirmed by recent studies. Layshock *et al.*, showed that excluding the dibenzopyrenes when assessing air quality most likely results in a substantial underestimation of the health risk.¹² A calculated increase in lifetime risk of developing lung cancer for residents in Beijing, China, due to PAHs in air PM, may be 1 out of 10 000 to over 6 out of 100 and half of that risk was estimated to be related to levels of the dibenzopyrenes.¹² The importance of including high-molecular weight PAHs, such as the dibenzopyrenes, has also been emphasized in other studies showing a significant contribution to the estimated excess lifetime cancer risk of air PM.^{11,13,55} It should be noted that B[j]A was not included in any of the above mentioned studies which probably further resulted in an underestimation of the cancer risk.

In this study we focused on excess lifetime cancer risk from exposure to inhaled air PM PAHs. We are aware of the limitations in the estimation such as the shortcomings of using RPFs which in the case of B[j]A are based on data from Ames test and *in vivo* dermal and intraperitoneal exposure when assessing the cancer risk for inhaled air PM. Further to give a

more comprehensive view of excess lifetime cancer risk, the PAH levels should be monitored over the whole year to include the seasonal variations in PAH constituents. It is very apparent in the three Stockholm samples that the time of sampling greatly affects the PAH levels and in turn the cancer risk estimate.

SUPPORTING INFORMATION

Supporting information available: Tables S1-S3 (air PM B[a]P_{eq} levels, chemical analysis of air PM samples and fold induction of DNA damage signaling) and Figures S1-S2 (GC/MS chromatograms). This material is available free of charge via the internet at <http://pubs.acs.org>.

ACKNOWLEDGEMENTS

The authors are grateful to Dr. Avram Gold and Dr. Zhenfa Zhang from the University of North Carolina at Chapel Hill, USA for kindly providing the B[j]A standard for the present study. This work was supported by Stockholm University, Karolinska Institutet and the Swedish Research Council Formas (R&D project No. 2012-478).

REFERENCES

1. Loomis, D.; Grosse, Y.; Lauby-Secretan, B.; El Ghissassi, F.; Bouvard, V.; Benbrahim-Tallaa, L.; Guha, N.; Baan, R.; Mattock, H.; Straif, K., The carcinogenicity of outdoor air pollution. *Lancet Oncol* **2013**, *14* (13), 1262-3.

- 461 2. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans, Some
462 Non-Heterocyclic Polycyclic Aromatic Hydrocarbons and Some Related Exposures. IARC
463 Monographs on the Evaluation of Carcinogenic Risks to Humans. International Agency for
464 Research on Cancer, Ed. IARC Press: Lyon, France., 2010; Vol. 92.
- 465 3. Boström, C. E.; Gerde, P.; Hanberg, A.; Jernström, B.; Johansson, C.; Kyrklund, T.;
466 Rannug, A.; Törnqvist, M.; Victorin, K.; Westerholm, R., Cancer risk assessment, indicators,
467 and guidelines for polycyclic aromatic hydrocarbons in the ambient air. *Environ Health*
468 *Perspect* **2002**, *110* (Suppl 3), 451-88.
- 469 4. World Health Organisation (WHO), Air Quality Guidelines for Europe, Second
470 Edition. WHO Regional Publications, European Series, No. 91. World Health Organization
471 (WHO) Regional Office for Europe, Ed. WHO: Copenhagen, Denmark., 2000.
- 472 5. Jarvis, I. W. H.; Dreij, K.; Mattsson, A.; Jernström, B.; Stenius, U., Interactions
473 between polycyclic aromatic hydrocarbons in complex mixtures and implications for cancer
474 risk assessment. *Toxicology* **2014**, *321*, 27-39.
- 475 6. Luch, A., On the impact of the molecule structure in chemical carcinogenesis. In
476 *Molecular, Clinical and Environmental Toxicology*, Luch, A., Ed. Birkhäuser Basel: 2009;
477 Vol. 99, pp 151-179.
- 478 7. Siddens, L. K.; Larkin, A.; Krueger, S. K.; Bradfield, C. A.; Waters, K. M.; Tilton, S.
479 C.; Pereira, C. B.; Lohr, C. V.; Arlt, V. M.; Phillips, D. H., et al., Polycyclic aromatic
480 hydrocarbons as skin carcinogens: comparison of benzo[a]pyrene, dibenzo[def,p]chrysene
481 and three environmental mixtures in the FVB/N mouse. *Toxicol Appl Pharmacol* **2012**, *264*
482 (3), 377-86.
- 483 8. Brown, A. S.; Brown, R. J.; Coleman, P. J.; Conolly, C.; Sweetman, A. J.; Jones, K.
484 C.; Butterfield, D. M.; Sarantaridis, D.; Donovan, B. J.; Roberts, I., Twenty years of

485 measurement of polycyclic aromatic hydrocarbons (PAHs) in UK ambient air by nationwide
 486 air quality networks. *Environ Sci Process Impacts* **2013**, *15* (6), 1199-215.

487 9. Okona-Mensah, K. B.; Battershill, J.; Boobis, A.; Fielder, R., An approach to
 488 investigating the importance of high potency polycyclic aromatic hydrocarbons (PAHs) in the
 489 induction of lung cancer by air pollution. *Food Chem Toxicol* **2005**, *43* (7), 1103-16.

490 10. Bergvall, C.; Westerholm, R., Identification and determination of highly carcinogenic
 491 dibenzopyrene isomers in air particulate samples from a street canyon, a rooftop, and a
 492 subway station in Stockholm. *Environ Sci Technol* **2007**, *41* (3), 731-7.

493 11. Jia, Y.; Stone, D.; Wang, W.; Schrlau, J.; Tao, S.; Simonich, S. L., Estimated
 494 reduction in cancer risk due to PAH exposures if source control measures during the 2008
 495 Beijing Olympics were sustained. *Environ Health Perspect* **2011**, *119* (6), 815-20.

496 12. Layshock, J.; Simonich, S. M.; Anderson, K. A., Effect of dibenzopyrene
 497 measurement on assessing air quality in Beijing air and possible implications for human
 498 health. *J Environ Monit* **2010**, *12* (12), 2290-8.

499 13. Menichini, E.; Merli, F., Dibenzopyrenes, other PAHs with molecular weight 302,
 500 and selected carcinogenic PAHs seldom determined: identification and one-year
 501 quantification in urban air. *Intern J Environ Anal Chem* **2012**, *92* (14), 1609-1625.

502 14. Slezakova, K.; Pires, J. C.; Castro, D.; Alvim-Ferraz, M. C.; Delerue-Matos, C.;
 503 Morais, S.; Pereira, M. C., PAH air pollution at a Portuguese urban area: carcinogenic risks
 504 and sources identification. *Environ Sci Pollut Res Int* **2013**, *20* (6), 3932-45.

505 15. Mass, M. J.; Jeffers, A. J.; Ross, J. A.; Nelson, G.; Galati, A. J.; Stoner, G. D.;
 506 Nesnow, S., Ki-ras oncogene mutations in tumors and DNA adducts formed by
 507 benz[j]aceanthrylene and benzo[a]pyrene in the lungs of strain A/J mice. *Mol Carcinog* **1993**,
 508 *8* (3), 186-92.

- 509 16. Nesnow, S.; Gold, A.; Sangaiah, R.; Slaga, T. J., Mouse skin tumor-initiating activity
510 of benz[j]aceanthrylene in SENCAR mice. *Cancer Lett* **1993**, 73 (2-3), 73-6.
- 511 17. Holme, J. A.; Bjorge, C.; Soderlund, E. J.; Brunborg, G.; Becher, R.; Lag, M.;
512 Schwarze, P. E.; Ross, J.; Nelson, G.; Nesnow, S., Genotoxic effects of cyclopenta-fused
513 polycyclic aromatic hydrocarbons in isolated rat hepatocytes and rabbit lung cells.
514 *Carcinogenesis* **1993**, 14 (6), 1125-31.
- 515 18. Nesnow, S.; Lasley, J.; Curti, S.; Ross, J.; Nelson, G.; Sangaiah, R.; Gold, A.,
516 Morphological transformation and DNA adduct formation by benz[j]aceanthrylene and its
517 metabolites in C3H10T1/2CL8 cells: evidence for both cyclopenta-ring and bay-region
518 metabolic activation pathways. *Cancer Res* **1991**, 51 (22), 6163-9.
- 519 19. Johnsen, N. M.; Brunborg, G.; Haug, K.; Scholz, T.; Holme, J. A., Metabolism and
520 activation of cyclopenta polycyclic aromatic hydrocarbons in isolated human lymphocytes,
521 HL-60 cells and exposed rats. *Chem Biol Interact* **1998**, 114 (1-2), 77-95.
- 522 20. Johnsen, N. M.; Nyholm, S. H.; Haug, K.; Scholz, T.; Holme, J. A., Metabolism and
523 activation of cyclopenta polycyclic aromatic hydrocarbons in liver tissue from rats and
524 humans. *Chem Biol Interact* **1998**, 113 (3), 217-37.
- 525 21. Grimmer, G.; Jacob, J.; Dettbarn, G.; Naujack, K. W., Determination of polycyclic
526 aromatic hydrocarbons, azaarenes, and thiaarenes emitted from coal-fired residential furnaces
527 by gas chromatography/mass spectrometry. *Fresenius' Zeitschrift für analytische Chemie*
528 **1985**, 322 (6), 595-602.
- 529 22. Schmidt, W.; Grimmer, G.; Jacob, J.; Dettbarn, G., Polycyclic aromatic hydrocarbons
530 and thiaarenes in the emission from hard-coal combustion. *Toxicol Environ Chem* **1986**, 13
531 (1-2), 1-16.

- 532 23. Kamens, R. M.; Fulcher, J. N.; Ball, L. M.; Gold, A., Analysis of cyclopentafused
533 isomers of benz(a)anthracene in wood smoke. *Biomed Environ Mass Spectrom* **1987**, *14* (8),
534 369-74.
- 535 24. Gao, H.; Zhou, L.; Ma, M. Q.; Chen, X. G.; Hu, Z. D., Composition and source of
536 unknown organic pollutants in atmospheric particulates of the Xigu District, Lanzhou,
537 People's Republic of China. *Bull Environ Contam Toxicol* **2004**, *72* (5), 923-30.
- 538 25. Li, Z.; Wu, Y.; Zhao, Y.; Wang, L.; Zhu, H.; Qin, L.; Feng, F.; Wang, W., Analysis of
539 coal tar pitch and smoke extract components and their cytotoxicity on human bronchial
540 epithelial cells. *J Hazard Mater* **2011**, *186* (2-3), 1277-82.
- 541 26. Rodgman, A.; Smith, C. J.; Perfetti, T. A., The composition of cigarette smoke: a
542 retrospective, with emphasis on polycyclic components. *Hum Exp Toxicol* **2000**, *19* (10),
543 573-95.
- 544 27. Suzuki, S.; Kaneko, T.; Tsuchiya, M., Characterization of C12 to C45 polycyclic
545 aromatic hydrocarbons and fused-ring heterocyclic compounds in coal tar pitch and airborne
546 total suspended particle. *J Environ Chem* **1996**, *6* (4), 511-520.
- 547 28. Wang, Q.; Kobayashi, K.; Zhou, M.; Lu, S.; Dong, S.; Nakajima, D.; Sekigushi, K.;
548 Terasaki, M., Study on the size-segregated distribution of 37 species of polycyclic aromatic
549 hydrocarbons in urban atmospheric fine particles of Japan. In *Air Pollution XXII*, Longhurst,
550 J.; Brebbia, C., Eds. WIT Press: 2014; pp 131-143.
- 551 29. Bergvall, C.; Westerholm, R., Determination of 252-302 Da and tentative
552 identification of 316-376 Da polycyclic aromatic hydrocarbons in Standard Reference
553 Materials 1649a Urban Dust and 1650b and 2975 Diesel Particulate Matter by accelerated
554 solvent extraction-HPLC-GC-MS. *Anal Bioanal Chem* **2008**, *391* (6), 2235-48.
- 555 30. Sadiktsis, I.; Koegler, J. H.; Benham, T.; Bergvall, C.; Westerholm, R., Particulate
556 associated polycyclic aromatic hydrocarbon exhaust emissions from a portable power

generator fueled with three different fuels – A comparison between petroleum diesel and two biodiesels. *Fuel* **2014**, *115* (0), 573-580.

31. Knasmüller, S.; Parzefall, W.; Sanyal, R.; Ecker, S.; Schwab, C.; Uhl, M.; Mersch-Sundermann, V.; Williamson, G.; Hietsch, G.; Langer, T., et al., Use of metabolically competent human hepatoma cells for the detection of mutagens and antimutagens. *Mutat Res* **1998**, *402* (1-2), 185-202.

32. Jarvis, I. W. H.; Bergvall, C.; Bottai, M.; Westerholm, R.; Stenius, U.; Dreij, K., Persistent activation of DNA damage signaling in response to complex mixtures of PAHs in air particulate matter. *Toxicol Appl Pharmacol* **2013**, *266* (3), 408-418.

33. Niziolek-Kierecka, M.; Dreij, K.; Lundstedt, S.; Stenius, U., γ H2AX, pChk1, and Wip1 as potential markers of persistent DNA damage derived from dibenzo[a,l]pyrene and PAH-containing extracts from contaminated Soils. *Chem Res Toxicol* **2012**, *25* (4), 862-72.

34. Tarantini, A.; Maitre, A.; Lefebvre, E.; Marques, M.; Marie, C.; Ravanat, J. L.; Douki, T., Relative contribution of DNA strand breaks and DNA adducts to the genotoxicity of benzo[a]pyrene as a pure compound and in complex mixtures. *Mutat Res* **2009**, *671* (1-2), 67-75.

35. Mosmann, T., Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* **1983**, *65* (1-2), 55-63.

36. Karlsson, H. L.; Nilsson, L.; Moller, L., Subway particles are more genotoxic than street particles and induce oxidative stress in cultured human lung cells. *Chem Res Toxicol* **2005**, *18* (1), 19-23.

37. Carpenter, A. E.; Jones, T. R.; Lamprecht, M. R.; Clarke, C.; Kang, I. H.; Friman, O.; Guertin, D. A.; Chang, J. H.; Lindquist, R. A.; Moffat, J., et al., CellProfiler: image analysis software for identifying and quantifying cell phenotypes. *Genome Biol* **2006**, *7* (10), R100.

- 581 38. Huber, P. J., The behavior of maximum likelihood estimates under nonstandard
582 conditions. *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and*
583 *Probability, Volume 1: Statistics*, 221-233, University of California Press, Berkeley, CA, USA
584 **1967**.
- 585 39. Minnesota Department of Health., Guidance for evaluating the cancer potency of
586 polycyclic aromatic hydrocarbon (PAH) mixtures in environmental samples. 2014 October
587 31.
- 588 40. Nesnow, S.; Leavitt, S.; Easterling, R.; Watts, R.; Toney, S. H.; Claxton, L.;
589 Sangaiah, R.; Toney, G. E.; Wiley, J.; Fraher, P., et al., Mutagenicity of cyclopenta-fused
590 isomers of benz(a)anthracene in bacterial and rodent cells and identification of the major rat
591 liver microsomal metabolites. *Cancer Res* **1984**, *44* (11), 4993-5003.
- 592 41. Sangaiah, R.; Gold, A.; Toney, G. E.; Toney, S. H.; Claxton, L.; Easterling, R.;
593 Nesnow, S., Benz[j]aceanthrylene: a novel polycyclic aromatic hydrocarbon with bacterial
594 mutagenic activity. *Mutat Res* **1983**, *119* (3), 259-66.
- 595 42. Umbuzeiro, G. A.; Kummrow, F.; Morales, D. A.; Alves, D. K.; Lim, H.; Jarvis, I. W.
596 H.; Bergvall, C.; Westerholm, R.; Stenius, U.; Dreij, K., Sensitivity of salmonella YG5161
597 for detecting PAH-associated mutagenicity in air particulate matter. *Environ Mol Mutagen*
598 **2014**, *55* (6), 510-7.
- 599 43. Solhaug, A.; Refsnes, M.; Lag, M.; Schwarze, P. E.; Husoy, T.; Holme, J. A.,
600 Polycyclic aromatic hydrocarbons induce both apoptotic and anti-apoptotic signals in
601 Hepa1c1c7 cells. *Carcinogenesis* **2004**, *25* (5), 809-19.
- 602 44. Buterin, T.; Hess, M. T.; Luneva, N.; Geacintov, N. E.; Amin, S.; Kroth, H.; Seidel,
603 A.; Naegeli, H., Unrepaired fjord region polycyclic aromatic hydrocarbon-DNA adducts in
604 ras codon 61 mutational hot spots. *Cancer Res* **2000**, *60* (7), 1849-56.

- 605 45. Dreij, K.; Seidel, A.; Jernström, B., Differential removal of DNA adducts derived
606 from anti-diol epoxides of dibenzo[a,l]pyrene and benzo[a]pyrene in human cells. *Chem Res*
607 *Toxicol* **2005**, 18 (4), 655-64.
- 608 46. Geacintov, N. E.; Broyde, S.; Buterin, T.; Naegeli, H.; Wu, M.; Yan, S.; Patel, D. J.,
609 Thermodynamic and structural factors in the removal of bulky DNA adducts by the
610 nucleotide excision repair machinery. *Biopolymers* **2002**, 65 (3), 202-10.
- 611 47. Jarvis, I. W. H.; Bergvall, C.; Morales, D. A.; Kummrow, F.; Umbuzeiro, G. A.;
612 Westerholm, R.; Stenius, U.; Dreij, K., Nanomolar levels of PAHs in extracts from urban air
613 induce MAPK signaling in HepG2cells. *Toxicol Lett* **2014**, 229 (1), 25-32.
- 614 48. Westerholm, R.; Bergvall, C.; Sadiktsis, I.; Johansson, C.; Stenius, U., Mätning av
615 starkt carcinogena dibensopyrener i jämförelse med humancarcinogenen bens(a)pyren
616 [B(a)P] i Stockholmsluft från vägtrafik. *Report (in Swedish) to the Stockholm county council.*
617 *Stockholm (2012), p. 1* **2012**.
- 618 49. Ravindra, K.; Sokhi, R.; Van Grieken, R., Atmospheric polycyclic aromatic
619 hydrocarbons: Source attribution, emission factors and regulation. *Atmospheric Environment*
620 **2008**, 42 (13), 2895-2921.
- 621 50. Mattsson, A.; Jernström, B.; Cotgreave, I. A.; Bajak, E., H2AX phosphorylation in
622 A549 cells induced by the bulky and stable DNA adducts of benzo[a]pyrene and
623 dibenzo[a,l]pyrene diol epoxides. *Chem Biol Interact* **2009**, 177 (1), 40-7.
- 624 51. Rogakou, E. P.; Boon, C.; Redon, C.; Bonner, W. M., Megabase chromatin domains
625 involved in DNA double-strand breaks in vivo. *J Cell Biol* **1999**, 146 (5), 905-16.
- 626 52. Audebert, M.; Zeman, F.; Beaudoin, R.; Pery, A.; Cravedi, J. P., Comparative potency
627 approach based on H2AX assay for estimating the genotoxicity of polycyclic aromatic
628 hydrocarbons. *Toxicol Appl Pharmacol* **2012**, 260 (1), 58-64.

- 629 53. Bartos, T.; Cupr, P.; Klanova, J.; Holoubek, I., Which compounds contribute most to
630 elevated airborne exposure and corresponding health risks in the Western Balkans? *Environ*
631 *Int* **2009**, 35 (7), 1066-71.
- 632 54. Amarillo, A. C.; Tavera Busso, I.; Carreras, H., Exposure to polycyclic aromatic
633 hydrocarbons in urban environments: Health risk assessment by age groups. *Environ Pollut*
634 **2014**, 195C, 157-162.
- 635 55. Wang, J.; Chen, S.; Tian, M.; Zheng, X.; Gonzales, L.; Ohura, T.; Mai, B.; Simonich,
636 S. L., Inhalation cancer risk associated with exposure to complex polycyclic aromatic
637 hydrocarbon mixtures in an electronic waste and urban area in South China. *Environ Sci*
638 *Technol* **2012**, 46 (17), 9745-52.
- 639
- 640

Fig. 1. Detection of B[j]A in air PM samples. GC/MS chromatograms obtained in SIM mode of air PM samples from Stockholm (A), Limeira (B), Stockholm spiked with B[j]A (C) and Limeira spiked with B[j]A (D). 1: benzo[b]fluoranthene, 2: benzo[k]fluoranthene, 3: benzo[j]fluoranthene, 4: benzo[a]fluoranthene, 5: unknown, 6: benz[j]aceanthrylene, 7: benzo[e]pyrene, 8: benzo[a]pyrene-D₁₂, 9: benzo[a]pyrene, 10: perylene.

Fig. 2. B[j]A reduce cell viability and induce DNA damage stronger than B[a]P. HepG2 cells were exposed to B[j]A, B[a]P or control solvent (DMSO) for 48 h. MTT assay (A) of cells exposed to 0.001-1 μM B[j]A or 0.01-4 μM B[a]P where cell viability is presented as percent of DMSO treated cells. Comet assay (B) was performed on cells exposed to 0.3 and 1 μM B[j]A or B[a]P, or DMSO and 100 cells were scored (n=2). Sigmoidal dose response curve fit (n=6) was applied in A. One-way ANOVAs with Bonferroni's post test was applied in A and B, * $p < 0.05$ compared DMSO. Immunostaining (C) of γH2AX (red) and DAPI (blue) in cells exposed to 0.3 μM B[a]P or B[j]A, or DMSO. Scatter plot (D) of counted cells and foci from immunostaining with statistical significance obtained using a linear regression model.

Fig. 3. Stronger activation of DNA damage signaling proteins in cells exposed to B[j]A compared to B[a]P. HepG2 cells were exposed to PAHs for 48 h and protein levels were measured with Western blot. In (A) the effect of equimolar concentration of B[j]A and B[a]P (1 μ M) on pChk1, γ H2AX, p53, pp53 and p21. Cdk2 was used as loading control. In (B) dose response of 0.001-1 μ M B[j]A and 0.1-5 μ M B[a]P on pChk1 and γ H2AX. In (C) densitometric analysis of pChk1 and γ H2AX. Densitometric analysis of DBC was previously published in Jarvis et al., 2013. Non-linear curve fit, sigmoidal dose response, n=4.

709 **Table 1.** Information of the air PM samples collected in Stockholm and Limeira.

Name	Abbreviation	Sampling period	Duration (hours)	Sampled air (m ³)	Total PAH (pg/m ³)	B[a]P (pg/m ³ , n=3)	B[j]A (pg/m ³ , n=3)	Total PM (mg)
Stockholm 1	STO1	Jan 18-21, 2013	71	5141	4440 ± 138	289 ± 4	12.7 ± 0.2	36.8
Stockholm 2	STO2	Sep 10-17, 2013	168	12265	351 ± 4.1	17.3 (SD<0.02)	1.57 ± 0.07	91.3
Stockholm 3	STO3	Dec 10-17, 2013	167	11034	793 ± 7.1	49.5 ± 0.5	3.14 ± 0.15	60.3
Limeira 1	LMR1	Jul 12, 2010	24	2352	7958 ± 158	560 ± 10	19.6 ± 0.8	263
Limeira 2	LMR2	Jul 19, 2010	24	2441	10693 ± 297	899 ± 17	30.2 ± 0.7	234

711 **Table 2.** Contribution of B[j]A levels to the excess lifetime cancer cases from exposure to air
712 PM. Based on B[a]P_{eq} levels and the WHO unit risk value of 8.7×10^{-5} ng/m³ B[a]P.⁴

Air PM	Excess lifetime cancer cases / 100 000 people				
	B[a]P	B[a]P _{eq} excl. B[j]A	B[a]P _{eq} incl. B[j]A ^a		
			RPF of B[j]A = 10 ^b	RPF of B[j]A = 30 ^c	RPF of B[j]A = 60 ^d
STO1	2.5	11.3	+ 1.1	+ 3.4	+ 6.7
STO2	0.15	0.77	+ 0.14	+ 0.41	+ 0.82
STO3	0.43	2.3	+ 0.3	+ 0.9	+ 1.7
LMR1	4.9	16.2	+ 1.7	+ 5.1	+ 10.2
LMR2	7.8	22.7	+ 2.6	+ 7.9	+ 15.7

713 ^aincrease in cancer cases compared to B[a]P_{eq} excl. B[j]A; ^b from^{16,40-41}, ^c our fold induction data for γH2AX
714 (Table S2), ^d from¹⁵

715

716

717 **Abstract Art**

718