**Karolinska
Institutet**

Karolinska Institutet

<http://openarchive.ki.se>

This is a Peer Reviewed Accepted version of the following article, accepted for publication in Environmental and molecular mutagenesis.

2017-07-07

Sensitivity of Salmonella YG5161 for detecting PAH-associated mutagenicity in air particulate matter

Umbuzeiro, Gisela A; Kummrow, Fabio; Morales, Daniel A; Alves, Debora Kristina M; Lim, Hwanmi; Jarvis, Ian W H; Bergvall, Christoffer; Westerholm, Roger; Stenius, Ulla; Dreij, Kristian

Environ Mol Mutagen. 2014 Jul;55(6):510-7.

<http://doi.org/10.1002/em.21861>

<http://hdl.handle.net/10616/45988>

If not otherwise stated by the Publisher's Terms and conditions, the manuscript is deposited under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

**Karolinska
Institutet**

This is the peer reviewed version of the following article
Environ Mol Mutagen. 2014 Jul;55(6):510-7. which has
been published in final form at:

<http://dx.doi.org/10.1002/em.21861>

**Sensitivity of Salmonella YG5161 for detecting PAH-
associated mutagenicity in air particulate matter.**

**Umbuzeiro, GA; Kummrow, F; Morales, D.A.; Alves, DK;
Lim, H; Jarvis, Ian WH; Bergvall, Christoffer;
Westerholm, Roger; Stenius, Ulla; Dreij, K**

Access to the published version may require subscription.
Published with permission from: **Wiley**

**Sensitivity of Salmonella YG5161 for Detecting PAH-Associated Mutagenicity in Air
Particulate Matter**

Running title: Sensitivity of YG5161 for detection of PAH-mutagenicity in air

Gisela A. Umbuzeiro^{1*}, Fábio Kummrow², Daniel Alexandre Morales¹, Debora Kristina M. Alves³, Hwanmi Lim⁵, Ian W. H. Jarvis⁴, Christoffer Bergvall⁵, Roger Westerholm⁵, Ulla Stenius⁴, and Kristian Dreij⁴

¹Faculty of Technology, State University of Campinas (Unicamp), Limeira, SP, Brazil.

²Institute of Environmental, Chemical and Pharmaceutical Sciences, Federal University of São Paulo (Unifesp), Diadema, SP, Brazil.

³Pharmaceutical Sciences Faculty, University of São Paulo (USP), São Paulo, SP, Brazil.

⁴Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden.

⁵Department of Analytical Chemistry, Stockholm University, Stockholm, Sweden.

Keywords: PAHs, Salmonella/microsome; nitro-PAHs, mutagenicity; air pollution

* Corresponding author: Tel: +55 19 21133419; E-mail: giselau@ft.unicamp.br

ABSTRACT

The Salmonella/microsome assay is the most used assay for the evaluation of air particulate matter (PM) mutagenicity and a positive correlation between strain TA98 responses and benzo[a]pyrene (B[a]P) levels in PM has been found. However, it seems that the major causes of PM mutagenicity in this assay are the nitro and oxy-PAHs. Salmonella YG5161, a 30-times more responsive strain to B[a]P has been developed. To verify if YG5161 strain was sufficiently sensitive to detect mutagenicity associated with B[a]P mutagenicity, PM samples were collected in Brazil and Sweden, extracted with toluene and tested in the Salmonella/microsome microsuspension assay. PAHs and B[a]P were determined and the extracts were tested with YG5161 and its parental strain TA1538. The extracts were also tested with YG1041 and its parental strain TA98. For sensitivity comparisons, we tested B[a]P and 1-nitropyrene (1-NP) using the same conditions. The minimal effective dose of B[a]P was 155 ng/plate for TA1538 and 7 ng/plate for YG5161. Although the maximum tested dose, 10 m3/plate containing 9 ng of B[a]P in the case of Brazilian sample, was sufficient to elicit a response in YG5161, mutagenicity was detected at a dose as low as 1 m3/plate (0.9 ng). This is probably caused by nitro-compounds that have been shown to be even more potent than B[a]P for YG5161. It seems that the mutagenicity of B[a]P present in PM is not detectable even with the use of YG5161 unless more efficient separation to remove the nitro-compounds from the PAH extract is performed.

1 INTRODUCTION

2 The mutagenicity of airborne particulate matter (PM) can be attributed to at least 500
3 identified compounds from different chemical classes [Claxton et al., 2004]. Among these,
4 benzo[a]pyrene (B[a]P), along with other polycyclic aromatic hydrocarbons (PAHs), have
5 received great attention because of their recognized carcinogenic and mutagenic potential
6 [Srogi, 2007].

7 The Salmonella/microsome assay is the most widely used method for the evaluation of
8 the mutagenic activity of pure compounds and environmental samples [Claxton et al., 2010],
9 including atmospheric samples [Claxton and Woodall Jr, 2007]. The assay is sensitive to
10 several PAHs, including B[a]P [DeMarini et al., 2011; Brito et al., 2013]. Some studies have
11 correlated the mutagenic activity detected in the Salmonella/microsome assay with the levels
12 of B[a]P and other non-substituted PAHs present in the samples [Viras et al., 1990; Nielsen et
13 al., 1996; Claxton and Woodall Jr., 2007; Srogi, 2007], although it does not seem to
14 demonstrate a direct relationship. The primary components responsible for the mutagenicity
15 of air particulate matter in the Salmonella assay seem to be nitro and oxy-PAHs [Claxton et
16 al., 2004; Sharma et al., 2007; Umbuzeiro et al., 2008 a, b; Walgraeve et al., 2010]. One
17 explanation for this could be that the typically used strains (TA98 and TA100) are more
18 sensitive to nitro- and oxy- PAHs than to non-substituted PAHs [Claxton et al., 2004; Enya et
19 al., 1997; Enya et al., 1998; Kummrow et al., 2006; Franco et al., 2010]. To enhance the
20 sensitivity of the Salmonella/microsome assay to non-substituted PAHs Matsui et al. [2006]
21 developed the YG5161 strain which is more responsive to B[a]P and other non-substituted
22 PAHs than its parental strain TA1538. The YG5161 strain overexpresses DNA polymerase
23 IV, and has the *dinB* gene of *Escherichia coli* encoded in the pYG768 plasmid, which also
24 confers ampicillin resistance to the strain. DNA polymerase IV facilitates the error-prone
25 bypass of the DNA guanine adducts formed by polycyclic aromatic compounds which, after

1 repair, will lead to the deletion of two base pairs and consequently shifting of the reading
2 frame [Matsui et al., 2006]. These authors suggested the possibility of using YG5161 as a
3 major strain for the detection of the mutagenicity of non-substituted PAHs such as B[a]P.
4 Because this compound needs to be metabolized to react with DNA [Uppstad et al., 2010], the
5 addition of S9 mix is required for its detection in the Salmonella/mutagenicity assay.

6 Some strains have also been developed to be more sensitive to different compounds.
7 For example YG1041 strain, a derivative of TA98, is more sensitive to nitroarenes and
8 aromatic amines because it overproduces nitroreductase and *O*-acetyltransferases, both
9 important enzymes in the activation of such compounds [Hagiwara et al., 1993]. Similarly,
10 the strain YG7108 which is derived from TA1535 is more responsive to alkylating agents
11 [Yamada et al., 1997]. Both strains have been used in the identification of the types of
12 compounds predominantly responsible for the mutagenic activity of a test samples
13 [Umbuzeiro et al., 2011]. Mutlu et al. [2013] demonstrated the applicability of strains with
14 different sensitivities for analyzing environmental samples. In a hierarchical clustering
15 analysis they showed that although PAHs, aromatic amines, and nitro-compounds were
16 present in diesel exhaust extracts, oxy-PAHs were the cause of much of the mutagenicity.

17 When atmospheric particulate air samples are tested they usually demonstrate a clear
18 increase with YG1041 in relation to TA98, indicating that the mutagenicity is related to nitro-
19 and oxy- PAHs and not B[a]P, although PAHs are present in those samples when chemical
20 analyses are performed [DeMarini et al., 2004; Umbuzeiro et al., 2008a,b].

21 The objective of this work was to verify if the YG5161 strain in the
22 Salmonella/microsome microsuspension assay was sufficiently sensitive to B[a]P
23 mutagenicity in air particulate matter collected in two different locations, Limeira, Brazil and
24 Stockholm, Sweden. We determined the total PAH and B[a]P levels and tested the same
25 extracts with YG5161 and its parental strain, TA1538. We also tested the same extracts with

YG1041 and its parental strain, TA98. For sensitivity comparisons we tested several concentrations of B[a]P and 1-nitropyrene (1-NP) using the same conditions.

MATERIAL AND METHODS

Sampling sites

Total atmospheric particulate matter (PM) was collected at two sites: the campus of the Faculty Technology at UNICAMP, Limeira, Brazil (LIMEIRA) and the campus of the Stockholm University, Stockholm, Sweden (STHOLM). The LIMEIRA site is impacted by heavy traffic including cars and trucks, industrial emissions, and sugar cane growing and harvesting activities, including biomass burning. The mutagenic potencies of previously evaluated extracts from this site using the Salmonella/microsome assay [Alves, 2011] and surrounding areas [Umbuzeiro et al., 2008a, b] were among the highest potencies found in the literature. The STHOLM site is also impacted by heavy traffic with emissions from cars and trucks as the main source of pollution. PM samples from this site have recently been used to investigate the toxicological impact of PAHs in air PM [Jarvis et al., 2013a,b].

Total PM samples from Limeira were collected at street level on a glass-fiber filter (254 × 233 mm; 0.33 mm pore size. Energética Ind. Com. LTDA, Rio de Janeiro, RJ, Brazil) using a high-volume sampler (Energética Ind. Com. LTDA, Rio de Janeiro, RJ, Brazil) operated at an average flow rate of 1,130 L/min for 24 h. The air total PM samples from Stockholm, Sweden were collected at roof-top level (22 m from the ground) on a fluorocarbon-coated glass fiber filter (235 mm of diameter; Fiberfilm Filters, Pallflex, Pall Corporation, Putnam, CT, USA) with an average flow rate of 1,209 L/min for 71 h. The total PM concentrations for Limeira were 95.8 ug/m³ and for Stockholm, 7.16 ug/m³.

Organic extraction and PAH analysis

Extraction was performed by pressurized fluid extraction using an ASE 200 accelerated solvent extraction system (Dionex Co., Sunnyvale, CA, USA) using toluene for 5 x 30 min. This extraction procedure was previously developed and validated for analysis of PAHs in air PM [Bergvall and Westerholm, 2008]. Aliquots of the extracts were evaporated gently under a nitrogen stream at 55°C until dryness and re-dissolved in dimethylsulfoxide (DMSO) for the biological assays. Portions of the extracts were analyzed for 42 PAHs (Table I) using silica solid phase extraction (SPE) cartridges and on-line liquid chromatography-gas chromatography/mass spectrometry as described in detail elsewhere [Sadiktsis et al., 2014]. A blank filter was also extracted and analyzed as an analytical control.

Salmonella/microsome microsuspension assay

The extracts were assayed in the Salmonella/microsome microsuspension assay [Kado et al., 1983; DeMarini et al., 1989] in dose-response experiments using the *Salmonella* strains TA1538, TA98, YG5161 and YG1041 with and without metabolic activation (S9). Strains were kindly provided by Dr. Takehiko Nohmi, except for TA98, which was kindly provided by Dr. Larry Claxton. Table II summarizes the genetic characteristics of the strains. B[a]P and 1-NP were also tested with the four strains in dose response experiments. Overnight cultures (around 10^9 cells/mL) were concentrated 5-fold by centrifugation (10,000xg at 4 °C for 10 min) and resuspended into 0.015 M sodium phosphate buffer. A volume of 50 µL of cell suspension, 50 µL of 0.015 M sodium phosphate buffer or S9 mix, and 5 µL of the sample were incubated at 37 °C for 90 min without shaking. To the mixture, 2 mL of molten agar was added and poured onto a minimal agar plate. Colonies were counted after 66 h of incubation at 37 °C by hand with the aid of a stereomicroscope. Toxicity was also carefully evaluated by observing the background of the agar plates. Metabolic activation was provided by Aroclor

1254-induced Sprague Dawley rat liver S9 mix (MolTox, Boone, NC) prepared at 4 % v/v and supplemented with the required co-factors [Mortelmans and Zeiger, 2000].

PM extracts were tested at 0.02, 0.2, 1, 2, 5 and 10 m³ per plate. Positive controls without S9 were 4-nitroquinoline-oxide (4NQO) at 0.125 µg/plate for TA98 and YG5161 and 4-nitro-o-phenylenediamine (4NOP) at 2.5 µg/plate for YG1041. With S9, 2-aminoanthracene (2AA) at 0.625 µg/plate was used as a positive control for TA1538, TA98 and YG5161 and at 0.03125 µg/plate for YG1041. Duplicates of each concentration were tested, except for the negative control that was tested in triplicate. DMSO was used as negative control and to dilute the extracts and positive controls. The extract of a clean (blank) filter was also tested.

Data were analyzed with the Salanal computer program using the Bernstein model [Bernstein et al., 1982]. Samples were considered positive when a significant difference among the tested doses and the negative control (ANOVA) and a significant positive dose response were observed. Results were expressed as revertants per m³ and per mg of PM equivalent. Also, the minimum effective dose (MED) to elicit a positive response was calculated for B[a]P and 1-NP. MED was defined as the dose that provided a doubling of revertants based on the calculated linear regression curve.

RESULTS AND DISCUSSION

In this study, we evaluated the mutagenicity of PM organic extracts from two cities: Limera, Brazil and Stockholm, Sweden. The concentration of PM collected in LIMEIRA was 95.8 µg/m³ and in STHOLM was 7.16 µg/m³. The total PAH content in the LIMEIRA sample was 10.66 ng/m³ with 0.9 ng/m³ of B[a]P, and the STHOLM sample contained 4.43 ng/m³ with 0.29 ng/m³ of B[a]P (Table I). The mean of the number of revertants per plate obtained for the four strains including negative, blank, positive controls and calculated potencies are presented in Table III for LIMEIRA and Table IV for STHOLM. In LIMEIRA,

the potency for TA98 without S9 was 100 revertants/m³, similar what had been reported for other cities in Brazil [Sato et al., 1995; Ducatti and Vargas, 2003; Umbuzeiro et al., 2008 a, b]. The potency of the STHOLM sample was 4 times lower (25 revertants/m³). The potency values decreased for both locations when S9 was used (Tables III and IV). According to Claxton et al. [2004] this is a first indication of a response of nitroaromatics. In general the samples from LIMEIRA were more mutagenic than the STHOLM in all strains/conditions tested, especially with TA98 with S9 (13-fold difference). However, because only one sample from each site was analyzed it was not possible to compare the mutagenicity of the two sites with a high level of confidence. A comprehensive study, which includes additional sampling at the same season of the year, is being conducted to provide additional information on the potencies at both sites.

To establish how sensitive each strain was to B[a]P, the prototypical non-substituted PAH, and 1-NP, a typical nitroaromatic, their mutagenicities were assessed using the same protocol used to evaluate the PM samples.

YG5161 was the most sensitive strain for B[a]P showing a MED of 7 ng/plate in contrast to 77.5 ng/plate for TA1538, 265 ng/plate for YG1041, and 1,400 ng/plate for TA98 (Table V). Although several non-substituted PAHs were detected in both PM samples (Table II), the potency of the YG5161 response did not provide a typical B[a]P response, which would be an increase of 30-fold in relation to TA1538 (Table V). This is most likely due to the fact that the amount of B[a]P present in the 10 m³ of PM (the maximum dose tested) was 9 ng/plate and the MED of B[a]P for YG5161 is 7 ng/plate (Table V). All the comparisons were made with S9 because non-substituted PAHs require S9 to be activated.

YG1041, as expected, demonstrated much higher sensitivity to 1-NP than any other strain, with a MED of 0.03 ng/plate. MEDs for TA98, TA1538 and YG5161 were 0.45, 1.8 and 1.8 ng/plate, respectively (Table VI). 1-NP was chosen as an example for testing purpose,

1 but other nitrocompounds (e.g. dinitropyrenes, nitrobenzanthrones) are more mutagenic and
2 are likely to be present in PM samples. For both PM samples we observed a typical response
3 of nitroaromatics because the potency of YG1041 was 35 times higher than TA98. This
4 comparison was made without S9 because all the potencies decreased with S9. This behavior
5 is consistent with the hypothesis that nitroaromatics are the major cause of the mutagenic
6 effect.

7 The data suggest that B[a]P is not the compound predominantly causing the mutagenic
8 responses of the PM analyzed in this study, but that nitroaromatics such as 1-NP seems to be
9 causing the observed effect. Chemical analysis of nitroaromatics could be performed in the
10 same extracts in order to determine their contribution to the mutagenicity.

11 Other authors also used YG5161 to test different samples. Sharma et al. [2007] used
12 TA98, YG1041, and YG5161 strains, in the Salmonella/microsome microsuspension assay to
13 evaluate the genotoxicity of organic fractions of PM collected from an incineration energy
14 plant and urban air. The fraction containing moderately polar PAHs, alkyl-PAHs and *O*- and
15 *S*-heterocyclics was mutagenic only in the presence of S9 mix, indicating that compounds
16 such as B[a]P could be responsible for the observed effect. Because the mutagenic activity
17 with the YG5161 strain was lower in comparison to TA98 it is plausible to suspect that non-
18 substituted PAHs are not predominantly contributing to the observed mutagenic activity.
19 Conversely, the fraction containing N-heterocyclics, nitro-, amino- and oxy-PAHs was more
20 mutagenic without S9 and higher in YG1041 than TA98, suggesting that nitroarenes and
21 aromatic amines are likely to be the major cause of the mutagenic activity.

22 Maertens et al. [2009] used the Salmonella/microsome assay with several
23 strains/conditions, including TA98, YG1041 and YG5161 to, evaluate the mutagenicity of
24 tobacco and marijuana smoke condensates. All of the tested smoke condensates presented
25 positive responses. The mutagenic potencies obtained with YG1041 and with YG5161 were

higher than the ones with TA98 indicating that aromatic amines and non-substituted PAHs were contributing to the mutagenicity. Although the authors identified B[a]P and benzo[a]anthracene in the samples, they contributed to less than 0.1% of the total mutagenicity indicating that there are other major mutagens in the mixture.

Yauk et al. [2012] observed mutagenic activity in cigarette smoke condensate extracts using the Salmonella/microsome assay with the TA98, YG1041, and YG5161 strains in the presence of S9 mix. The authors observed that the potencies with YG1041 were greater than those for TA98 but the increase of YG5161 potencies in relation to TA98 was not significant, indicating that non-substituted PAHs could not fully explain the mutagenicity.

Therefore we conclude that even the highly sensitive strain to B[a]P, YG5161, does not seem to be able to distinguish the response of B[a]P in samples containing nitroaromatics. We observe that only 1 m³ is sufficient to elicit a positive response in the Salmonella/microsuspension assay for the majority of the strains/conditions. Both for the LIMEIRA and STHOLM air samples.

Therefore we can suggest that although associations were observed between B[a]P content and mutagenicity, this compound does not totally explain the mutagenicity of PM air samples tested in the Salmonella/microsome assay unless amounts of B[a]P are higher than 7 ng/plate and nitrocompounds are eliminated by the fractionation procedure.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank CAPES for the master fellowship of DKM Alves and DA Morales. We also thank Dr. Larry D. Claxton and Dr. Errol Zeiger for the valuable suggestions. This work was

supported by the Swedish Research Council Formas; Cancer- och Allergifonden and Stockholm University.

REFERENCES

- Alves DKM. 2011. A aplicabilidade de combinações seletivas de linhagens *S. typhimurium* na caracterização da mutagenicidade de amostras de ar. São Paulo: Departamento de Toxicologia e Análises Toxicológicas, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo. pp. 1-166.
- Bergvall C, Westerholm R. 2008. Determination of 252–302 Da and tentative identification of 316–376 Da polycyclic aromatic hydrocarbons in Standard Reference Materials 1649a Urban Dust and 1650b and 2975 Diesel Particulate Matter by accelerated solvent extraction-HPLC-GC-MS. *Anal Bioanal Chem* 391:2235-2248.
- Bernstein L, Kaldor J, McCann J, Pike MC. 1982. An empirical approach to the statistical analysis of mutagenesis data from the *Salmonella* test. *Mutat Res* 97:267-287.
- Brito KCT, Lemos CT, Rocha JAV, Mielli AC, Matzenbacher C, Vargas VMF. 2013. Comparative genotoxicity of airborne particulate matter (PM_{2.5}) using *Salmonella*, plants and mammalian cells. *Ecotoxicol Environ Saf* 94:14-20.
- Claxton LD, Matthews PP, Warren SH. 2004. The genotoxicity of ambient outdoor air, a review: *Salmonella* mutagenicity. *Mutat Res* 567:347-399.
- Claxton LD, Umbuzeiro GA, DeMarini DM. 2010. The *Salmonella* mutagenicity assay: the stethoscope of genetic toxicology for the 21st Century. *Environ Health Perspect* 118:1515-1522.
- Claxton LD, Woodall Jr GM. 2007. A review of the mutagenicity and rodent carcinogenicity of ambient air. *Mutat Res* 636:36-94.

- 1 DeMarini DM, Brooks LR, Warren SH, Kobayashi T, Gilmour MI, Singh P. 2004. Bioassay-
2 directed fractionation and Salmonella mutagenicity of automobile and forklift diesel
3 exhaust particles. Environ Health Perspect 112:814-819.
- 4 DeMarini DM, Dallas MM, Lewtas J. 1989. Cytotoxicity and effect on mutagenicity of
5 buffers in a microsuspension assay. Teratog Carcinog Mutagen 9:287-295.
- 6 DeMarini DM, Hanley DM, Warren SH, Adams LD, King LC. 2011. Association between
7 mutation spectra and stable and unstable DNA adduct profiles in *Salmonella* for
8 benzo[*a*]pyrene and dibenzo[*a,l*]pyrene. Mutat Res 714:17-25.
- 9 Ducatti A, Vargas VMF. 2003. Mutagenic activity of airborne particulate matter as an
10 indicative measure of air pollution. Mutat Res 540:67-77.
- 11 Enya T, Kawanishi M, Suzuki H, Matsui S, Hisamatsu Y. 1998. An unusual DNA adduct
12 derived from the powerfully mutagenic environmental contaminant 3-
13 nitrobenzanthrone. Chem Res Toxicol 11:1460-1467.
- 14 Enya T, Suzuki H, Watanabe T, Hirayama T, Hisamatsu Y. 1997. 3-Nitrobenzanthrone, a
15 powerful bacterial mutagen and suspected human carcinogen found in diesel exhaust
16 and airborne particulates. Environ Sci Technol 31:2772-2776.
- 17 Franco A, Kummrow F, Umbuzeiro GA, Vasconcellos PC, Carvalho LRF. 2010. Occurrence
18 of polycyclic aromatic hydrocarbons derivatives and mutagenicity study in extracts of
19 PM10 collected in São Paulo, Brazil. Rev Bras Toxicol 23:1-10.
- 20 Hagiwara Y, Watanabe M, Oda Y, Sofuni T, Nohmi T. 1993. Specificity and sensitivity of
21 *Salmonella typhimurium* YG1041 and YG1042 strains possessing elevated levels of
22 both nitroreductase and acetyltransferase activity. Mutat Res 291:171-180.
- 23 Jarvis IWH, Bergvall C, Bottai M, Westerholm R, Stenius U, Dreij K. 2013a. Persistent
24 activation of DNA damage signaling in response to complex mixtures of PAHs in air
25 particulate matter. Toxicol Appl Pharmacol 266:408-418.

- Jarvis IWH, Bergvall C, Morales DA, Kummrow F, Umbuzeiro GA, Westerholm R, Stenius U, Dreij K. 2013b. Complex mixtures of PAHs in air particulate matter stimulate an inflammatory response involving the MEK4/JNK/AP-1 pathway. *Toxicol Appl Pharmacol*, under revision.
- Kado NY, Langley D, Eisenstatd E. 1983. A simple modification of the *Salmonella* liquid incubation assay. *Mutat Res* 121:25-32.
- Kummrow F, Rech CM, Coimbra CA, Umbuzeiro GA. 2006. Blue rayon-anchored technique/*Salmonella* microsome microsuspension assay as a tool to monitor for genotoxic polycyclic compounds in Santos estuary. *Mutat Res* 609:60-67.
- Maertens RM, White PA, Rickert W, Levasseur G, Douglas GR, Bellier PV, McNamee JP, Thuppall V, Walker M, Desjardins S. 2009. The genotoxicity of mainstream and side stream marijuana and tobacco smoke condensates. *Chem Res Toxicol* 22:1406-1414.
- Maron DM, Ames BN. 1983. Revised methods for the *Salmonella* mutagenicity test. *Mutat Res* 113:173-215.
- Matsui K, Yamada M, Imai M, Yamamoto K, Nohmi T. 2006. Specificity of replicative and SOS-inducible DNA polymerases in frameshift mutagenesis: mutability of *Salmonella typhimurium* strains overexpressing SOS-inducible DNA polymerases to 30 chemical mutagens. *DNA Repair* 5:465-478.
- Mortelmans K, Zeiger E. 2000. The Ames *Salmonella*/microsome mutagenicity assay. *Mutat Res* 455:29-60.
- Mutlu E, Warren SH, Matthews PP, King C, Linak WP, Kooter IM, Schmid JE, Ross JA, Gilmour MI, DeMarini DM. 2013. Bioassay-directed fractionation and sub-fractionation for mutagenicity and chemical analysis of diesel exhaust particles. *Environ Mol Mutagen* 54:719-736.

- 1 Nielsen T, Jørgensen HE, Larsen JC, Poulsen M. 1996. City air pollution of polycyclic
2 aromatic hydrocarbons and other mutagens: occurrence, sources and health effects. *Sci*
3 *Total Environ* 189/190:41-49.
- 4 Sadiktsis I, Koegler JH, Benham T, Bergvall C, Westerholm R. 2014. Particulate associated
5 polycyclic aromatic hydrocarbon exhaust emissions from a portable power generator
6 fueled with three different fuels – A comparison between petroleum diesel and two
7 biodiesels. *Fuel* 115: 573-580.
- 8 Sato MIZ, Umbuzeiro GA, Coimã CA, Coelho MCLS, Sanches PS, Alonso CD, Martins
9 MT. 1995. Mutagenicity of airborne particulate organic material from urban and
10 industrial areas of São Paulo, Brazil. *Mutat Res* 335:317-330.
- 11 Sharma AK, Jensen KA, Rank J, White PA, Lundstedt S, Gagne R, Jacobsen NR, Kristiansen
12 J, Vogel U, Wallin H. 2007. Genotoxicity, inflammation and physico-chemical
13 properties of fine particle samples from an incineration energy plant and urban air.
14 *Mutat Res* 633:95-111.
- 15 Srogi K. 2007. Monitoring of environmental exposure to polycyclic aromatic hydrocarbons: a
16 review. *Environ Chem Lett* 5:169-195.
- 17 Umbuzeiro GA, Franco A, Magalhães D, Castro FJV, Kummrow F, Rech CM, Carvalho
18 LRF, Vasconcellos PC. 2008a. A preliminary characterization of the mutagenicity of
19 air particulate matter collected during sugar cane harvesting using the
20 *Salmonella/microsome* microsuspension assay. *Environ Mol Mutagen* 49:249-255.
- 21 Umbuzeiro GA, Franco A, Martins MH, Kummrow F, Carvalho LRF, Schmeiser HH,
22 Leykauf J, Stiborova M, Claxton LD. 2008b. Mutagenicity and DNA adduct formation
23 of PAH, nitro-PAH, and oxy-PAH fractions of air particulate matter from São Paulo,
24 Brazil. *Mutat Res* 652:72-80.

- 1 Umbuzeiro GA, Machala M, Weiss J. 2011. Diagnostic tools for effect-directed analysis of
2 mutagens, AhR agonists, and endocrine disruptors. In: Brack W, editor. Effect-
3 directed analysis of complex environmental contamination. Springer-Verlag,
4 Heidelberg. p. 69-82.
- 5 Uppstad H, Øvrebø S, Haugen A, Møllerup S. 2010. Importance of CYP1A1 and CYP1B1 in
6 bioactivation of benzo[a]pyrene in human lung cell lines. *Toxicol Lett* 192:221-228.
- 7 Viras LG, Athanasiou K, Siskos PA. 1990. Determination of mutagenic activity of airborne
8 particulates and of the benzo[α]pyrene concentrations in Athens atmosphere. *Atmos*
9 *Environ* 24B:267-274.
- 10 Walgraeve C, Demeestere K, Dewulf J, Zimmermann R, Van Langenhove H. 2010.
11 Oxygenated polycyclic aromatic hydrocarbons in atmospheric particulate matter:
12 Molecular characterization and occurrence. *Atmos Environ* 44:1831-1846.
- 13 Yamada M, Matsui K, Sofuni T, Nohmi T. 1997. New tester strains of *Salmonella*
14 *typhimurium* lacking O6-methylguanine DNA methyltransferases and highly sensitive
15 to mutagenic alkylating agents. *Mutat Res* 381:15-24.
- 16 Yauk CL, Williams A, Buick JK, Chen G, Maertens RM, Halappanavar S, White PA. 2012.
17 Genetic toxicology and toxicogenomic analysis of three cigarette smoke condensates
18 in vitro reveals few differences among full-flavor, blonde, and light products. *Environ*
19 *Mol Mutagen* 53:281-296.
- 20
21
22
23
24
25

Contribution of the Authors

Gisela A. Umbuzeiro: designed experiments, performed the Salmonella/microsome assay, analyzed data and drafted the manuscript

Fábio Kummrow: designed experiments, performed the Salmonella/microsome assay, analyzed data and drafted the manuscript

Daniel Alexandre Morales: designed experiments, performed the Salmonella/microsome assay and analyzed data

Debora Kristina M. Alves: designed experiments, performed the Salmonella/microsome assay and analyzed data

Hwanmi Lim: performed the chemical analysis and analyzed data

Ian W. H. Jarvis: designed experiments, analyzed data and drafted the manuscript

Christoffer Bergvall: designed experiments, analyzed data and drafted the manuscript

Roger Westerholm: designed experiments, analyzed data and drafted the manuscript

Ulla Stenius: designed experiments, analyzed data and drafted the manuscript

Kristian Dreij: designed experiments, analyzed data and drafted the manuscript

TABLE I. PAHs content (pg/m³) in extracts of the PM samples collected in Limeira, Brazil and Stockholm, Sweden.

PAH	LIMEIRA (pg/m ³)		STHOLM (pg/m ³)	
	Mean	SD	Mean	SD
Phenanthrene	798	27	250	2
Anthracene	180	14	33.6	0.7
3-Methylphenanthrene	47.0	2.5	23.8	0.7
2-Methylphenanthrene	65.4	2.1	32.0	1.3
2-Methylantracene	14.8	0.7	5.75	0.52
9-Methylphenanthrene	34.5	1.5	18.0	1.4
1-Methylphenanthrene	43.3	4.2	37.2	0.8
4H-Cyclopenta[def]phenanthrene	58.9	2.0	47.8	0.1
2-Phenylanthracene	35.0	2.9	26.9	2.6
3,6-Dimethylphenanthrene	5.03	0.20	1.70	0.11
3,9-Dimethylphenanthrene	22.9	1.5	8.31	0.75
Fluoranthene	671	6	368	6
Pyrene	895	25	439	8
1-Methylfluoranthene	40.9	3.5	46.1	3.7
Benzo[a]fluorene	28.7	1.2	27.0	2.4
Benzo[b]fluorene	16.2	1.5	13.0	0.8
2-Methylpyrene	30.8	2.9	18.4	1.4
4-Methylpyrene	52.3	4.7	30.7	2.9
1-Methylpyrene	47.7	3.9	32.1	3.2
Benzo[ghi]fluoranthene	346	15	207	8
Benzo[c]phenanthrene	48.5	3.4	77.6	2.1
Benzo[b]naphto(1,2-d)thiophene	9.95	0.31	2.16	0.71
Benz[a]anthracene	332	3	306	10
3-Methylchrysene	39.1	2.3	21.2	2.0
2-Methylchrysene	61.0	3.3	47.9	3.5
6-Methylchrysene	40.2	2.5	32.6	2.5
1-Methylchrysene	47.1	2.6	51.5	3.3
Benzo[b]fluoranthene	974	19	465	8
Benzo[k]fluoranthene	440	5	180	4
Benzo[e]pyrene	1050	31	352	13
Benzo[a]pyrene	899	17	289	4
Perylene	164	6	45.1	0.2
Indeno[1,2,3-cd]fluoranthene	65.4	1.5	28.2	0.3
Indeno[1,2,3-cd]pyrene	908	28	230	12
Dibenz[a,h]anthracene	55.9	2.7	35.6	3.3
Picene	76.2	2.2	30.9	1.7
Benzo[ghi]perylene	1620	35	351	7
Dibenzo[a,l]pyrene	4.79	0.37	5.14	0.28
Dibenzo[a,e]pyrene	15.7	0.6	32.4	2.0
Coronene	366	8	171	10
Dibenzo[a,i]pyrene	9.23	0.28	3.66	0.17
Dibenzo[a,h]pyrene	3.03	0.11	2.53	0.19
Total	10,663		4,427	

TABLE II. Summary of the main genetic characteristics of *Salmonella typhimurium* specific strains.

Strain	Description	Reversion event	Plasmids	Reference
TA1538	<i>his</i> D3052, Δ (<i>uvrB</i> , <i>bio</i>), <i>rfa</i>	Frameshift	No plasmids	Maron and Ames, [1983]
TA98	<i>his</i> D3052, Δ (<i>uvrB</i> , <i>bio</i>), <i>rfa</i> , Ap ^r	Frameshift	pKM101	Maron and Ames, [1983]
YG1041	<i>his</i> D3052, Δ (<i>uvrB</i> , <i>bio</i>), <i>rfa</i> , Ap ^r and Km ^r , NR and O-AT overproducing strain	Frameshift	pKM101, pYG233	Hagiwara et al. [1993]
YG5161	<i>his</i> D3052, Δ (<i>uvrB</i> , <i>bio</i>), <i>rfa</i> , Ap ^r , DNA Pol IV overproducing	Frameshift	pYG768	Matsui et al. [2006]

his – mutation in the histidine operon
 Δ *uvrB* – deletion of *uvrB* gene
 Δ *bio* – deletion of biotin gene
rfa – mutation cause partial loss of the lipopolysaccharide barrier
Ap^r – resistant to ampicillin
NR – nitroreductase
O-AT – O-acetyltransferase
DNA Pol. IV – DNA Polymerase IV
Km^r – resistant to Kanamycin

TABLE III. Mutagenicity for LIMEIRA PM extracts in the Salmonella/microsome microsuspension assay with TA98, YG1041, TA1538 and YG5161 without and with S9.

Doses		Mean of Number of Revertants/plate and <i>Standard Deviation (SD)</i>							
m ³ /plate	TA98				YG1041				
	-S9		+S9		-S9		+S9		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
Negative control	52.7	4.51	53.3	14.4	157.7	17.1	145.7	9.1	
0.02	53.5	2.12	58.0	7.1	262.0	24.1	151.5	17.7	
0.2	68.5	6.4	59.5	0.7	849.5	7.8	329.5	16.3	
1	144.0*	22.6	69.0	8.5	1,922.5	160.5	1,344.0	73.5	
2	207.5	45.9	59.0	2.8	3,583.5	231.2	3,312.5	50.2	
5	595.5	37.5	76.0	9.9	4,145.5	381.1	5,308.0	151.3	
10	1,110.0	70.7	133.0	18.4	3,861.5	362.7	6,355.5	0.7	
Blank Filter	60.0	1.4	47.0	2.8	146.0	30.4	137.0	11.3	
Positive control	927.0	101.1	1,518.0	352.8	908.0	173.9	2,344.0	7	
Potency revertants/m³	100		6.3		3,500		1,100		

Doses		Mean of Number of Revertants/plate and <i>Standard Deviation (SD)</i>							
m ³ /plate	TA1538				YG5161				
	-S9		+S9		-S9		+S9		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
Negative control	15.0	2.0	12.7	2.5	26.3	4.0	28.0	2.65	
0.02	16.5	0.7	14.5	2.1	23.5	0.7	29.5	7.8	
0.2	28.0	2.8	18.5	2.1	43.0	7.1	34.0	7.1	
1	89.5	0.7	52.0	9.9	99.0	2.8	57.5	2.1	
2	146.0	18.3	69.0	14.1	136.0	4.2	85.5	3.5	
5	306.5	24.7	147.5	13.4	272.0	4.2	162.0	7.1	
10	429.0	38.2	209.0	1.4	395.0	8.5	179.0	28.3	
Blank Filter	20.0	1.4	21.0	9.2	18.0	0.7	29.0	6.4	
Positive control	392.0	50.2	2,451.0	746.0	677.0	111.0	3,114.0	514.1	
Potency revertants/m³	62		28		61		27		

*Shaded values represent the number of revertants that equal or more than twice the negative controls

TABLE IV. Mutagenicity data for STHOLM PM extracts of the Salmonella/microsome microsuspension assay with TA98, YG1041, TA1538 and YG5161 without and with S9.

Doses	Mean of Number of Revertants/plate and <i>Standard Deviation (SD)</i>							
m ³ /plate	TA98				YG1041			
	-S9		+S9		-S9		+S9	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Negative control	52.7	4.5	53.3	14.4	157.7	17.1	145.7	9.1
0.02	58.5	6.4	51.5	3.5	173.5	6.4	157.0	5.7
0.2	61.0	12.7	43.0	1.4	364.5	2.1	173.5	16.3
1	73.0	1.4	50.0	5.7	936.0	135.7	418.5	4.9
2	93.0	14.1	58.0	8.5	2,108.0	159.8	683.5	37.5
5	195.0*	59.4	70.5	3.5	3,790.0	181.0	1,059.5	33.2
10	728.0	31.1	98.5	0.7	4,695.0	29.7	2,885.5	191.6
Blank filter	50.0	11.3	57.0	14.8	178.0	24.0	153.0	3.5
Positive control	927.0	101.1	1,518.0	352.8	908.0	173.9	2,344.0	171.1
Potency revertants/m³	25		4.8		890		270	

Doses	Mean of Number of Revertants/plate and <i>Standard Deviation (SD)</i>							
m ³ /plate	TA1538				YG5161			
	-S9		+S9		-S9		+S9	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Negative control	15.0	2.0	12.7	2.5	26.3	4.0	28.0	2.6
0.02	15.0	1.4	11.5	3.5	26.5	2.1	20.0	0.0
0.2	24.0	2.8	20.5	3.5	27.5	9.2	28.5	2.1
1	38.0	0.0	30.0	4.2	40.0	0.0	36.0	2.8
2	66.0	8.5	47.0	1.4	55.5	0.7	58.0	2.8
5	162.5	9.2	93.5	19.1	129.0	5.6	101.5	16.3
10	264.5	41.7	117.0	1.4	213.5	9.2	143.0	5.7
Blank filter	14.0	4.2	18.0	7.1	26.0	2.8	25.0	7.8
Positive control	392.0	50.2	2,451.0	746.0	677.0	111.0	3,114.0	514.1
Potency revertants/m³	26		12		18		15	

*Shaded values represent the number of revertants that equal or more than twice the negative controls

TABLE V. Mutagenic responses in the *Salmonella*/microsome microsuspension assay for benzo[a]pyrene with the strains, TA98, YG1041, TA1538 and YG5161 in the presence of S9.

Doses ng/plate	Mean of Number of Revertants/plate and <i>Standard Deviation (SD)</i>							
	TA98		YG1041		TA1538		YG5161	
	Mean	<i>SD</i>	Mean	<i>SD</i>	Mean	<i>SD</i>	Mean	<i>SD</i>
Negative control	28.0	7.1	159.0	21.2	15.5	0.7	21.0	5.7
0.5							32.0	7.1
1.0							28.5	0.7
2.0							34.0	2.8
4.0							48.5	3.5
7.8	32.5	3.5	164.5	29.0	14.5	0.7	47.5	2.1
15.6	31.0	1.7	183.0	35.4	25.0	5.7	71.5	6.4
31.2	37.5	3.5	210.5	10.6	30.5	4.9	114.0	1.4
62.5	30.5	7.8	214.0	21.2	44.0	17.0	110.5	12.0
125	37.5	9.2	306.0	31.1	49.5	12.0	155.0	8.5
250	42.5	6.4	339.0	38.2	68.0	4.2	203.5	19.1
500	39.0	5.7	462.5	17.7	98.5	10.6	230.0	5.7
1,000	60.0*	14.1	471.0	9.9	125.5	27.6	285.0	1.4
2,000	63.0	11.3	470.5	0.7	106.0	5.7	271.5	2.1
Potency revertants/ng	0.02		0.6		0.2		3.0	
MED**	1,400		265		77.5		7	

*Shaded values represent the number of revertants that equal or more than twice the negative controls

**MED = Minimal Effective Dose in ng/plate

TABELA VI. Mutagenic responses in the *Salmonella*/microsome microsuspension assay for 1-nitropyrene with the strains, TA98, YG1041, TA1538 and YG5161 in the absence of S9.

Doses ng/plate	Mean of Number of Revertants/plate and <i>Standard Deviation (SD)</i>							
	TA98		YG1041		TA1538		YG5161	
	<i>Mean</i>	<i>SD</i>	Mean	<i>SD</i>	Mean	<i>SD</i>	Mean	<i>SD</i>
Negative control	42.5	4.9	201.0	21.2	18.0	0	24.0	4.2
0.01			257.5	10.6				
0.02			310.5	23.3				
0.03			383.0	35.4				
0.04			410.5	2.1				
0.05			506.5	24.7				
0.1	48.0	2.8	791.5	3.5	18.5	0.7	33.5	10.6
0.3	58.0	15.6	3,063.5	101.1	17.5	7.8	29.0	5.7
0.7	101.0*	7.1	4,353.0	110.3	28.5	6.4	45.5	0.7
1.0	129.5	0.7			19.0	5.7	36.0	5.7
2.5	266.0	41.0			36.5	3.5	68.0	11.3
5.0	536.5	140.7			63.5	0.7	76.5	13.4
10	1,001.0	75.0			139.5	46.0	135.0	33.9
20	1,883.5	82.7			595.0	110.3	334.0	19.8
Potency revertants/ng	94		5,900		10		13	
MED**	0.45		0.03		1.8		1.8	

*Shaded values represent the number of revertants that equal or more than twice the negative controls

**MED - Minimal Effective Dose in ng/plate