**Karolinska
Institutet**

Karolinska Institutet

<http://openarchive.ki.se>

This is a Peer Reviewed Accepted version of the following article, accepted for publication in Gut.

2018-01-11

Fetal and early life antibiotics exposure and very early onset inflammatory bowel disease : a population-based study

Örtqvist, Anne K; Lundholm, Cecilia; Halfvarson, Jonas; Ludvigsson, Jonas F; Almqvist, Catarina

Gut. 2019 Feb;68(2):218-225.

British Society of Gastroenterology

<http://doi.org/10.1136/gutjnl-2017-314352>

<http://hdl.handle.net/10616/46186>

If not otherwise stated by the Publisher's Terms and conditions, the manuscript is deposited under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

**Karolinska
Institutet**

This is an author produced version of a paper accepted by **Gut**. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Fetal and early life antibiotics exposure and very early onset inflammatory bowel disease: a population-based study

Örtqvist, Anne K; Lundholm, Cecilia; Halfvarson, Jonas; Ludvigsson, Jonas F; Almqvist, Catarina

Gut : 10 January 2018. [Published Online First]

DOI: [10.1136/gutjnl-2017-314352](https://doi.org/10.1136/gutjnl-2017-314352)

Access to the published version may require subscription.
Published with permission from: **BMJ**

**Fetal and early life antibiotics exposure and very early onset inflammatory
bowel disease – a population-based study**

Short title: Antibiotics and inflammatory bowel disease

AUTHORS

Anne K. Örtqvist, postdoctoral researcher¹, Cecilia Lundholm, biostatistician¹, Jonas Halfvarson,
senior lecturer², Jonas F. Ludvigsson, professor^{1,3}, Catarina Almqvist, professor^{1,4}

1. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, PO Box 281,
171 77 Stockholm, Sweden

2. Department of Gastroenterology, Faculty of Medicine and Health, Örebro University, 701 82
Örebro, Sweden

3. Department of Pediatrics, Örebro University Hospital, 701 85 Örebro, Sweden

4. Pediatric Allergy and Pulmonology Unit at Astrid Lindgren Children's Hospital, Karolinska
University Hospital, 17176 Stockholm, Sweden

ABBREVIATIONS

aHR – adjusted Hazard Ratio

CD - Crohn's disease

cHR – crude Hazard Ratio

CI – Confidence Interval

IBD – Inflammatory Bowel Disease

MBR – Medical Birth Register

NPR – National Patient Register

26 PcV – Phenoxymethylpenicillin
27 PIN – Personal Identification Number
28 SPDR – Swedish Prescribed Drug Register
29 TPR – Total Population Register
30 UC – Ulcerative Colitis
31 VEO-IBD – Very Early Onset Inflammatory Bowel Disease

32

33

34 **CORRESPONDING AUTHOR**

35 Professor Catarina Almqvist, MD PhD

36 Dept. Medical Epidemiology and Biostatistics

37 PO Box 281, Karolinska Institutet

38 SE 171 77 Stockholm, SWEDEN

39 Email. catarina.almqvist@ki.se

40 Telephone. +46 70 116 0852

41

42 **KEY WORDS:** antibiotics; Crohn’s disease; Ulcerative colitis; population-based registers; very
43 early onset (VEO) IBD

44 **WORD COUNT:** 4309

45 **REFERENCES:** 56

46 **TABLES:** 5

47 **FIGURES:** 2

48

49

ABSTRACT

Objective Earlier studies on antibiotics exposure and development of inflammatory bowel disease (IBD, Crohn's disease and ulcerative colitis) may have been biased by familial factors and gastroenteritis. We aimed to estimate the association between antibiotics during pregnancy or infantile age and very early onset (VEO-) IBD.

Design In this cohort study of 827 239 children born in Sweden 2006-2013, we examined the link between exposure to systemic antibiotics and VEO-IBD (diagnosis <6 years of age), using Cox proportional hazard regression models. Information on antibiotics and IBD was retrieved from the nationwide population-based Swedish Prescribed Drug Register and the National Patient Register. We specifically examined potential confounding from parental IBD and gastroenteritis.

Results Children exposed to antibiotics during pregnancy were at increased risk of IBD compared to general population controls (adjusted hazard ratio (aHR) 1.93; 95% confidence interval (CI) 1.06-3.50). Corresponding aHRs were 2.48 (1.01-6.08) for Crohn's disease (CD) and 1.25 (0.47-3.26) for ulcerative colitis (UC) respectively. For antibiotics in infantile age, the aHR for IBD was 1.11 (0.57-2.15); for CD 0.72 (0.27-1.92) and 1.23 (0.45-3.39) for UC. Excluding children with gastroenteritis 12 months prior to the first IBD diagnosis retained similar aHR for antibiotics during pregnancy and CD, while the association no longer remained significant for IBD.

Conclusion We found that exposure to antibiotics during pregnancy, but not in infantile age, is associated with an increased risk of VEO-IBD regardless of gastroenteritis. The risk increase for exposure in pregnancy may be due to changes in the microbiota.

72 **Summary “box”**

73 **What is already known about this subject?** Very early onset (VEO-) inflammatory bowel
74 disease (IBD) has gradually become more common. One potential risk factor for VEO-IBD is
75 antibiotic exposure during pregnancy and in infantile age.

76 **What are the new findings?** In this population-based study of more than 800 000 children, we
77 found a positive association between antibiotics exposure during pregnancy, but not in infantile
78 age, and later VEO-IBD.

79 **How might it impact on clinical practice in the foreseeable future?** The risk increase for
80 exposure in pregnancy may be due to changes in the microbiota, which could have an impact on
81 care of pregnant women. However, the absolute risk of disease was very low, and antibiotics
82 during pregnancy should still be used when needed.

83

INTRODUCTION

Inflammatory bowel disease (IBD), comprising Crohn's disease (CD) and ulcerative colitis (UC), is characterized by chronic inflammation of the gastrointestinal tract. Symptoms related to the disease include diarrhoea, rectal bleeding, abdominal pain and weight loss. Although children (diagnosed at <18 years of age) may present with these classical symptoms, non-specific symptoms such as growth failure, anaemia and other extra-intestinal manifestations are also common manifestations of paediatric IBD.¹ Interestingly, patients with very early onset (VEO)-IBD (defined as diagnosed before six years of age) seem to represent a specific entity, possibly more likely to present with rectal bleeding due to a colonic phenotype (in CD)² and a family history of IBD.²⁻⁵ This disease entity has gradually become more common and recent data suggest that the increase in IBD incidence is more pronounced in patients with VEO-IBD than among children ≥ 6 years old, although the overall numbers are still very small compared to young adults.⁶

The pathogenesis is characterized by a complex interaction between genetics, an aberrant mucosal immune response to gut microbiota, disruption of gut barrier, and environmental triggers. While genetic factors clearly play an important role in the etiology of IBD,⁷⁻¹⁰ they cannot explain the recent rise in disease incidence or the proband concordance rate of 38-62% in monozygotic twins with CD (even less in UC).^{11 12} This has encouraged a search for environmental factors and led to the identification of factors such as smoking, oral contraceptives and appendectomy due to appendicitis (data on appendectomy have however been contradictory)¹³ as potential risk factors in adulthood IBD while less is known for childhood IBD.¹⁴⁻¹⁹

To what extent pathophysiologic mechanism of IBD can be applied to VEO-IBD remains partly unknown. It has been proposed that the influence of genetics might be more pronounced in VEO-IBD, due to the effect of rare variants with a high penetrance for IBD.²⁰ However, the role of environmental risk factors in VEO-IBD is largely unknown.

One potential risk factor for VEO-IBD is antibiotic exposure during pregnancy and in infantile age. Ungaro *et al.* reported an increased risk of IBD following antibiotic exposure, especially in children, in a recent meta-analysis of eleven observational studies.²¹ The association was limited to individuals with newly onset CD and not seen for UC. Some earlier studies have also found a positive dose-response relationship between antibiotics exposure and later IBD, but whether this also applies to VEO-IBD remains unknown.

Therefore, we aimed to estimate the association between antibiotics during pregnancy or infantile age and VEO-IBD (CD and UC), while adjusting for parental IBD, as well as taking gastroenteritis and number of doses into account in a population-based study.

METHOD

Study population and register linkage

This nationwide prospective population-based register study included all children born January 2006 to December 2013, identified together with their mothers from the Swedish Medical Birth Register (MBR). Fathers to the children were identified through the Swedish Multi-generation Register.²² We used the Swedish Prescribed Drug Register (SPDR)²³ and the National Patient Register (NPR)²⁴ to retrieve data on exposure and outcome. The SPDR contains complete data on all dispensed drugs from pharmacies since July 2005 while the NPR began in 1964, became nationwide in 1987 (inpatient diagnoses only), and added hospital-based outpatient visit data in 2001. Data from the different registers were linked using the unique personal identity number assigned to all residents in Sweden.²⁵

Variables

Antibiotic exposure was defined as filled prescriptions of relevant ATC codes (Anatomical Therapeutic Chemical): J01A-J01X (systemic antibiotics), and were obtained through the SPDR. We categorised antibiotics into two groups: “systemic antibiotics” which included any type of antibiotics; and “PcV” (Phenoxymethylpenicillin), which is by far the most commonly prescribed type of antibiotic in Sweden both during pregnancy and in childhood.²⁶

Our outcome IBD was defined as having ≥ 2 diagnoses of either CD or UC according to relevant International Classification of Disease (ICD)-10 codes K50 and K51 respectively recorded in the NPR, with onset before 6 years of age. Crohn’s disease was defined as ≥ 2 CD diagnoses but never an ulcerative colitis diagnosis and UC was defined as ≥ 2 UC diagnoses but never a CD diagnosis. These outcome definitions have previously been validated by Jakobsson *et al.* who found a positive predictive value (PPV) of 93% (95% CI: 87–97) for

IBD, 90% (77–97) for UC and 81% (67–91) for CD, when compared to the Copenhagen criteria.²⁷ While IBD-unclassified is now regarded as a separate entity (and some data suggests that it may represent up to 20% of the total pediatric IBD population),²⁸ this paper focused on CD and UC.

Co-variables; Through the MBR we retrieved information on the child's date of birth, sex, gestational age (days), mode of delivery (vaginal or caesarean section), maternal smoking at first visit to the antenatal care clinic (yes/no), maternal age at delivery (≤ 19 , 20–24, 25–29, 30–34, or ≥ 35 years), and parity (child's birth order at current delivery; first-born (1) or not (≥ 2)). Data from Cnattingius et al. suggest a high quality of data in the MBR with a coverage of >98% of all births in Sweden.²⁹

Maternal and paternal IBD was defined from the NPR similarly to that of the children (two records of either: ICD-8: 563.00, 563.10, 569.02; ICD-9: 555, and ICD-10: K50 for CD; and ICD-8: 563.98, 563.99; ICD-9: 556, and ICD-10: K51 for UC). Information on highest level of education of either parent (0–12 vs >12 years) was identified through the Longitudinal integration database for health insurance and labour market studies and parents' country of birth (Sweden or other) was identified from the Total Population Register (TPR).

We defined the beginning of the pregnancy (conception date) as date of birth minus gestational age in days. We furthermore divided pregnancy duration into trimester 1 (day 1–91), 2 (day 92–189), and 3 (day 190+) to examine time-varying effects of fetal antibiotics exposure. To ensure that the full pregnancy would be covered by this study (especially antibiotics exposure in the first trimester in offspring born close to the starting point of the SPDR), we restricted our study population to children with estimated conception date as on or after July 1st, 2005. Information on migration and death was obtained from the TPR to be able to define end of

172 follow-up. Offspring to women who immigrated to Sweden during pregnancy were excluded
173 from the study (n=14 030).

174
175 **Statistical analysis**

176 The risk of IBD, and CD and UC separately, in children exposed to antibiotics during pregnancy
177 and in infantile age compared to unexposed children, was examined in Cox proportional hazard
178 models. Attained age was used as the underlying time scale and clustering within families was
179 taken into account by using a sandwich estimator for the standard errors. Follow-up ended with
180 first IBD diagnosis, emigration, death or end of study period (December 31st, 2014 i.e at least one
181 year of follow-up), whichever happened first. The proportional hazards assumption was tested
182 using Schoenfeld's residuals (*p*-values of 0.58 for exposure during pregnancy and 0.84 for
183 exposure in infantile age). Antibiotic exposure during pregnancy was regarded as exposed or non-
184 exposed at the start of follow-up, while exposure after birth was modelled as a time(age)-varying
185 exposure, i.e. a model in which all individuals start as unexposed and then the exposure status
186 changes at the time an individual becomes exposed. However, as there was no sign of non-
187 proportional hazards we did not allow for the effect of the exposure to vary over time.

188 Potential confounders were identified based on the Directed Acyclic Graphs (DAG)
189 concept,³⁰ and the final models were adjusted for: mother's and father's history of IBD, parental
190 education, mother's and father's country of birth (in analyses of exposure during pregnancy and
191 in childhood) and mode of delivery (in analysis of exposure in infantile age) as shown in *Figure*
192 *1*. Those with missing information on these variables were excluded (n=2848 for exposure in
193 pregnancy and n=3274 for exposure in infantile age) and complete case analyses were performed.
194 Crude and adjusted Hazard Ratios (HR) are presented.

Additional analyses: Sensitivity analyses were performed by excluding all individuals who had been diagnosed with gastroenteritis, either bacterial or viral, according to ICD 10 codes A00-A09, within 12 months prior to onset of their first IBD diagnosis. The risk of IBD with onset after 2 years of age in children exposed to systemic antibiotics in the first year of life, was further tested to investigate potential information bias such as misclassification or reverse causation. This since a possible misclassification of first occurrence of IBD symptoms as gastroenteritis, treated with antibiotics, may induce an association between antibiotics and IBD. A potential dose-relationship between number of antibiotics prescriptions in childhood (1, 2 or ≥ 3 as a continuous variable) and IBD was tested. Dispensed prescriptions of the same type of antibiotics within 7 days from the prior dispense was counted as one dispense. Too few women had filled more than one prescription of antibiotics during pregnancy to be able to perform dose-response-analyses. The role of timing of systemic antibiotics exposure during the fetal period (1st vs 2nd vs 3rd trimester) and IBD was further explored. We also examined if the risk increase by antibiotic exposure was dependent on parity by adding interaction terms between parity (first-born vs. non-first born) and antibiotic exposure.

STATA statistical software (version 14) was used for all statistical analyses.

The study was approved by the Regional Ethical Review board in Stockholm, Sweden.

RESULTS

The final study population consisted of 827 239 children (*Figure 2*). Some 12 606 children emigrated during follow-up, and another 2428 died.

Overall, 17% (n=140 665) of the children had been exposed to antibiotics during pregnancy, and 5% (n=40 116) had been exposed on two or more occasions. In infantile age, 65% (n=539 809) had been exposed to systemic antibiotics at least once, and 373 802 (70% of those exposed at all) had filled two or more prescriptions (*Table 1*).

222 **Table 1.** Descriptive table of the study population.

Variable	All		IBD		CD		UC	
	N	%	n	%	n	%	n	%
Sex	827 239	100.0	51	100.0	20	100.0	24	100.0
<i>Males</i>	425 212	51.4	26	51.0	12	60.0	11	45.8
<i>Females</i>	402 027	48.6	25	49.0	8	40.0	13	54.2
Systemic antibiotic exposure during pregnancy	140 665	17.0	15	29.4	7	35.0	5	20.8
<i>Phenoxymethylpenicillin (PcV)</i>	60 701	7.3	8	15.7	4	20.0	3	12.5
<i>Pivmecillinam</i>	34 264	4.1	2	3.9	1	5.0	1	4.2
<i>Nitrofurantoin</i>	30 904	3.7	4	7.8	1	5.0	2	8.3
<i>Other</i>	42 672	5.2	5	9.8	4	20.0	0	0
Number of prescriptions during pregnancy (any)*								
<i>1</i>	100 549	12.2	10	19.6	4	20.0	3	12.5
<i>2</i>	26 357	3.2	3	5.9	2	10.0	1	4.2
<i>≥3</i>	13 759	1.7	2	3.9	1	5.0	1	4.2
Systemic antibiotic exposure in infantile age	539 809	65.3	43	84.3	16	80.0	20	83.3
<i>Phenoxymethylpenicillin (PcV)</i>	460 283	55.6	38	74.5	14	70.0	17	70.8
<i>Amoxicillin</i>	152 052	18.4	15	29.4	7	35.0	5	20.8
<i>Flucloxacillin</i>	80 814	9.8	13	25.5	5	25.0	4	16.7
<i>Other</i>	204 387	24.7	28	54.9	10	50.0	11	45.8
Number of prescriptions in infantile age (any)*								
<i>1</i>	166 007	20.1	3	5.9	1	5.0	2	8.3
<i>2</i>	111 761	13.5	8	15.7	2	10.0	6	25.0
<i>≥3</i>	262 041	31.7	32	62.7	13	65.0	12	50.0
Mothers' diagnoses								
<i>IBD</i>	6 780	0.8	7	13.7	4	20.0	2	8.3
<i>CD</i>	1 943	0.2	2	3.9	2	10.0	0	0.0
<i>UC</i>	3 598	0.4	5	9.8	2	10.0	2	8.3
Fathers' diagnoses								
<i>IBD</i>	7 333	0.9	4	7.8	0	0.0	2	8.3
<i>CD</i>	1 868	0.2	0	0.0	0	0.0	0	0.0
<i>UC</i>	3 972	0.5	1	2.0	0	0.0	0	0.0
Parents' highest achieved education								
<i>0-12 years</i>	337 312	40.8	18	35.3	8	40.0	7	29.2
<i>>12 years</i>	485 926	58.7	33	64.7	12	60.0	17	70.8
<i>Missing</i>	4 001	0.5	0	0.0	0	0.0	0	0.0
Mother's country of birth								
<i>Sweden</i>	646 032	78.1	42	82.4	18	90.0	20	83.3
<i>Other</i>	181 207	21.9	9	17.7	2	10.0	4	16.7
Father's country of birth								

<i>Sweden</i>	630 372	76.2	39	76.5	18	90.0	19	79.2
<i>Other</i>	178 669	21.6	12	23.5	2	10.0	5	20.8
<i>Missing</i>	18 198	2.2	0	0.0	0	0.0	0	0.0
Mode of delivery								
<i>Vaginal</i>	679 693	82.2	43	84.3	16	80.0	20	83.3
<i>Caesarean Section</i>	147 107	17.8	8	15.7	4	20.0	4	16.7
<i>Missing</i>	439	0.1	0	0.0	0	0.0	0	0.0
Maternal smoking at first visit to the antenatal care clinic								
<i>Yes</i>	51 892	6.3	3	5.9	1	5.0	1	4.2
<i>No</i>	741 727	89.7	46	90.2	17	85.0	23	95.8
<i>Missing</i>	33 620	4.1	2	3.9	2	10.0	0	0.0
Maternal age at delivery (years)								
≤ 19	12 060	1.5	0	0.0	0	0.0	0	0.0
20-24	105 879	12.8	4	7.8	2	10.0	1	4.2
25-29	237 650	28.7	18	35.3	7	35.0	7	29.2
30-34	287 319	34.7	18	35.3	5	25.0	12	50.0
≥ 35	184 330	22.3	11	21.6	6	30.0	4	16.7
<i>Missing</i>	1	0.0	0	0.0	0	0.0	0	0.0
Parity (at current delivery)								
<i>1</i>	361 332	43.7	26	51.0	11	55.0	10	41.7
≥ 2	465 907	56.3	25	49.0	9	45.0	14	58.3

223

224 *Any* – Any type of systemic antibiotics (ATC: J01)

225 * Independent on time in relation to diagnosis (i.e. either before or after onset of diagnosis)

226 “Other” antibiotics during pregnancy – Tetracyclines (J01A), penicillins with extended spectrum
227 (J01CA) (except pivmecillinam), beta-lactamase sensitive penicillins (J01CE) (except
228 phenoxymethylpenicillin), beta-lactamase resistant penicillins (J01CF), combinations of
229 penicillins, incl. beta-lactamase inhibitors (J01CR), cephalosporins (J01DB-DD), monobactams
230 (J01DF), carbapenems (J01DH), trimethoprim (J01EA), combinations of sulfonamides and
231 trimethoprim (J01EE), macrolides (J01FA), lincosamides (J01FF), other aminoglycosides
232 (J01GB), fluoroquinolones (J01MA), other antibacterials (J01X) (except nitrofurantoin).

233 “Other” antibiotics during childhood – Tetracyclines (J01A), penicillins with extended spectrum
234 (J01CA) (except amoxicillin), beta-lactamase resistant penicillins (J01CF) (except
235 flucloxacillin), combinations of penicillins, incl. beta-lactamase inhibitors (J01CR),
236 cephalosporins (J01DB-DE), monobactams (J01DF), carbapenems (J01DH), trimethoprim
237 (J01EA), combinations of sulfonamides and trimethoprim (J01EE), macrolides (J01FA),
238 lincosamides (J01FF), other aminoglycosides (J01GB), fluoroquinolones (J01MA), other
239 antibacterials (J01X).

240

The median time from conception to exposure to systemic antibiotics was 125 days (interquartile range (IQR) 64-197 days), while the median age at first exposure to systemic antibiotics in childhood was 1.3 years (IQR 0.8-2.1 years).

In total, 51 children with IBD (CD and/or UC), 20 with CD (but never an UC diagnosis) and 24 with UC (but never a CD diagnosis) could be identified through the NPR. Approximately 14% of the children with IBD had a mother with IBD and 8% had a father with IBD, compared to those children without IBD, where the corresponding numbers for parental IBD were less than 1% (0.8% mothers and 0.9% fathers) (*Table 1*). The median age of the first IBD diagnosis was 2.0 years (IQR 0.9-4.1 years).

Antibiotics during pregnancy

Table 2 presents crude and adjusted Hazard Ratios (aHR) and 95% Confidence Intervals (CI) for the association between exposure to antibiotics during pregnancy and IBD, CD and UC respectively. There was a 93% significantly increased risk of IBD in children exposed to systemic antibiotics during pregnancy (aHR 1.93, 95% CI 1.06-3.50). An association remained for CD (aHR 2.48, 95% CI 1.01-6.08), but not for UC (aHR 1.25, 95% CI 0.47-3.26). The aHR seemed to remain when restricting systemic antibiotic exposure to PcV (aHR for IBD 2.15, 95% CI 1.02-4.56), although no longer significant for CD (aHR 2.85, 95% CI 0.96-8.45). . Still, no significant association was found between “PcV” and UC.

In total, six children had been diagnosed with gastroenteritis 12 months prior to the first diagnosis of IBD, where five had been diagnosed with a viral or unspecified gastroenteritis and colitis (ICD 10: A09) and one had been diagnosed with a bacterial gastroenteritis with *Clostridium difficile* (ICD 10: A047). In sensitivity analyses, excluding these individuals, similar aHR remained for systemic antibiotics and CD (aHR 2.51, 95% CI 0.96-6.56), although no longer

264 significant, and with lower non-significant risk estimates for systemic antibiotics and IBD (aHR
265 1.68, 95% CI 0.88-3.21) (*Table 2*).

Table 2. Crude and adjusted Hazard ratios and 95% confidence intervals for inflammatory bowel disease (IBD) in relation to exposure to systemic antibiotics and “PcV” during pregnancy, and results from sensitivity analysis excluding those who have been diagnosed with gastroenteritis (GE) 12 months prior to onset of the first IBD diagnosis.

Antibiotics	IBD			Crohn’s disease (CD)			Ulcerative colitis (UC)		
	n	cHR (95% CI)	aHR (95% CI)	n	cHR (95% CI)	aHR (95% CI)	n	cHR (95% CI)	aHR (95% CI)
<i>Any</i>	15	1.96 (1.07-3.57)	1.93 (1.06-3.50)	7	2.52 (1.01-6.28)	2.48 (1.01-6.08)	5	1.24 (0.47-3.32)	1.25 (0.47-3.26)
<i>exl.GE</i>	12	1.70 (0.88-3.29)	1.68 (0.88-3.21)	6	2.55 (0.95-6.86)	2.51 (0.96-6.56)	4	1.05 (0.36-3.08)	1.05 (0.36-3.08)
<i>PcV</i>	8	2.19 (1.03-4.66)	2.15 (1.02-4.56)	4	2.92 (0.99-8.67)	2.85 (0.96-8.45)	3	1.71 (0.51-5.71)	1.68 (0.51-5.97)
<i>exl.GE</i>	7	2.17 (0.97-4.84)	2.13 (0.96-4.75)	3	2.50 (0.73-8.62)	2.45 (0.71-8.42)	3	1.87 (0.56-6.34)	1.85 (0.55-6.21)

269

270 cHR – Crude Hazard Ratio; 3 945 000 Person-years

271 aHR – Adjusted Hazard Ratio; 3 874 000 Person-years; analyses adjusted for mother’s and father’s history of IBD, parental education,
272 mother’s and father’s country of birth. Those with missing information on these variables are excluded (no cases excluded).

273 n – cases, *i.e.* those who have been exposed to antibiotics prior to onset of IBD/UC/CD diagnosis

274 *exl.GE* – excluding gastroenteritis, either bacterial or viral, according to ICD 10 codes A00-A09 12 months prior to onset of the first
275 IBD diagnosis.

276 *PcV* – Phenoxymethylpenicillin

277 The role of timing of systemic antibiotics exposure during pregnancy and IBD was
 278 further explored, where aHR for the first trimester was 1.59 (95% CI 0.64-3.97), second trimester
 279 1.23 (95% CI 0.45-3.40) and with a significant association during the third trimester (aHR 2.57
 280 95% CI 1.10-6.01) (*Table 3*). Too few cases were available for separate analyses of exposure in
 281 different trimesters and CD/UC.

282 **Table 3.** Crude and adjusted Hazard ratios and 95% confidence intervals for the association
 283 between systemic antibiotics and inflammatory bowel disease (IBD) in different trimesters during
 284 pregnancy.

285

IBD				
Trimester	n	cHR (95% CI)	aHR (95% CI)	
First	5	1.59 (0.63-4.01)	1.59 (0.64-3.97)	286
Second	4	1.27 (0.46-3.51)	1.23 (0.45-3.40)	287
Third	6	2.60 (1.11-6.10)	2.57 (1.10-6.01)	288

289 cHR – Crude Hazard Ratio; 3 945 000 Person-years

290 aHR – Adjusted Hazard Ratio; 3 874 000 Person-years; analyses adjusted for mother's and
 291 father's history of IBD, parental education, mother's and father's country of birth. Those with
 292 missing information on these variables are excluded (no cases excluded).

293 n – cases, *i.e.* those who have been exposed to antibiotics prior to onset of IBD

Antibiotics in infantile age

Table 4 presents crude and adjusted Hazard Ratios (aHR) and 95% Confidence Intervals (CI) for the association between exposure to antibiotics in infantile age and IBD, CD and UC respectively. No significant associations were found for systemic antibiotics and “PcV” for IBD, CD or UC, where aHR varied between 0.72 (95% CI 0.27-1.92) (CD), 1.23 (95% CI 0.45-3.39) (UC) and 1.11 (95% CI 0.57-2.15) (IBD) for systemic antibiotics and 0.87 (95% CI 0.33-2.27) (CD), 1.20 (95% CI 0.51-2.81) (UC) and 1.25 (95% CI 0.70-2.26) (IBD) for “PcV”. The associations remained non-significant after excluding individuals with gastroenteritis for both systemic antibiotics and “PcV” and all outcomes (IBD, CD and UC).

Table 4. Crude and adjusted Hazard ratios and 95% confidence intervals for the association between exposure to systemic antibiotics and “PcV” in infantile age and inflammatory bowel disease (IBD), and results from sensitivity analysis excluding those who had been diagnosed with gastroenteritis (GE) 12 months prior to onset of the first IBD diagnosis.

Antibiotics	IBD			Crohn's disease (CD)			Ulcerative colitis (UC)		
	n	cHR (95% CI)	aHR (95% CI)	n	cHR (95% CI)	aHR (95% CI)	n	cHR (95% CI)	aHR (95% CI)
<i>Any</i>	25	1.13 (0.58-2.20)	1.11 (0.57-2.15)	9	0.72 (0.27-1.94)	0.72 (0.27-1.92)	11	1.22 (0.44-3.37)	1.23 (0.45-3.39)
<i>exl.GE</i>	23	1.28 (0.60-2.73)	1.26 (0.60-2.64)	7	0.66 (0.22-1.99)	0.66 (0.23-1.92)	11	1.39 (0.46-4.25)	1.40 (0.46-4.25)
<i>PcV</i>	22	1.27 (0.70-2.31)	1.25 (0.70-2.26)	8	0.88 (0.33-2.31)	0.87 (0.33-2.27)	9	1.19 (0.50-2.80)	1.20 (0.51-2.81)
<i>exl.GE</i>	21	1.55 (0.80-3.00)	1.52 (0.80-2.90)	7	1.07 (0.36-3.21)	1.06 (0.37-3.10)	9	1.29 (0.52-3.23)	1.31 (0.53-3.22)

306

307 cHR – Crude Hazard Ratio; 3 945 000 Person-years

308 aHR – Adjusted Hazard Ratio; 3 874 000 Person-years; analyses adjusted for mother's and father's history of IBD, parental education,
309 mother's and father's country of birth and mode of delivery. Those with missing information on these variables are excluded (no cases
310 excluded).

311 n – cases, *i.e.* those who have been exposed to antibiotics prior to onset of IBD/UC/CD diagnosis

312 *exl.GE* – excluding gastroenteritis, either bacterial or viral, according to ICD 10 codes A00-A09 12 months prior to onset of the first
313 IBD diagnosis.

314 *PcV* – Phenoxymethylpenicillin

Exploring the association between individuals who had been exposed to systemic antibiotics during the first year of life and the risk of first IBD diagnosis from 2 years of age, the aHR was 1.49 (95% CI 0.69-3.22). Furthermore, no significant interaction ($p=0.48$) was found between systemic antibiotics and parity and IBD, where the aHR in first-born was 0.93 (95% CI 0.39-2.22) and 1.38 in non-first-borns (95% CI 0.58-3.27) (data not tabulated).

No dose-response relationship was found between increasing number of prescriptions for systemic antibiotics in infantile age and IBD (*Table 5*).

Table 5. Crude and adjusted Hazard ratios and 95% confidence intervals for inflammatory bowel disease (IBD) in relation to increasing number of antibiotic prescriptions.

Systemic antibiotics, filled prescriptions	IBD		
	n	cHR (95% CI)	aHR (95% CI)
1	6	0.73 (0.28-1.92)	0.73 (0.28-1.89)
2	9	1.87 (0.81-4.34)	1.85 (0.80-4.30)
≥3	10	1.15 (0.48-2.75)	1.12 (0.47-2.75)

cHR – Crude Hazard Ratio; 3 945 000 Person-years

aHR – Adjusted Hazard Ratio; 3 874 000 Person-years; analyses adjusted for mother's and father's history of IBD, parental education, mother's and father's country of birth and mode of delivery. Those with missing information on these variables are excluded (no cases excluded).

n – cases, *i.e.* those who have been exposed to antibiotics prior to onset of IBD diagnosis

DISCUSSION

In this nationwide population-based birth cohort study of more than 800 000 children, we found a positive association between antibiotics exposure during pregnancy and later VEO-IBD and CD, but not UC. The aHR seemed to remain when restricting systemic antibiotic exposure to PcV, although no longer significant for CD. The risk estimates for systemic antibiotics and CD remained similar after exclusion of children with gastroenteritis 12 months prior to their first IBD diagnosis. No association was found between systemic antibiotics or “PcV” in infantile age and later VEO-IBD, CD or UC, independent of exclusion of individuals with gastroenteritis prior to their first IBD diagnosis. This is important as gastroenteritis may have represented undiagnosed IBD, and resulted in antibiotics treatment occurring after IBD rather than preceding it. Furthermore, there was no significant association for children who had been exposed to systemic antibiotics during the first year of life and the risk of first IBD diagnosis from 2 years of age, or between increasing numbers of filled prescriptions for antibiotics and IBD.

Previous literature

While recent research has confirmed that antibiotic use is associated with an increased risk of IBD, including pediatric CD, less attention has been paid to VEO-IBD. In a recent meta-analysis, Ungaro *et al.* examined 11 studies,³¹⁻⁴¹ with four focusing on paediatric IBD.^{34 37 39 40} The authors regarded only three of these as paediatric,^{34 37 39} and when pooling their data the HR was substantially higher than in the overall meta-analysis (HR 2.75, 95% CI 1.72-4.38) for children only).

Antibiotics may influence the risk of acquiring IBD in several ways. Firstly antibiotics may be causally related to IBD, potentially by a mediating effect on the microbiome, causing a reduced diversity and an increased dysbiosis.⁴² The microbiome interacts with the host

through production of short-chain fatty acids (including butyrate), induction of the mucosal immune system, stimulation of the local nervous system but also through interaction with the lamina propria by modification of the gut barrier function.⁴³ A dysbiosis in the gut microbiota, characterized by reduction of beneficial bacteria such as *Faecalibacterium prausnitzii*, and Ruminococcaceae and an increase of pathogens or pathobionts, has consistently been shown in patients with IBD, especially ileal CD.⁴⁴ Compared to previous studies we did not find any association between antibiotics treatment in infantile age and IBD, maybe due to that VEO-IBD is partly a different entity from later-onset IBD.^{31 39} On the other hand, we found a two-fold increased risk in offspring to mothers receiving antibiotics during pregnancy. We suggest such antibiotics exposure may be detrimental to the risk of VEO-IBD in the offspring. The gut microbiota in pregnant women resembles that of healthy non-pregnant women during the first two trimesters, but undergoes substantial changes during the third trimester.⁴⁵ In our study, the highest risk of later VEO-IBD was seen in mothers exposed to antibiotics in the last trimester (aHR 2.57 95% CI 1.10-6.01), i.e. just before birth. Recent animal research indicates that antibiotics administered during pregnancy have substantial effects on the offspring microbiome (reduced bacterial diversity), but may also influence the immune response in the offspring⁴⁵ and increase susceptibility to develop colonic inflammation.⁴⁶

While it has long been thought that the intestinal tract is sterile at birth, recent data suggest that the microbial colonization process may be initiated already in utero,⁴⁷ a process that may be affected by antibiotics late in pregnancy. Furthermore, it has been suggested in studies of repeated fecal samples from term infants that the use of intrapartum PcV prophylaxis, to prevent early onset group B streptococcal infection in newborns, alters the offspring microbiome,^{48 49} although others have found very few differences between antibiotic-exposed and non-antibiotic-

exposed infants.⁵⁰ Meanwhile maternal intake of probiotics influence the expression of toll-like receptors in infant meconium,⁵¹ indicating that fetal exposure of antibiotics may play an important role for the development microbiota and the immune system of the child.⁴⁸ While a causative association between antibiotics during pregnancy and VEO-IBD thus seems plausible, we cannot exclude possible confounding from an intrauterine infection with fever of the mother and a raised inflammatory reaction, causing epigenetic imprinting in the fetus and subsequent VEO-IBD. Bernstein and colleagues recently suggested that individuals with IBD were no more likely than controls to have been born to mothers with peripartum infections, however they were diagnosed at an earlier age than those whose mothers did not have an infection.⁵² Unfortunately, we had no data on the indications of the mother's antibiotic treatment during pregnancy or on either symptoms (or date of symptom onset) or genotype in patients, neither were we able to examine the microbiota per se in children with VEO-IBD

In the meta-analysis by Ungaro *et al.* also fluoroquinolones were highly linked to IBD (pooled OR=1.79, 95% CI 1.03-3.12).²¹ During the study period fluoroquinolones were not recommended for younger children in Sweden and only accounted for 0.2% of all antibiotic prescriptions, we therefore chose not to examine fluoroquinolone exposure and IBD separately. The fact that the highest risk estimates for IBD have previously been shown for metronidazole²¹ and fluoroquinolones²¹ (both used in the treatment of IBD, although not currently recommended⁵³) suggest that reverse causation may have been an issue in earlier studies demonstrating an positive association between antibiotics treatment and later IBD.^{31 39} We did not find any association for IBD when we restricted our analysis to antibiotics in the first year of life and our outcome to IBD onset beyond 2 years of age. That sub-analysis is similar to the analysis by Ungaro *et al.*²¹ limiting their dataset to studies with ≥ 1 year of exclusion time between

antibiotics exposure and IBD (HR 1.50, 95% CI 1.44-1.57), while we found an aHR of 1.49 with broad confidence intervals (95% CI 0.69-3.22). Reverse causation is obviously not a concern regarding the possible link between antibiotics exposure during pregnancy and risk of VEO-IBD in the offspring.

Strengths and limitations

The main strength of this study is the nationwide cohort, based on prospective information retrieved from high quality population-based register, thereby eliminating recall bias. In addition, we were able to carry out important sub-analyses such as exclusion of individuals with gastroenteritis prior to their first IBD diagnosis, as well as including a time window between exposure and outcome, analyses that allow us to study the potential influence by information bias. Furthermore, we were able to consider familial factors such as parental history of IBD, parents' country of birth and socioeconomic factors including education level.

We acknowledge a number of limitations. Despite our use of a nationwide register-based cohort of more than 800 000 children, we could only identify 51 cases of IBD in total. Thus, the number of individuals in each subgroup (CD, UC) was quite low, which is reflected by the rather wide confidence intervals and may also be of concern in the adjusted analyses, however crude and adjusted estimates were very similar. Furthermore, we acknowledge the potential difficulties to diagnose UC in this age group, why stratification of patients into UC versus CD may not be completely accurate. This means that even a small number of misclassified patients may have affected our conclusions, since some of the results are of borderline significance, specifically in the sensitivity analyses where we excluded individuals whose potential first episode of IBD could have been misclassified as gastroenteritis. Nevertheless, we believe that the strength of our data lies in the results seen in IBD in general, which is not dependent on

stratification into UC versus CD. Misclassification of a child's IBD diagnosis after maternal exposure to antibiotics in pregnancy is most likely non-differential, whereas it could be differential for exposure in infantile age. The misclassification of father's diagnoses related to maternal exposure during pregnancy and to exposure in infantile age is most likely non-differential, whereas misclassification of mother's diagnoses after exposure during pregnancy could be differential, but most likely non-differential for exposure in infantile age. A non-differential misclassification of the outcome will generally bias towards the null, whereas differential misclassification could lead to both higher and lower risk estimates.

Overall, the limited follow-up time of our study means that we were unable to examine the long-term effect of fetal and early life antibiotics on IBD in adulthood, which may also be one possible explanation to the low number of identified cases and lack of association between antibiotic exposure in childhood and later IBD. While a previous validation of IBD, using the same definition (requiring ≥ 2 diagnoses of IBD) found a positive predictive value (PPV) of 93%,²⁷ we acknowledge that this PPV was calculated in a population of a much higher median age and we cannot rule out that the PPV for ≥ 2 IBD records in young age is different. As data from primary out-patient care are not available in the NPR, the sensitivity of our study could be lower than in real life. However, we believe that the sensitivity of IBD in the NPR is high for children as paediatric IBD patients are managed by hospital-based specialists and closely monitored with visits every three to six months.²⁸ In addition, to minimize the risk of false positive cases and to increase the specificity, we used ≥ 2 diagnoses for our outcome, even though this could mean that we probably excluded some true cases with a lower sensitivity as result. The small number of cases also limited our possibility to perform sibling analysis, which otherwise would have helped us to control for all factors siblings share (both genetic and environmental).

Unfortunately, no population-based data on exposure to antibiotics in inpatient care are available in Sweden today. We have previously shown that 13% of vaginal deliveries are associated with intrapartum exposure to antibiotics.⁵⁴ The corresponding number for elective caesarean section was 14%, and 63% for emergency caesarean section. Thus, while the majority of antibiotics in Sweden (87%) are prescribed in out-patient care⁵⁵, we did not capture those exposed to antibiotics during the immediate intrapartum period, but were able to adjust for mode of delivery as a co-variate. Exposure of antibiotics was furthermore defined as having filled a prescription of antibiotics, which is not equivalent to adherence to treatment.⁵⁶ Finally, the number of children receiving different subtypes of antibiotics was limited, wherefore we were only able to perform stratified analyses on PcV.

Conclusion and implications

In conclusion, we found an association between antibiotics exposure during pregnancy, specifically during the third trimester, but not in infantile age, and subsequent development of VEO-IBD. Our results may indicate that antibiotic exposure in late pregnancy can lead to changes in the microbiome of the child, however further research is needed to confirm our findings. In addition, the absolute risk of VEO-IBD is very low, and antibiotics during pregnancy should still be used when needed.

ACKNOWLEDGEMENTS

We direct our great appreciation to Christina Norrby and Marcus Boman who contributed with excellent data collection and management.

COMPETING INTERESTS

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf. AKÖ, CL, JH, JFL and CA claim no conflict of interest related to the submitted work.

FUNDING

Financial support was provided from the Swedish Research Council through the Swedish Initiative for Research on Microdata in the Social And Medical Sciences (SIMSAM) framework grant no 340-2013-5867, grants provided by the Stockholm County Council (ALF-projects), the Swedish Heart-Lung Foundation and the Swedish Asthma and Allergy Association's Research Foundation.

AUTHOR CONTRIBUTIONS

The study was initiated by JFL and CA, and designed by AKÖ, CL, JFL and CA. AKÖ and CL performed the statistical analysis and wrote the initial draft together with JFL, JH and CA. All authors contributed with invaluable support for data analyses, interpretation of findings and critical revision of the article. CA obtained the financial support. All authors had full access to data, reviewed and approved the final version of the article submitted for publication. AKÖ, CL, JFL, JH and CA are the guarantors for the study and accept full responsibility for the work, had access to the data, and controlled the decision to publish.

REFERENCES

1. Rosen MJ, Dhawan A, Saeed SA. Inflammatory Bowel Disease in Children and Adolescents. JAMA pediatrics 2015;**169**(11):1053-60.
2. Gupta N, Bostrom AG, Kirschner BS, et al. Presentation and disease course in early- compared to later-onset pediatric Crohn's disease. The American journal of gastroenterology 2008;**103**(8):2092-8.
3. Paul T, Birnbaum A, Pal DK, et al. Distinct phenotype of early childhood inflammatory bowel disease. Journal of clinical gastroenterology 2006;**40**(7):583-6.
4. Heyman MB, Kirschner BS, Gold BD, et al. Children with early-onset inflammatory bowel disease (IBD): analysis of a pediatric IBD consortium registry. The Journal of pediatrics 2005;**146**(1):35-40.
5. Bequet E, Sarter H, Fumery M, et al. Incidence and Phenotype at Diagnosis of Very-early-onset Compared with Later-onset Paediatric Inflammatory Bowel Disease: A Population-based Study [1988-2011]. Journal of Crohn's & colitis 2017;**11**(5):519-26.
6. Benchimol EI, Mack DR, Nguyen GC, et al. Incidence, outcomes, and health services burden of very early onset inflammatory bowel disease. Gastroenterology 2014;**147**(4):803-13.e7; quiz e14-5.
7. Satsangi J, Parkes M, Louis E, et al. Two stage genome-wide search in inflammatory bowel disease provides evidence for susceptibility loci on chromosomes 3, 7 and 12. Nat Genet 1996;**14**(2):199-202.
8. Hampe J, Cuthbert A, Croucher PJ, et al. Association between insertion mutation in NOD2 gene and Crohn's disease in German and British populations. Lancet 2001;**357**(9272):1925-8.

- 519 9. Ogura Y, Bonen DK, Inohara N, et al. A frameshift mutation in NOD2 associated with
520 susceptibility to Crohn's disease. *Nature* 2001;**411**(6837):603-6.
- 521 10. Jostins L, Ripke S, Weersma RK, et al. Host-microbe interactions have shaped the genetic
522 architecture of inflammatory bowel disease. *Nature* 2012;**491**(7422):119-24.
- 523 11. Ek WE, D'Amato M, Halfvarson J. The history of genetics in inflammatory bowel disease.
524 *Annals of gastroenterology : quarterly publication of the Hellenic Society of*
525 *Gastroenterology* 2014;**27**(4):294-303.
- 526 12. Zhulina Y, Hahn-Stromberg V, Shamikh A, et al. Subclinical inflammation with increased
527 neutrophil activity in healthy twin siblings reflect environmental influence in the
528 pathogenesis of inflammatory bowel disease. *Inflammatory bowel diseases*
529 2013;**19**(8):1725-31.
- 530 13. Parian A, Limketkai B, Koh J, et al. Appendectomy does not decrease the risk of future
531 colectomy in UC: results from a large cohort and meta-analysis. *Gut* 2017;**66**(8):1390-97.
- 532 14. Harries AD, Baird A, Rhodes J. Non-smoking: a feature of ulcerative colitis. *Br Med J (Clin*
533 *Res Ed)* 1982;**284**(6317):706.
- 534 15. Tysk C, Lindberg E, Jarnerot G, et al. Ulcerative colitis and Crohn's disease in an unselected
535 population of monozygotic and dizygotic twins. A study of heritability and the influence
536 of smoking. *Gut* 1988;**29**(7):990-6.
- 537 16. Andersson RE, Olaison G, Tysk C, et al. Appendectomy is followed by increased risk of
538 Crohn's disease. *Gastroenterology* 2003;**124**(1):40-6.
- 539 17. Andersson RE, Olaison G, Tysk C, et al. Appendectomy and protection against ulcerative
540 colitis. *N Engl J Med* 2001;**344**(11):808-14.
- 541 18. Radford-Smith GL, Edwards JE, Purdie DM, et al. Protective role of appendicectomy on
542 onset and severity of ulcerative colitis and Crohn's disease. *Gut* 2002;**51**(6):808-13.

19. Khalili H, Granath F, Smedby KE, et al. Association Between Long-term Oral Contraceptive Use and Risk of Crohn's Disease Complications in a Nationwide Study. *Gastroenterology* 2016;**150**(7):1561-67.e1.
20. Snapper SB. Very-Early-Onset Inflammatory Bowel Disease. *Gastroenterology & hepatology* 2015;**11**(8):554-6.
21. Ungaro R, Bernstein CN, Gearry R, et al. Antibiotics associated with increased risk of new-onset Crohn's disease but not ulcerative colitis: a meta-analysis. *The American journal of gastroenterology* 2014;**109**(11):1728-38.
22. Ekbom A. The Swedish Multi-generation Register. *Methods Mol Biol* 2011;**675**:215-20.
23. Wettermark B, Hammar N, Fored CM, et al. The new Swedish Prescribed Drug Register--opportunities for pharmacoepidemiological research and experience from the first six months. *Pharmacoepidemiol Drug Saf* 2007;**16**(7):726-35.
24. Ludvigsson JF, Andersson E, Ekbom A, et al. External review and validation of the Swedish national inpatient register. *BMC Public Health* 2011;**11**(1):450.
25. Ludvigsson JF, Otterblad-Olausson P, Pettersson BU, et al. The Swedish personal identity number: possibilities and pitfalls in healthcare and medical research. *European journal of epidemiology* 2009;**24**(11):659-67.
26. Adriaenssens N, Coenen S, Versporten A, et al. European Surveillance of Antimicrobial Consumption (ESAC): outpatient antibiotic use in Europe (1997-2009). *J Antimicrob Chemother* 2011;**66 Suppl 6**:vi3-12.
27. Jakobsson GL, Sternegard E, Olen O, et al. Validating inflammatory bowel disease (IBD) in the Swedish National Patient Register and the Swedish Quality Register for IBD (SWIBREG). *Scandinavian journal of gastroenterology* 2017;**52**(2):216-21.

- 566 28. Ludvigsson JF, Busch K, Olen O, et al. Prevalence of paediatric inflammatory bowel disease
567 in Sweden: a nationwide population-based register study. *BMC gastroenterology*
568 2017;**17**(1):23.
- 569 29. Cnattingius S, Ericson A, Gunnarskog J, et al. A quality study of a medical birth registry.
570 *Scand J Soc Med* 1990;**18**(2):143-8.
- 571 30. Greenland S, Pearl J, Robins JM. Causal diagrams for epidemiologic research. *Epidemiology*
572 (Cambridge, Mass) 1999;**10**(1):37-48.
- 573 31. Card T, Logan RF, Rodrigues LC, et al. Antibiotic use and the development of Crohn's
574 disease. *Gut* 2004;**53**(2):246-50.
- 575 32. Van Kruiningen HJ, Joossens M, Vermeire S, et al. Environmental factors in familial Crohn's
576 disease in Belgium. *Inflammatory bowel diseases* 2005;**11**(4):360-5.
- 577 33. Gearry RB, Richardson AK, Frampton CM, et al. Population-based cases control study of
578 inflammatory bowel disease risk factors. *Journal of gastroenterology and hepatology*
579 2010;**25**(2):325-33.
- 580 34. Shaw SY, Blanchard JF, Bernstein CN. Association between the use of antibiotics in the first
581 year of life and pediatric inflammatory bowel disease. *The American journal of*
582 *gastroenterology* 2010;**105**(12):2687-92.
- 583 35. Han DY, Fraser AG, Dryland P, et al. Environmental factors in the development of chronic
584 inflammation: a case-control study on risk factors for Crohn's disease within New
585 Zealand. *Mutation research* 2010;**690**(1-2):116-22.
- 586 36. Castiglione F, Diaferia M, Morace F, et al. Risk factors for inflammatory bowel diseases
587 according to the "hygiene hypothesis": a case-control, multi-centre, prospective study in
588 Southern Italy. *Journal of Crohn's & colitis* 2012;**6**(3):324-9.

37. Virta L, Auvinen A, Helenius H, et al. Association of repeated exposure to antibiotics with the development of pediatric Crohn's disease--a nationwide, register-based finnish case-control study. *American journal of epidemiology* 2012;**175**(8):775-84.
38. Margolis DJ, Fanelli M, Hoffstad O, et al. Potential association between the oral tetracycline class of antimicrobials used to treat acne and inflammatory bowel disease. *The American journal of gastroenterology* 2010;**105**(12):2610-6.
39. Hviid A, Svanstrom H, Frisch M. Antibiotic use and inflammatory bowel diseases in childhood. *Gut* 2011;**60**(1):49-54.
40. Kronman MP, Zaoutis TE, Haynes K, et al. Antibiotic exposure and IBD development among children: a population-based cohort study. *Pediatrics* 2012;**130**(4):e794-803.
41. Shaw SY, Blanchard JF, Bernstein CN. Association between the use of antibiotics and new diagnoses of Crohn's disease and ulcerative colitis. *The American journal of gastroenterology* 2011;**106**(12):2133-42.
42. Gevers D, Kugathasan S, Denson LA, et al. The treatment-naïve microbiome in new-onset Crohn's disease. *Cell host & microbe* 2014;**15**(3):382-92.
43. Imhann F, Vich Vila A, Bonder MJ, et al. Interplay of host genetics and gut microbiota underlying the onset and clinical presentation of inflammatory bowel disease. *Gut* 2016.
44. Halfvarson J, Brislawn CJ, Lamendella R, et al. Dynamics of the human gut microbiome in inflammatory bowel disease. *Nature microbiology* 2017;**2**:17004.
45. Koren O, Goodrich JK, Cullender TC, et al. Host remodeling of the gut microbiome and metabolic changes during pregnancy. *Cell* 2012;**150**(3):470-80.
46. Munyaka PM, Eissa N, Bernstein CN, et al. Antepartum Antibiotic Treatment Increases Offspring Susceptibility to Experimental Colitis: A Role of the Gut Microbiota. *PloS one* 2015;**10**(11):e0142536.

47. Collado MC, Rautava S, Aakko J, et al. Human gut colonisation may be initiated in utero by distinct microbial communities in the placenta and amniotic fluid. *Scientific reports* 2016;**6**:23129.
48. Nogacka A, Salazar N, Suarez M, et al. Impact of intrapartum antimicrobial prophylaxis upon the intestinal microbiota and the prevalence of antibiotic resistance genes in vaginally delivered full-term neonates. *Microbiome* 2017;**5**(1):93.
49. Azad MB, Konya T, Persaud RR, et al. Impact of maternal intrapartum antibiotics, method of birth and breastfeeding on gut microbiota during the first year of life: a prospective cohort study. *BJOG : an international journal of obstetrics and gynaecology* 2016;**123**(6):983-93.
50. Jauregui F, Carton M, Panel P, et al. Effects of intrapartum penicillin prophylaxis on intestinal bacterial colonization in infants. *Journal of clinical microbiology* 2004;**42**(11):5184-8.
51. Rautava S, Collado MC, Salminen S, et al. Probiotics modulate host-microbe interaction in the placenta and fetal gut: a randomized, double-blind, placebo-controlled trial. *Neonatology* 2012;**102**(3):178-84.
52. Bernstein CN, Burchill C, Targownik LE, et al. Maternal Infections That Would Warrant Antibiotic Use Antepartum or Peripartum Are Not a Risk Factor for the Development of IBD: A Population-Based Analysis. *Inflammatory bowel diseases* 2017;**23**(4):635-40.
53. Gomollon F, Dignass A, Annese V, et al. 3rd European Evidence-based Consensus on the Diagnosis and Management of Crohn's Disease 2016: Part 1: Diagnosis and Medical Management. *Journal of Crohn's & colitis* 2017;**11**(1):3-25.
54. Ahlen KM, Ortqvist AK, Gong T, et al. Antibiotic Treatment and Length of Hospital Stay in Relation to Delivery Mode and Prematurity. *PloS one* 2016;**11**(10):e0164126.

- 636 55. Hellman J O-LB, Bengtsson B, Greko C. SWEDRES-SVARM 2016. Use of antimicrobials
637 and occurrence of antimicrobial resistance in Sweden Secondary SWEDRES-SVARM
638 2016. Use of antimicrobials and occurrence of antimicrobial resistance in Sweden
639 [http://www.sva.se/globalassets/redesign2011/pdf/om_sva/publikationer/swedres_svarm20](http://www.sva.se/globalassets/redesign2011/pdf/om_sva/publikationer/swedres_svarm2016.pdf)
640 [16.pdf](http://www.sva.se/globalassets/redesign2011/pdf/om_sva/publikationer/swedres_svarm2016.pdf).
641 56. Kardas P. Patient compliance with antibiotic treatment for respiratory tract infections. J
642 Antimicrob Chemother 2002;**49**(6):897-903.
643
644

Figure legends and footnotes for figures

Figure 1. A Directed Acyclic Graph (DAG) depicting the included variables in the final models.

Footnote:

The DAG can be applied to various analyses, for example, a study of the association between antibiotic exposure and VEO-IBD. A directed arrow between these two variables indicates that the exposure is associated with the outcome. DAGs can be used to identify all potential confounders that may exist, in order to know which variables that should be adjusted for in the analyses. When the exposure and the outcome are common causes for a third factor (a common effect), this factor is called a collider. Pathways through colliders are closed, unless the collider is adjusted for which will then open the path and potentially cause spurious associations. In this DAG, parental education is a potential collider through parental IBD, however, with adjustment of parental IBD, this backdoor pathway is closed.

* Mode of delivery was only included in analysis of antibiotic exposure in infantile age.

Figure 2. Flow chart of final study population.

Footnote:

*The first estimated conception date, based on date of birth and gestational age in days, is July 1, 2005, and the first child in the cohort was born January 8, 2006. The last estimated conception date was July 7, 2013, and the child was born after 164 days on December 30, 2013.

MBR – Medical Birth Register

MGR – Multi-generation Register

SPDR – Swedish Prescribed Drug Register

LISA – Longitudinal integration database for health insurance and labour market studies

TPR – Total Population Register