



Karolinska  
Institutet

Karolinska Institutet

<http://openarchive.ki.se>

---

This is a Peer Reviewed Accepted version of the following article, accepted for publication in *Allergy*.

2018-11-26

# Allergen-specific IgE over time in women before, during and after pregnancy

Hedman, Anna M; Lundholm, Cecilia; Scheynius, Annika; Alm, Johan; Andolf, Annika; Pershagen, Göran; Almqvist, Catarina

---

*Allergy*. 2019 Mar;74(3):625-628.

Wiley

<http://doi.org/10.1111/all.13662>

<http://hdl.handle.net/10616/46574>

*If not otherwise stated by the Publisher's Terms and conditions, the manuscript is deposited under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.*

## Allergen-specific IgE over time in women before, during and after pregnancy

Anna. M. Hedman, PhD<sup>1</sup>, Cecilia. Lundholm, MSc<sup>1</sup>, Annika. Scheynius, MD, PhD<sup>2,3</sup>, Johan. Alm, MD, PhD<sup>2</sup>, Ellika. Andolf, MD, PhD<sup>4</sup>, Göran. Pershagen, MD, PhD<sup>5,6</sup>, Catarina. Almqvist, MD, PhD<sup>1,7</sup>

### Affiliations:

<sup>1</sup> Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

<sup>2</sup> Department of Clinical Science and Education, Södersjukhuset, Karolinska Institutet, and Unit Sachs' Children and Youth Hospital, Södersjukhuset Stockholm, Sweden

<sup>3</sup> Clinical Genomics, Science for Life Laboratory, Stockholm, Sweden

<sup>4</sup> Department of Clinical Sciences, Danderyd Hospital, Stockholm, Sweden

<sup>5</sup> Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

<sup>6</sup> Centre for Occupational and Environmental Medicine, Stockholm County Council, Stockholm, Sweden

<sup>7</sup> Pediatric Allergy and Pulmonology Unit, Astrid Lindgren Children's Hospital, Karolinska University Hospital, Stockholm, Sweden

20 **Corresponding author:**

21 Anna M. Hedman

22 Department of Medical Epidemiology and Biostatistics

23 PO Box 281, Karolinska Institutet, SE-171 77 Stockholm, Sweden

24 Phone: +46 (0)8-52485285

25 E-mail: [anna.hedman@ki.se](mailto:anna.hedman@ki.se)

26 **Email addresses:** [anna.hedman@ki.se](mailto:anna.hedman@ki.se); [cecilia.lundholm@ki.se](mailto:cecilia.lundholm@ki.se); [annika.scheynius@ki.se](mailto:annika.scheynius@ki.se);

27 [johan.alm@ki.se](mailto:johan.alm@ki.se); [ellika.andolf@sll.se](mailto:ellika.andolf@sll.se); [goran.pershagen@ki.se](mailto:goran.pershagen@ki.se); [catarina.almqvist@ki.se](mailto:catarina.almqvist@ki.se)

28 **Word count:** 1080

29

30 **Summary**

31 The trajectory of IgE levels before, during and after pregnancy in sensitized individuals is  
32 characterized by significant increase in specific IgE to birch allergens but not to other allergens  
33 after multiple testing. This increase may warrant some surveillance in the antenatal care for  
34 those with clinical symptoms.

35 **Keywords:** IgE, change, pregnancy, cohort

36 **Abbreviations:**

37 asIgE = allergen-specific IgE

38 LOQ = limit of quantification

## 39 **To the Editor:**

40 A strong association has been found between IgE reactivity to allergens and asthma, eczema  
41 and rhinitis but these diseases also exist in non-sensitized individuals (1). During pregnancy, IgG,  
42 IgM and IgA have been found to decrease between first and third trimester and to increase  
43 postpartum but the decrease in IgG was mainly due to hemodilution (2). For IgE, studies report  
44 a decrease in total IgE concentration in plasma from pregnancy to after delivery in allergic  
45 mothers (3), however, this decrease was not observed for allergen-specific IgE (asIgE) to  
46 common inhalant allergens (3). In order to tolerate the semiallogeneic fetus a regulated  
47 homeostasis between the Th1/Th2/Th17 (with additional Th9 and Treg) cell subsets and their  
48 dynamic and complex inter-relationship is essential during pregnancy (4, 5) To our knowledge,  
49 no study has included measures of asIgE *before* conception to study the trajectory of asIgE in  
50 sensitized women. Our aim was to investigate changes of asIgE before, during and after  
51 pregnancy.

52 In the prospective longitudinal cohort study Born into Life, we followed 106 women during  
53 2010-2013 living in Stockholm before, during and after pregnancy (6). In total 413 blood  
54 samples were collected at baseline (before conception, n=102), at gestational week 10-14  
55 (n=64) and 26-28 (n=93), at admission to hospital (n=74), and 3 days after delivery (n=80).  
56 Plasma from these blood samples at all time-points were analyzed for IgE-reactivity against a  
57 mixture of 11 inhalant allergens, Phadiatop® and 6 food allergens, fx5® (Thermo Fisher  
58 Scientific, Uppsala, Sweden). If the test screened positive for Phadiatop®  $\geq 0.35$  kU<sub>A</sub>/L, asIgE  
59 antibodies to the most common single airborne allergens; cat, dog, horse-epithelium, birch and

60 timothy were analysed. We did not include house dust mite allergens, since they are not  
61 abundant in Sweden. If screening positive for fx5<sup>®</sup>  $\geq 0.35$  kU<sub>A</sub>/L the sample was subsequently  
62 analysed for asIgE antibodies to the single allergens extract of egg, milk, cod, wheat and soy. To  
63 investigate change over time in each specific allergen (as well as all airborne allergens taken  
64 together), a mixed tobit-model was applied. Correction for multiple testing was made by the  
65 Bonferroni method. Since very few women (n=5) tested positive for fx5<sup>®</sup> we only present  
66 descriptive statistics for these food allergens. Details on study population, data-sources,  
67 variables and statistical analyses are provided in this article's online supporting information. The  
68 Regional Ethics Review Board in Stockholm, Sweden, granted ethical approval. Written informed  
69 consent was obtained from the study participants.

70 In total, n=38 women screened positive for Phadiatop<sup>®</sup>, n=5 women screened positive for fx5<sup>®</sup>,  
71 n=4 women screened positive for both Phadiatop<sup>®</sup> and fx5<sup>®</sup> and n=67 women were negative for  
72 both Phadiatop<sup>®</sup> and fx5<sup>®</sup>. See Table E1 for baseline descriptive in this article's online  
73 supporting information.

74 Figure 1a and Table E2 in this article's online supporting information displays the geometric  
75 mean for IgE airborne allergens concentration over time with 95% Confidence Interval (CI). The  
76 highest geometric mean concentration of asIgE for all time points was found for birch. The  
77 geometric mean concentration of IgE to birch was 0.57 (95% CI 0.39-0.83) kU<sub>A</sub>/L. At baseline,  
78 before conception, the IgE concentration to birch was 0.41 (0.19-0.88) kU<sub>A</sub>/L and 3 days after  
79 delivery 0.72 (0.31-1.70) kU<sub>A</sub>/L. The geometric mean concentration of IgE to cat was 0.21 (0.16-  
80 0.28) kU<sub>A</sub>/L, at baseline 0.21 (0.11-0.40) kU<sub>A</sub>/L and 0.24 (0.09-0.63) kU<sub>A</sub>/L week 10-14 during  
81 pregnancy. Figure 1b shows the specific food allergens over time with 95% CI where IgE to soy

82 displays the highest concentration although low in numbers (max n=5) for all food-allergens,  
83 Table E2.

84 Table 1 shows results from the multivariate mixed tobit model. Levels of IgE to birch increased  
85 significantly over time, with  $\beta=1.006$  (95% CI: 1.002-1.010), which means that by one week  
86 progress in pregnancy the IgE geometric mean concentration increased by 0.6%. This remained  
87 significant after correction for multiple testing using Bonferroni, with similar results for those  
88 with at least one test above 0.10 kU<sub>A</sub>/L. The largest regression coefficient was found when all  
89 airborne allergens were taken together in one analysis,  $\beta=1.007$ , and the smallest was seen for  
90 timothy,  $\beta=1.003$  however non-significant.

91 In summary, we report a small but statistically significant increase over time from  
92 preconception to postpartum for asIgE to birch allergen, but no changes for other allergens. This  
93 finding is in line with previous findings that birch-induced cytokine levels were increased during  
94 pregnancy compared to post-partum in sensitized women with allergic symptoms (7). While  
95 they measured Th2-like cytokines and we assessed levels of asIgE, Th2-like cytokines and IgE has  
96 been shown to be related in a series of cellular reactions (5). Th2 cells are suggested to be the  
97 key T-helper cell subset responsible for allergic disease (5), and normal pregnancies have been  
98 described as an adequate balance for Th1/Th2 immunity which is slightly shifted to Th2-type  
99 immunity with additional regulation of the Th17 and Treg cells (4). Thus, the Th2-like immunity  
100 associated with allergic disease could be beneficial for the maintenance of a successful  
101 pregnancy. On the other hand, pregnancy might enhance the already Th2-skewed immunity of  
102 allergic women, expose the foetus to a strong Th2 environment, and perhaps lead to allergy  
103 development later in life (8). Since there is a strong correlation between sensitization and

104 allergic symptoms, our findings may warrant some surveillance of allergic women with clinical  
105 symptoms in the antenatal care.

106 The strength of this study is the longitudinal design with repeated measurements before  
107 conception, during pregnancy and after delivery. In addition, our study population was very  
108 homogeneous regarding age, geography and education. We also controlled for season, which is  
109 particularly important for the specific birch and timothy allergens at this geographical latitude,  
110 and we applied multiple test correction, making type I error less likely. The limitations include a  
111 small cohort size resulting in low statistical power. Normal aging could be an issue, however,  
112 findings from the follow-up by European Community Respiratory Health Survey rather showed  
113 that aging was associated with decreased levels of sensitization (9). Methodological issues  
114 relating to the analyses of biological samples could affect the results although the analyses were  
115 all performed on the same instrument, randomized and using the same batch reagents.  
116 Furthermore, being sensitized to birch is very common in this region and we can not rule out  
117 that there is also an effect for the other allergens, including timothy, although not statistically  
118 significant due to lower power.

119 In conclusion, we found a small but statistically significant increase in IgE levels to birch  
120 allergens in women over time before, during and after pregnancy but no statistically significant  
121 change in levels of asIgE for other inhalant allergens or food allergens.

122

123 *Anna M. Hedman, PhD<sup>1</sup>*

124 *Cecilia Lundholm, MSc*<sup>1</sup>

125 *Annika Scheynius, MD, PhD*<sup>2,3</sup>

126 *Johan Alm, MD, PhD*<sup>2</sup>

127 *Ellika Andolf, MD, PhD*<sup>4</sup>

128 *Göran Pershagen, MD, PhD*<sup>5,6</sup>

129 *Catarina Almqvist, MD, PhD*<sup>1,7</sup>

130 <sup>1</sup> Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm,  
131 Sweden.

132 <sup>2</sup> Department of Clinical Science and Education, Södersjukhuset, Karolinska Institutet, and Unit  
133 Sachs' Children and Youth Hospital, Södersjukhuset Stockholm, Sweden

134 <sup>3</sup> Clinical Genomics, Science for Life Laboratory, Stockholm, Sweden

135 <sup>4</sup> Department of Clinical Sciences, Danderyd Hospital, Stockholm, Sweden

136 <sup>5</sup> Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

137 <sup>6</sup> Centre for Occupational and Environmental Medicine, Stockholm County Council, Stockholm,  
138 Sweden

139 <sup>7</sup> Pediatric Allergy and Pulmonology Unit, Astrid Lindgren Children's Hospital, Karolinska  
140 University Hospital, Stockholm, Sweden

141 **Funding sources:** The Swedish Research Council, partly through the Swedish initiative for

142 Research on Microdata in the Social And Medical Sciences (SIMSAM) framework grant no 340-



143 2013-5867, grants provided by the Stockholm County Council (ALF-projects), the Strategic  
144 Research Program in Epidemiology at Karolinska Institutet, the Department of Clinical Sciences  
145 at Danderyd Hospital, the Swedish Medical Research Council, the Swedish Heart-Lung  
146 Foundation, and the Swedish Asthma and Allergy Association's Research Foundation.

147 **Competing financial interests:** The authors declare no financial conflict of interest

148 **Author Contributions:** CA, GP and EA wrote the proposal, AH analysed the data and drafted the  
149 paper, CL helped with analysis. AS, JA, EA, GP, CA and CL helped in study conception and  
150 designing, data interpretation and gave significant inputs in revising the manuscript. All authors  
151 read and approved the final manuscript.

152

## 153 **References**


- 154 1. WAO. White book on allergy. Milwaukee, Wisconsin, United States of America: World Allergy  
155 Association (WAO),2011.
- 156 2. Ailus KT. A follow-up study of immunoglobulin levels and autoantibodies in an unselected  
157 pregnant population. *Am J Reprod Immunol*. 1994;**31**:189-96.
- 158 3. Sandberg M, Frykman A, Jonsson Y, Persson M, Ernerudh J, Berg G, et al. Total and allergen-  
159 specific IgE levels during and after pregnancy in relation to maternal allergy. *J Reprod Immunol*.  
160 2009;**81**:82-8.
- 161 4. Saito S, Nakashima A, Shima T, Ito M. Th1/Th2/Th17 and regulatory T-cell paradigm in  
162 pregnancy. *Am J Reprod Immunol*. 2010;**63**:601-10.
- 163 5. Berker M, Frank LJ, Gessner AL, Grassl N, Holtermann AV, Hoppner S, et al. Allergies - A T cells  
164 perspective in the era beyond the TH1/TH2 paradigm. *Clin Immunol*. 2017;**174**:73-83.
- 165 6. Smew AI, Hedman AM, Chiesa F, Ullemar V, Andolf E, Pershagen G, et al. Limited association  
166 between markers of stress during pregnancy and fetal growth in 'Born into Life', a new prospective birth  
167 cohort. *Acta Paediatr*. 2018.
- 168 7. Abelius MS, Jedenfalk M, Ernerudh J, Janefjord C, Berg G, Matthiesen L, et al. Pregnancy  
169 modulates the allergen-induced cytokine production differently in allergic and non-allergic women.  
170 *Pediatr Allergy Immunol*. 2017;**28**:818-24.
- 171 8. Abelius MS, Lempinen E, Lindblad K, Ernerudh J, Berg G, Matthiesen L, et al. Th2-like chemokine  
172 levels are increased in allergic children and influenced by maternal immunity during pregnancy. *Pediatr  
173 Allergy Immunol*. 2014;**25**:387-93.

174 9. Amaral AFS, Newson RB, Abramson MJ, Anto JM, Bono R, Corsico AG, et al. Changes in IgE  
175 sensitization and total IgE levels over 20 years of follow-up. *J Allergy Clin Immunol*. 2016;**137**:1788-95.e9.

176

177

178



179  
 180 w = week during pregnancy  
 181 3 d = 3 days after delivery

182  
 183 **Figure 1a-b. IgE-reactivity to airborne (a) and food (b) allergens over time in women pre,**  
 184 **during and post pregnancy, presented as geographic means with 95% Confidence Intervals.**

185

186

187

188 **Table 1. Regression coefficients for time from mixed tobit model with 95% Confidence**  
 189 **Intervals.**

Allergen	All				At least one above 0.10 kU <sub>A</sub> /L			
	$\beta$	95% CI		p-value	$\beta$	95% CI		p-value
Airborne allergens (all)	1.007	0.991	1.024	0.388	1.006	0.993	1.020	0.346
Cat	1.005	0.998	1.012	0.136	1.005	0.998	1.012	0.142
Horse	1.005	0.998	1.011	0.160	1.005	0.998	1.011	0.170
Dog	1.004	0.998	1.009	0.208	1.004	0.998	1.009	0.215
Birch	<b>1.006</b>	<b>1.002</b>	<b>1.010</b>	<b>0.004</b>	<b>1.006</b>	<b>1.002</b>	<b>1.010</b>	<b>0.004</b>
Timothy	1.003	0.999	1.008	0.143	1.003	0.999	1.008	0.140

$\beta$  = Exponentiated regression coefficient for time (weeks) from analysis on log-transformed data

p-value cut-off with Bonferroni correction = 0.008

190

## 1 **Online Supporting Information**

### 2 **Study Population**

3 Born into Life originates from the larger LifeGene study which has been described in detail  
4 elsewhere (1). In short, LifeGene included index persons between 18-45 years who were also  
5 encouraged to invite their household members. At enrollment each participant was invited to a  
6 test center where biosamples were taken as well as physical measurements. A web-based  
7 questionnaire based on multifaceted questions regarding health, diet, lifestyle and diseases was  
8 administered at baseline and annually thereafter with a shorter version. Women who became  
9 pregnant in LifeGene in Stockholm during 2010-2013 were recruited to Born into Life. The  
10 women answered a web-based questionnaires at gestational week 10-14 and 26-18 regarding  
11 health, diseases, lifestyle and pregnancy.

### 12 **Blood samples and analyses of IgE-reactivity to allergens**

13 Blood samples from 106 women were collected before conception, at gestational week 10-14  
14 and 26-28, at admission to the hospital for delivery (cord-blood), and 3 days after delivery in  
15 conjunction with the Phenylketonuria screening of the child (2). A positive screening test for  
16 either common inhalant allergens (Phadiatop®) or food allergens (fx5®) is fulfilled if the  
17 concentration is  $\geq 0.35$  kU<sub>A</sub>/L. A total of n=155 tests from 38 women screened positive for  
18 Phadiatop® and n=15 tests from 5 women screened positive for fx5®. Based on this information  
19 we selected samples to analyse allergen-specific IgE (asIgE). We selected the samples from  
20 women who screened positive for one test and had at least one other test at another time-point

21 (regardless if this test was screened positive or not), in total n=131 tests from 33 women  
22 regarding airborne allergens and n=19 tests from 5 women regarding food allergens. These  
23 specific allergens were reported as continuous values, from <0.10 to >100 kU<sub>A</sub>/L, i.e. the lower  
24 limit of quantification (LOQ) was 0.10 kU<sub>A</sub>/L. All samples were analyzed at the Centre for Child  
25 Research, Södersjukhuset, Stockholm, Sweden.

## 26 **Data sources and variables**

27 Maternal age was retrieved from the medical birth records for each mother and child. Highest  
28 attained educational level (from mandatory secondary school to high school, university or  
29 other) was derived from LifeGene questionnaires. Educational level was used as a proxy for  
30 socioeconomic status (3). The covariate season of the year was coded as 1, if the sample was  
31 taken during the pollen season of the year in this geographical region (i.e., May-June-July) and 0  
32 otherwise (i.e., all other months).

## 33 **Statistical analyses**

34 Baseline descriptive statistics included frequencies and percentages for count variables. Chi-  
35 squares with Fisher's exact test were applied to the categorical variables. All data were initially  
36 analyzed for normality and extreme values. Due to positively skewed distribution of the IgE  
37 concentration, the values of the IgE were logarithmically transformed. For the descriptive  
38 statistics, samples less than LOQ (i.e., <0.10 kU<sub>A</sub>/L) were exchanged with fill values equal to  
39 LOQ/√2.

40 The tobit model is a regression model for truncated data method, analyzing laboratory  
41 measurements subject to detection or quantification limits (i.e., where non-detects are values  
42 below a given detection limit, which may not be observed, in this case from zero to 0.10 kU<sub>A</sub>/L)  
43 (4). In the mixed tobit model we assume an intercept,  $\alpha_i$ , for each woman, where  $\alpha_i \sim N(0, \sigma^2)$   
44 and that data are missing at random. Our explanatory variable was time, measured in weeks.  
45 The analyses were adjusted for season (at which season the measurement was taken), since it  
46 has been found that seasonal exposure to pollen influences IgE antibody levels in allergic  
47 individuals (5). For ease of interpretation, the regression coefficients ( $\log\beta$ ) and the  
48 corresponding CIs were exponentiated back to the original scale. Thus  $(1-\beta) \times 100$  can be  
49 interpreted percentage increase in asIgE geometric mean concentration associated with one  
50 week progression in pregnancy. We first included all data (everyone above the cut-off  $\geq 0.35$   
51 kU<sub>A</sub>/L for Phadiatop<sup>®</sup> and fx5<sup>®</sup>) in our analyses and then did sensitivity analyses using subjects  
52 with at least one test above the lower limit of quantification (LOQ) for each specific allergen  
53 (i.e.,  $\geq 0.10$  kU<sub>A</sub>/L.) Data management was performed using SAS 9.4 (SAS Institute, Cary, NC,  
54 USA) and data analyses was conducted using STATA/IC 15.0 for Windows (StataCorp LLC,  
55 College Station, TX, USA).

56

57 1. Almqvist C, Adami HO, Franks PW, Groop L, Ingelsson E, Kere J, et al. LifeGene--a large  
58 prospective population-based study of global relevance. *European Journal of Epidemiology*. 2011;**26**:67-  
59 77.

60

61 2. Smew AI, Hedman AM, Chiesa F, Ullemar V, Andolf E, Pershagen G, et al. Limited association  
62 between markers of stress during pregnancy and fetal growth in 'Born into Life', a new prospective birth  
63 cohort. *Acta Paediatrica*. 2018.

64

- 65 3. Winkleby MA, Jatulis DE, Frank E, Fortmann SP. Socioeconomic status and health: how  
66 education, income, and occupation contribute to risk factors for cardiovascular disease. *American*  
67 *Journal of Public Health*. 1992;**82**:816-20.  
68
- 69 4. Lubin JH, Colt JS, Camann D, Davis S, Cerhan JR, Severson RK, et al. Epidemiologic evaluation of  
70 measurement data in the presence of detection limits. *Environmental Health Perspectives*.  
71 2004;**112**:1691-6.  
72
- 73 5. Lagier B, Pons N, Rivier A, Chanal I, Chanez P, Bousquet J, et al. Seasonal variations of interleukin-  
74 4 and interferon-gamma release by peripheral blood mononuclear cells from atopic subjects stimulated  
75 by polyclonal activators. *Journal of Allergy and Clinical Immunology*. 1995;**96**:932-40.
- 76



## Online Supporting Information

**Table E1. Baseline descriptive characteristics of the cohort.**

	Phadiatop® negative and fx5® negative n=67	Phadiatop® positive n=38 (>0.35 kU <sub>A</sub> /L)	fx5® positive n=5 (>0.35 kU <sub>A</sub> /L)
Age (years)	n(%)	n (%)	n(%)
<19	0	0	0
20-24	1 (1.5)	1 (2.6)	0
25-29	9 (13.4)	11 (29.0)	1 (20.0)
29-34	36 (53.7)	17 (44.7)	2 (40.0)
>34	21 (31.3)	8 (21.1)	2 (40.0)
missing	0	1 (2.6)	0
Education (years)			
<9	0	0	0
10-12	4 (6.0)	5 (13.2)	1 (20.0)
>13	59 (88.1)	27 (71.1)	3 (60.0)
other	2 (3.0)	1 (2.6)	1 (20.0)
missing	2 (3.0)	5 (13.2)	0

n = nr of participants

**Table E2. Quantification of the specific allergens with geometric mean, 95% confidence interval and number of individuals.**

Allergen	Geometric mean (95% CI)	n	Baseline	n	Week 10-14	n	Week 26-28	n	Admission for delivery	n	3 days after delivery	n
<b>Cat</b>	0.21 (0.16-0.28)	131	0.21 (0.11-0.40)	31	0.24 (0.09-0.63)	17	0.22 (0.12-0.41)	32	0.17 (0.10-0.29)	25	0.21 (0.11-0.40)	26
<b>Horse</b>	0.13 (0.11-0.16)	131	0.13 (0.08-0.19)	31	0.17 (0.09-0.34)	17	0.13 (0.09-0.19)	32	0.11 (0.08-0.16)	25	0.14 (0.09-0.22)	26
<b>Dog</b>	0.17 (0.13-0.21)	131	0.15 (0.09-0.25)	31	0.25 (0.11-0.58)	17	0.16 (0.10-0.26)	32	0.14 (0.08-0.24)	25	0.18 (0.10-0.31)	26
<b>Birch</b>	0.57 (0.39-0.83)	131	0.41 (0.19-0.88)	31	0.78 (0.26-2.38)	17	0.45 (0.21-0.97)	32	0.71 (0.30-1.70)	25	0.72 (0.31-1.70)	26
<b>Timothy</b>	0.44 (0.32-0.62)	127	0.37 (0.19-0.74)	30	0.40 (0.15-1.05)	17	0.36 (0.19-0.70)	31	0.51 (0.23-1.12)	23	0.63 (0.30-1.35)	26
<b>Egg</b>	0.11 (0.09-0.14)	19	0.14 (0.07-0.26)	5	0.12 (0.07-0.19)	3	0.10 (0.07-0.16)	5	0.09 (0.06-0.15)	3	0.09 (0.06-0.15)	3
<b>Milk</b>	0.13 (0.09-0.18)	19	0.13 (0.06-0.27)	5	0.16 (0.07-0.36)	3	0.13 (0.06-0.28)	5	0.11 (0.04-0.29)	3	0.10 (0.05-0.23)	3
<b>Cod</b>	0.07 (0.07-0.07)	19	0.07 (0.07-0.07)	5	0.07 (0.07-0.07)	3	0.07 (0.07-0.07)	5	0.07 (0.07-0.07)	3	0.07 (0.07-0.07)	3
<b>Wheat</b>	0.25 (0.16-0.39)	19	0.20 (0.08-0.46)	5	0.46 (0.27-0.81)	3	0.23 (0.09-0.60)	5	0.25 (0.06-1.02)	3	0.24 (0.06-1.07)	3
<b>Soy</b>	0.23 (0.10-0.52)	18	0.19 (0.03-1.36)	4	0.37 (0.04-3.87)	3	0.18 (0.04-0.75)	5	0.25 (0.02-3.15)	3	0.25 (0.02-2.95)	3

CI=Confidence Interval

n =number of individuals