

Karolinska Institutet http://openarchive.ki.se

This is a Peer Reviewed Accepted version of the following article, accepted for publication in European Journal of Epidemiology.

2017-03-30

No association between macrolide treatment in infancy and later pyloric stenosis in Sweden

Ludvigsson, Jonas F; Lundholm, Cecilia; Örtqvist, Anne K; Almqvist, Catarina

Eur J Epidemiol. 2016 Mar;31(3):331-2. http://doi.org/10.1007/s10654-015-0114-6 http://hdl.handle.net/10616/45660

If not otherwise stated by the Publisher's Terms and conditions, the manuscript is deposited under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (http://creativecommons.org/licenses/by-nc-nd/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way. Copyright © 2015 Springer Science Business Media. This manuscript is subject to Springer Nature's Terms of Use, which permit users to view, print, copy, download and text and data-mine the content, for the purposes of academic research, subject always to the full conditions of use. Under no circumstances may the manuscript be shared or distributed under a Creative Commons, or other form of open access license, nor may it be reformatted or enhanced.

Title: No association between macrolide treatment in infancy and later pyloric stenosis in Sweden

Jonas F Ludvigsson, MD, PhD^{1,2} Cecilia Lundholm, MSc¹ Anne K. Örtqvist, MD, PhD¹ Catarina Almqvist, MD, PhD^{1,3}

 ¹ Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
² Department of Pediatrics, Örebro University Hospital, Örebro University, Örebro, Sweden
³ Astrid Lindgren Children's Hospital, Karolinska University Hospital, Stockholm, Sweden

Corresponding author:

Jonas F. Ludvigsson, Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, 171 77 Stockholm, Sweden Phone: +46 (0) 8-5248 2356 Fax: +46 (0) 8-31 49 75 Email: jonasludvigsson@yahoo.com

Word count (introduction to discussion): 642 Date of this version: November 24, 2015

ETHICAL APPROVAL

The study was approved by the Regional Ethical Review board in Stockholm, Sweden (DNR 2011/2012-31/5 and 2012/1179-32).

COMPETING INTERESTS

All authors have completed the ICMJE uniform disclosure form at <u>www.icmje.org/coi_disclosure.pdf</u>. JFL, CL, AKÖ, and CA claim no conflict of interest related to the submitted work.

FUNDING None.

Dear EJE,

In a recent paper in the BMJ, Lund *et al* reported an increased risk of infantile hypertrophic pyloric stenosis (IHPS) in infants exposed to macrolides, especially during days 0-13 (adjusted rate ratio (ARR)=29.8, 95%CI=16.4-54.1)¹. The excess risk decreased rapidly with exposure later in life (days 14-120: ARR=3.24; 95%CI=1.20-8.74). Also a retrospective cohort study of children to US uniformed personnel recorded in the TRICARE database found a positive association between macrolides and IHPS². In Sweden, one of the most common uses of erythromycin in newborns is to stimulate the gastrointestinal motility and we believe that the positive association between macrolides and IHPS is due to reverse causation ³. Lund *et al* also mention that some macrolides are used for dysmotility in smaller children.

In a nationwide cohort of 582,494 children born between July 2005 and December 2010 in Sweden we examined the use of macrolides and the risk of IHPS.

METHODS

Data on pregnancy and birth factors were obtained from the Swedish Medical Birth Registry⁴, while we used the Swedish Patient Registry⁵ (contains data on hospitalbased outpatient and inpatient care since 2001) to identify cases of IHPS (international classification of disease (ICD) codes; Q40.0, K31.1). We defined exposure as macrolide use (ATC code: J01FA) according to the Swedish Prescribed Drug Registry⁶. In Sweden, the Prescribed Drug Registry contains dispensed drug data but not data on over the counter drugs or drugs administered at hospital. A priori we had planned to use Cox regression to examine the risk of IHPS in children exposed to macrolides.

RESULTS

Four hundred and fifty (450) children had a diagnosis of IHPS (equivalent to 0.8 per 1000 births). The median age at diagnosis was 35 days, ranging from birth to 1038 days (four children had their first recorded diagnosis of IHPS beyond the age of 500 days).

Some 17,659 children had a record of macrolides, but only 32 had been prescribed macrolides (all had received erythromycin) during days 0-13 (the youngest was 5 days old) and 1275 children during days 14-120. In total 240 children had a record of macrolides by day 35 (238 (99%) with erythromycin). No infant in our study had a record of macrolides prior to IHPS, but 18 children had received macrolides after the IHPS diagnosis. No case among 240 exposed corresponds to a proportion of 0% with a 95% confidence interval of 0-1.5% based on the binomial distribution.

DISCUSSION

Assuming a risk ratio of 30 for the association between macrolides during day 0-13 and IHPS, there would be a probability of 47% to find no cases among 32 exposed, based on the IHPS incidence in Sweden. Thus our data do not contradict the findings of Lund *et al*, and we cannot rule out that early macrolide exposure contributes to IHPS, especially when administered in high doses. Besides, a major weakness of our study is that we did not have access to antibiotics administered in hospital. For newborns, hospital-administered antibiotics are an important exposure. Still, we were able to identify 240 children with macrolides in the first 35 days of life and none developed IHPS. The lack of IHPS cases among individuals exposed to macrolides meant that we were unable to adjust our data for sex, firstborn status, and prematurity.

While the lack of individuals with IHPS and prior macrolides does not allow us to calculate any risk estimate for the association with IHPS it strengthens our belief that macrolides is a rare cause of IHPS. None of the 240 children had a record of IHPS. Assuming the risk ratios estimated by Lund *et al* are true, the population attributable fraction based on Swedish macrolide prescription frequencies would be 0.6%, corresponding to approximately one case every 4th year.

In conclusion, the current study found no association between macrolide exposure and later IHPS, but our results were based on small numbers of exposed individuals.

REFERENCES

- 1. Lund M, Pasternak B, Davidsen RB, et al. Use of macrolides in mother and child and risk of infantile hypertrophic pyloric stenosis: nationwide cohort study. *BMJ*. 2014;348:g1908.
- 2. Eberly MD, Eide MB, Thompson JL, Nylund CM. Azithromycin in early infancy and pyloric stenosis. *Pediatrics*. 2015;135(3):483-488.
- 3. Ortqvist AK, Lundholm C, Kieler H, et al. Antibiotics in fetal and early life and subsequent childhood asthma: nationwide population based study with sibling analysis. *Bmj.* 2014;349:g6979.
- 4. Cnattingius S, Ericson A, Gunnarskog J, Kallen B. A quality study of a medical birth registry. *Scand J Soc Med.* 1990;18(2):143-148.
- 5. Ludvigsson JF, Andersson E, Ekbom A, et al. External review and validation of the Swedish national inpatient register. *BMC Public Health*. 2011;11(1):450.
- 6. Wettermark B, Hammar N, Fored CM, et al. The new Swedish Prescribed Drug Register--opportunities for pharmacoepidemiological research and experience from the first six months. *Pharmacoepidemiol Drug Saf.* 2007;16(7):726-735.