


Thesis for doctoral degree (Ph.D.)
2010

Thesis for doctoral degree (Ph.D.) 2010

Glutamate modulates temporomandibular joint bone tissue resorption in patients with early rheumatoid arthritis



Glutamate modulates temporomandibular joint bone tissue resorption in patients with early rheumatoid arthritis

Anna-Kari Hajati



Karolinska
Institutet

200
1810 – 2010 *Years*



Karolinska
Institutet

200
1810 – 2010 *Years*

Anna-Kari Hajati

Division of Clinical Oral Physiology
DEPARTMENT OF DENTAL MEDICINE
Karolinska Institutet, Stockholm, Sweden

Glutamate modulates temporomandibular joint bone tissue resorption in patients with early rheumatoid arthritis

Anna-Kari Hajati



**Karolinska
Institutet**

200
1810 – 2010 *Years*

Stockholm 2010

Cover image

Three-dimensional renderings of a temporomandibular joint condyle processed from cone-beam computerized tomography scans at the time for RA diagnosis and at one-year and two-year follow-up. The first illustration is a rendering of the condyle at the time for diagnosis and the others are visualizations of bone tissue changes showing bone resorption in blue, bone formation in red and no change in white. In this particular condyle bone tissue resorption was seen after one year and new bone formation was identified between the one-year and two-year examinations.

In collaboration with: Departments of Orthodontics and Computer Sciences, University of North Carolina, Chapel Hill, NC, USA.

Processed by: Felipe de Assis Robeiro Carvalho, doctoral student in orthodontics and Beatriz Paniagua, postdoctoral fellow in Computer sciences.

All previously published papers are reprinted with permission from the publisher.

Published by Karolinska Institutet.

© Anna-Kari Hajati, 2010

Printed by



www.reproprint.se

Gårdsvägen 4. 169 70 Solna

ISBN

978-91-7409-954-6

To Alexandra and Reza

How success is measured is different for each of us.
The first and most important step is realizing that the secret of success is inside.

Garry Kasparov

Main supervisor

Sigvard Kopp, Professor, Karolinska Institutet, Department of Dental Medicine, Division of Clinical Oral Physiology, Huddinge, Sweden

Supervisor

Per Alstergren, Associate Professor, Karolinska Institutet, Department of Dental Medicine, Division of Clinical Oral Physiology, Huddinge, Sweden

External mentor

Fredrik Bergstrand, DDS, Specialist in orthodontics. Professional Services Manager, 3M Unitek Corporation, Monrovia , USA

Opponent

Thomas Klit Pedersen, Associate Professor, University of Aarhus, Faculty of Health Sciences, School of Dentistry, Aarhus, Denmark.

Examining committee

EwaCarin Ekberg (chairman), Associate Professor, Malmö University, Faculty of Odontology, Department of Stomatognathic Physiology, Malmö, Sweden

Ingiäld Hafström, Professor, Karolinska Institutet, Department of Medicine, Karolinska University Hospital, Huddinge, Sweden

Manuel Patarroyo, Professor, Karolinska Institutet, Department of Dental Medicine, Division of Oral Biology, Huddinge, Sweden

Contents

| | |
|------------------------------------|----|
| Contents | 6 |
| Summary | 7 |
| Preface | 10 |
| Abbreviations | 11 |
| Introduction | 12 |
| Aims | 21 |
| Materials and methods | 23 |
| Results and discussion | 37 |
| General discussion | 49 |
| Conclusions | 52 |
| Populärvetenskaplig sammanfattning | 54 |
| Acknowledgements | 56 |
| References | 58 |

Summary

The general aim was to investigate the occurrence and development of TMJ bone tissue resorption and to identify predictors for progression during the two first years with rheumatoid arthritis.

The thesis comprises one cross-sectional part and one longitudinal part. The aim of the cross-sectional part was to investigate whether there was an association between the radiographic sign (TMJ erosions) or the clinical sign (crepitus) of TMJ bone tissue resorption and glutamate and if that relationship was influenced by serotonin, tumor necrosis factor, interleukin-1 β , interleukin-6, vascular endothelial growth factor, C-reactive protein, erythrocyte sedimentation rate, rheumatoid factor or sex hormones. The first aim of the longitudinal part was to investigate whether TMJ erosions or crepitus, glutamate, inflammatory mediators and markers changed over time and if the change in TMJ erosions was related to these factors. The second aim was to investigate the predictive value of TMJ erosions, glutamate and crepitus as well as the inflammatory mediators and markers for change in TMJ erosions over two years after diagnosis.

Both parts of the thesis comprised the same group of patients with recent RA diagnosis. They were 47 at first visit and 41 attended the two year follow-up. Similar examination protocols were used for the cross-sectional and the longitudinal part. The subjects were examined for TMJ erosions on cone-beam computerized tomographs and the clinical examination included the articular sound of crepitus. Blood samples were obtained and analyzed for glutamate, serotonin, tumor necrosis factor, interleukin-1 β , interleukin-6 and vascular endothelial growth factor, C-reactive protein, erythrocyte sedimentation rate and rheumatoid factor, as well as for estradiol and testosterone.

Significance of univariate correlations was calculated with Spearman rank correlation coefficient. To test the significance of the differences between visits the Friedman ANOVA on ranks was used. Multivariate regression was used to investigate the relative influence by independent variables on the presence, development or prediction of the TMJ erosions. Interaction analysis was performed by including an interaction term in addition to the main effects in the multiple linear regression model. The significance of the differences between groups was calculated with Fisher's exact test for variables on nominal scale and Mann-Whitney

U-test for variables on ordinal scale. A probability level of less than 0.05 was considered significant in all analyses.

Seventy-two percent of the patients showed TMJ erosions at the beginning of the study. TMJ erosion score was positively correlated to both glutamate and crepitus. There was also a positive correlation between erosions and C-reactive protein in the patients with estradiol levels ≥ 65 pmol/L. Multiple stepwise regression showed that TMJ crepitus ($p < 0.001$), glutamate ($p = 0.005$) and erythrocyte sedimentation rate ($p = 0.010$) explained 41% of the variation in the dependent variable TMJ bone erosion score ($p < 0.001$). Glutamate was positively correlated to TMJ erosion score in the patients without TMJ crepitus, with C-reactive protein < 3 mg/L, C-reactive protein < 3 mg/L and estradiol < 50 pmol/L as well as C-reactive protein < 3 mg/L and testosterone ≤ 1.2 pmol/L. Progression of the erosion score was found in 29% of the patients and 43% showed regression during the first two years after RA diagnosis. The progression of erosion score was explained to 34% by testosterone ($p = 0.012$) and interleukin- 1β ($p = 0.024$). Patients with higher than median level of glutamate at the first visit showed more reduction in erosion score than those with lower level ($p = 0.035$). TMJ erosion score at the first visit was negatively correlated to progression of the score ($r_s = -0.64$, $p < 0.001$). The multivariate analysis showed that erosion score was the only significant predictive factor that explained 53% of the regression ($p < 0.001$).

TMJ bone tissue resorption was found to be present in a majority of patients with early RA. Glutamate modulates TMJ bone tissue resorption in the early phase of disease with absence of crepitus and under the influence by systemic inflammation and sex hormones. Regression in TMJ bone tissue resorption was the most frequent finding over a two-year period directly after diagnosis. Progression however seems to be associated with both inflammatory and non-inflammatory mechanisms while progression of crepitus is associated with inflammation when glutamate is increased. Regression of TMJ bone tissue resorption could be predicted by early TMJ erosions and glutamate. The results indicate that TMJ bone tissue resorption can be expected early in RA.

Keywords

Arthritis, rheumatoid; Bone tissue resorption; Cone-beam computed tomography; C-reactive-protein; Erythrocyte sedimentation rate; Estradiol; Glutamate; Interleukin-1 β ; Interleukin-6; Rheumatoid factor; Serotonin; Temporomandibular joint; Testosterone; Tumor necrosis factor; Vascular endothelial growth factor.

Preface

This thesis is based on the following manuscripts, which will be referred to in the text by their Roman numerals:

- I. Endogenous glutamate in association with inflammatory and hormonal factors modulates bone tissue resorption of the temporomandibular joint in patients with early rheumatoid arthritis.

Hajati A-K, Alstergren P, Näsström K, Bratt J and Kopp S.

J Oral Maxillofac Surg. 2009 Sep;67(9):1895-903.

- II. Temporomandibular joint bone tissue resorption in patients with early rheumatoid arthritis can be predicted by joint crepitus and plasma glutamate level.

Hajati A-K, Alstergren P, Näsström K, Bratt J and Kopp S.

Mediators of Inflammation. 2010:627803.

- III. Bone tissue resorption in the temporomandibular joint and its relation to systemic inflammation and crepitus during the first two years of rheumatoid arthritis.

Hajati A-K, Alstergren P, Näsström K, Bratt J and Kopp S.

Manuscript.

- IV. Temporomandibular joint bone tissue resorption and plasma glutamate predict regression of articular bone tissue resorption in two years for patients with early diagnosed rheumatoid arthritis.

Hajati A-K, Alstergren P, Näsström K, Bratt J and Kopp S.

Manuscript.

Abbreviations

| | |
|--------------|------------------------------------------------------------|
| 5-HT | Serotonin (5-hydroxytryptamine) |
| BARFOT | Better Anti-Rheumatic Farmacotherapy |
| AMPA | α -amino-3-hydroxyl-5-methyl-4-isoxazole-propionate |
| ANOVA | Analysis of variance |
| CT | Computerized tomography |
| DAS28 | Disease activity score 28 |
| DMARD | Disease modifying anti-rheumatic drug |
| EIA | Enzyme immuno-assay |
| GC | Glucocorticosteroids |
| VEGF | Vascular endothelial growth factor |
| NMDA | N-methyl-D-aspartate |
| NSAID | Non-steroidal anti-inflammatory drug |
| OPG | Osteoprotegerin |
| Glu | Glutamate |
| IL-1 β | Interleukin-1beta |
| IL-6 | Interleukin-6 |
| TNF | Tumor necrosis factor |
| RA | Rheumatoid arthritis |
| RANK | Receptor activator of nuclear factor κ B |
| RANKL | Receptor activator of nuclear factor κ B Ligand |
| TMJ | Temporomandibular joint |

Introduction

RHEUMATOID ARTHRITIS

RA is one of the most common systemic inflammatory conditions with a worldwide prevalence of 0.5-1% (Kvien et al., 1997; Alamanos et al., 2006; Englund et al., 2010). It is more common in women with a female: male ratio of 3:1 (Symmons, 2002). The etiology of RA is still partially unknown but knowledge of the pathogenetic processes is increasing.

RA is mainly characterized by progressive articular cartilage degradation and bone tissue resorption as well as pain and loss of function in the musculoskeletal system. Damage to the articular cartilage and bone by the disease process accounts for a considerable part of the disability in RA (Scott et al., 2000). If untreated, it leads to severe disability as well as poor quality of life for most patients (Yelin et al., 1987; Hashimoto et al., 2001; Sokka, 2003).

The disease process starts years before the clinical onset and certain RA-specific autoantibodies, e.g. autoantibody against cyclic citrullinated peptide, are present in the blood years before the symptoms start (Aho et al., 1991; Aho et al., 1993; Rantapaa-Dahlqvist et al., 2003). Both experimental and clinical studies confirm that a subclinical phase precedes the clinically manifest arthritis (Kraan et al., 1998; Hayer et al., 2007). Hayer et al. (2007) showed that synovitis as well as osteoclast formation and activation is established in the articular synovium and underlying bone before the clinical symptoms appear.

Current diagnostic criteria for RA (Arnett et al., 1988) are mainly based on clinical findings typical for the chronic phase and have limitations regarding the possibilities to identify early phases of RA and to predict early disease development. Early diagnosis of RA has been considered important to identify individuals who will develop severe destructive disease, so that effective treatment can be initiated before irreversible damage occurs (van Aken et al., 2004).

The radiographic sign of erosion is generally considered to be an indicator of bone tissue resorption in RA. The presence of erosions in joints from hands and feet may serve as an estimate of RA activity (Forslind et al., 2009) and severity as it predicts for progression and poor prognosis when detected early in the disease (van der Heijde et al.,

1992). About 70 % of RA patients show erosions on plain radiographs of the small joints in hands and feet already at the time of diagnosis (van der Heijde, 1995; Goekoop-Ruiterman et al., 2005).

Temporomandibular joint rheumatoid arthritis

TMJ involvement is common in the late phase of RA with a prevalence varying from 5-86% depending on patient selection, diagnostic criteria and techniques used (Syrjanen, 1985; Tegelberg and Kopp, 1987; Lin et al., 2007). The most important clinical consequences of late phase TMJ RA are pain and functional disability, e.g. pain upon mouth opening and chewing or anterior open bite due to TMJ bone tissue resorption, which have a significant impact on the patients' daily activities and quality-of-life (Voog et al., 2003b).

The severity of TMJ RA is well correlated to the severity of general disease. Markers of inflammation such as C-reactive protein, erythrocyte sedimentation rate, rheumatoid factor and thrombocyte particle count have all been associated with the severity of TMJ RA (Goupille et al., 1993; Nordahl et al., 2001; Voog et al., 2003a; Lin et al., 2007). The destructive process does not necessarily correlate to local clinical signs and symptoms as radiographic signs of bone tissue resorption can be observed in a large proportion of asymptomatic patients (Goupille et al., 1990; Goupille et al., 1993; Celiker et al., 1995). RA involvement of the TMJ may therefore be undiagnosed and untreated early in the process. The question is to what extent radiographic signs of TMJ RA occur regardless of clinical signs and symptoms of the TMJ.

Radiographic examination of erosions in TMJ RA is difficult owing to the location of the joint and its relation to other cranial structures. However, when detected on conventional tomograms erosions in the late phase of TMJ RA have a frequency of about 50% (Gynther et al., 1996; Voog et al., 2003a) and progress in 19%-33% of the patients over one to four years (Nordahl et al., 2001; Voog et al., 2003a). It has been shown that the severity of radiographic changes in late TMJ RA is comparable to those of the metacarpophalangeal and metatarsophalangeal joints of the hands and feet (Akerman et al., 1991). The frequency and progression of TMJ bone tissue resorption in early RA is unknown as well as its predictive value for progression.

Crepitus is a common clinical sign of late phase TMJ RA with a reported frequency of 75% with a disease duration of more than five years (Tegelberg and Kopp, 1987). Crepitus is associated with cartilage and

bone tissue destruction in late phase TMJ RA (Kopp and Rockler, 1979; Akerman et al., 1988; Wiese et al., 2008). However, the relationship between crepitus and radiologic changes of the TMJ in RA patients is weak as 80% of joints with crepitus have shown evidence of TMJ bone tissue resorption whereas only 47% of joints with radiographic erosions had crepitus (Akerman et al., 1988). This indicates that articular cartilage and bone destruction may develop by different mechanisms in late phase TMJ RA. This relationship is not yet elucidated in recently diagnosed RA.

MECHANISMS OF BONE TISSUE RESORPTION

Bone is a dynamic tissue with continuous resorption and formation to adapt to its functional role. The bone turnover and remodeling is accomplished by coordinated intercellular signals between osteoblasts and osteoclasts which is dependent on the RANKL/RANK/OPG system under influence of mechanical loading and locally produced cytokines, growth factors and neuroregulators as well as hormones. For example, proinflammatory mediators as well as sex hormones influence the osteoclast differentiation and the osteoclastic bone resorptive activity (Takahashi et al., 1988; Lacey et al., 1998; Shevde et al., 2000; Michael et al., 2005; Galal et al., 2008).

RANKL modulates osteoclast differentiation and activity via the RANK receptor on the osteoclasts or pre-osteoclasts whereas OPG acts as a competitive antagonist to RANKL (Simonet et al., 1997). In patients with early RA the relation between RANKL and OPG predicts articular bone tissue resorption in hands and feet (Geusens et al., 2006). Activated T cells as well as synovial fibroblasts in active synovitis express more RANKL compared to those from patients with less active disease or from patients with osteoarthritis (Kong et al., 1999; Gravallese et al., 2000; Takayanagi et al., 2000; Kotake et al., 2001; Crotti et al., 2002; Haynes et al., 2003). This increase in RANKL expression is accompanied by a decrease in OPG expression in patients with active disease in comparison to patients with less active disease or healthy individuals (Haynes et al., 2003). The imbalance in RANKL, RANK and OPG expression in the synovial tissues probably favors local differentiation and activity of osteoclasts (Ainola et al., 2008). There is, however, only limited knowledge available about the influence by glutamate, serotonin, proinflammatory cytokines and hormones on the modulation of early TMJ RA bone tissue resorption.

Glutamate

Glutamate is an excitatory amino acid which is involved in peripheral mediation of inflammation and bone tissue remodeling in physiologic as well as in pathologic processes. Glutamate is also considered to be the mediator of load-related bone tissue remodeling (Skerry, 2002; Turner et al., 2002; Spencer and Genever, 2003; Mason, 2004). Glutamate is involved in bone resorption and formation through stimulation of N-methyl-D-aspartate (NMDA) receptors on osteoclasts and osteoblasts (Reynolds and Miller, 1988; Hollmann and Heinemann, 1994; Spencer et al., 2007). Glutamate activation of the NMDA receptor expressed on osteoclasts or osteoblasts regulate their expression of RANKL and RANK (Chenu et al., 1998; Peet et al., 1999; Genever and Skerry, 2001; Itzstein et al., 2001; Gu et al., 2002; Mentaverri et al., 2003; Ho et al., 2005; Spencer et al., 2007; Lin et al., 2008). A potential source of glutamate is the nerve fibers in the bone microenvironment (Chenu, 2002a; Chenu, 2002c). In RA, the sources of glutamate are activated thrombocytes, lymphocytes, macrophages, neutrophils, fibroblasts, local nerve fibers in the synovium as well as osteoblasts and activated osteoclasts in the bone tissue (Chenu et al., 1998; Lawand et al., 2000; Chenu, 2002b; Gravallese, 2002; Aliprandi et al., 2005; Tremolizzo et al., 2006; Spencer et al., 2007).

Local concentrations of glutamate are significantly elevated in acute and chronic inflammation in both animal models and humans (Omote et al., 1998; Lawand et al., 2000; McNearney et al., 2000). Elevated levels have also been reported from synovial tissues in RA (Lawand et al., 1997; McNearney et al., 2004; Tremolizzo et al., 2006). Glutamate is also elevated in plasma possibly partly as a result from activation of thrombocytes which release glutamate *ex vivo* when challenged with aggregating stimuli such as the rheumatoid factor (Trang et al., 1985; Aliprandi et al., 2005). Continuous thrombocyte stimulation with ensuing release of glutamate into the plasma may also be accomplished by proinflammatory cytokines such as TNF and IL-1 β since thrombocytes express cytokine receptors that reduce thrombocyte reuptake of glutamate (Loppnow et al., 1998; Aliprandi et al., 2005). There are thus reasons to believe that glutamate and the NMDA receptor are involved in the articular bone destruction occurring in RA.

MEDIATORS OF INFLAMMATION

Cytokines and serotonin are mediators involved in processes like inflammation and modulation of bone remodeling and turnover. TNF, IL-1 β , IL-6 and VEGF levels are increased both systemically in the blood and locally in the synovium and synovial fluid in RA where they have been associated with articular bone and cartilage destruction by enhancing RANKL expression and production of cartilage-destructive enzymes (Duff, 1994; Lacey et al., 1998; Ballara et al., 2001; Kwan Tat et al., 2004; Walsh and Gravallese, 2004).

TNF has the most prominent role in orchestrating the inflammatory process and immune response and is primarily produced by activated T-cells and macrophages in the inflamed synovium (MacNaul et al., 1990; Danning et al., 2000). TNF has the capacity to regulate osteoclast differentiation and function directly or indirectly, independent of its role in inflammation (Kobayashi et al., 2000; Komine et al., 2001).

TNF levels are increased in the synovial fluid and the synovial membrane of patients with late phase RA (Buchan et al., 1988; Saxne et al., 1988). There is also an increase in plasma and synovial fluid levels of cytokines in early-phase RA (Raza et al., 2005; Kokkonen et al., 2010). The TNF levels in plasma as well as in the TMJ synovial fluid are related to local TMJ bone tissue resorption (Nordahl et al., 1998; Nordahl et al., 2000; Voog et al., 2003a). Anti-TNF treatment of RA results in retardation of radiographic progression (Walsh and Gravallese, 2004).

IL-1 β is produced in high levels by activated macrophages and synovial fibroblasts but also by monocytes and osteoblasts during inflammation but not in healthy individuals where IL-1 β is undetectable in synovial fluid from the TMJ as well as in plasma (Eastgate et al., 1988; MacNaul et al., 1990; Dinarello, 1996; Alstergren et al., 1999; Danning et al., 2000). IL-1 β has strong bone-resorbing effects via receptors on preosteoclasts, osteoclasts and osteoblasts inducing osteoclast differentiation and activation (Tunyogi-Csapo et al., 2008; Jung et al., 2009). Similar to TNF, IL-1 β can indirectly regulate osteoclastogenesis by upregulating RANKL expression in osteoblasts (Hofbauer et al., 1999). IL-1 β can also promote the survival and function of mature osteoclasts (Jimi et al., 1999; Kobayashi et al., 2000; Suda et al., 2001).

Presence of IL-1 β in plasma and in the TMJ synovial fluid is related to TMJ destruction, expressed as anterior open bite, in patients with chronic inflammatory disorders (Alstergren et al., 1998). Neutralization of TNF and

IL-1 β by soluble IL-1 β and TNF receptors decreases the osteoclast formation and bone loss in experimentally induced arthritis (Walsh and Gravallese, 2004). In bone tissue resorption inhibition of IL-1 β by the IL-1 β receptor antagonist leads to decreased progression of radiographically assessed bone loss (van den Berg and Bresnihan, 1999).

IL-6 has a prominent role in inflammation-induced bone tissue resorption and is primarily produced by osteoblasts, which also express receptors for IL-6 (Bellido et al., 1996). IL-6 induces RANKL mRNA expression in murine osteoblastic cell lines (Nakashima et al., 2000). It is possible that IL-6 stimulates osteoblastic RANKL production and exerts an indirect effect on osteoclasts via the RANKL/RANK/OPG system (Kwan Tat et al., 2004). IL-6 synthesis in osteoblasts is also upregulated by decreased estrogen levels indicating a role in RA osteoporosis (Girasole et al., 1992; Poli et al., 1994).

VEGF plays a major role in RA angiogenesis by stimulating proliferation and migration of endothelial cells to form new blood vessels and also increases the permeability of the vessels (Brown et al., 1997; Jackson et al., 1997; Achen and Stacker, 1998; Paleolog, 2002). The source of VEGF includes fibroblasts, osteoblasts, macrophages, thrombocytes and lymphocytes (Brown et al., 1997). The expression is upregulated upon hypoxia mainly by macrophages and fibroblasts in the synovial membrane of RA patients and VEGF is therefore present in the synovial fluid of RA patients (Shweiki et al., 1992; Fava et al., 1994; Koch et al., 1994; Nagashima et al., 1995; Harada et al., 1998; Paleolog and Miotla, 1998).

Serum level of VEGF is elevated in patients with active RA compared to healthy individuals and it is related to disease activity (Nagashima et al., 1995; Harada et al., 1998; Ballara et al., 2001; Lee et al., 2001; Sone et al., 2001). Serum VEGF level in early RA was predictive of progression of bone tissue resorption after one year (Ballara et al., 2001). It was previously reported from a study of an arthritis model in mice that inhibition of angiogenesis also reduced the destruction of cartilage and bone (de Bandt et al., 2000; Miotla et al., 2000). Osteoclasts express VEGF receptors and stimulate osteoclastic bone tissue resorption in vitro and may therefore contribute to joint destruction by direct stimulation of osteoclasts and osteoclast precursors or indirectly by contributing to synovial hyperplasia (Nakagawa et al., 2000; Pap and Distler, 2005).

Peripheral serotonin is an endogenous mediator of inflammation which is involved in bone remodeling (Herbert and Schmidt, 1992; Pierce et al., 1995; Warden et al., 2005; Bliziotis et al., 2006; Modder et al., 2010). Serotonin has

a key role in bone formation via the 5-HT_{2B} receptor on osteocytes and osteoblasts (Bliziotis et al., 2001; Westbroek et al., 2001; Collet et al., 2008). Serotonin is involved in osteoblast induced inhibition of osteoclast differentiation by increasing OPG and decreasing RANKL secretion from osteoblasts (Gustafsson et al., 2006). Activation of serotonin receptors on osteoblasts, preosteoclasts and osteoclasts leads to bone tissue resorption and osteoclastogenesis (Battaglini et al., 2004; Yadav et al., 2008; Modder et al., 2010). Serotonin is mainly released from activated thrombocytes (Endresen, 1989; McNicol and Israels, 1999; Battaglini et al., 2004). It is present in bone marrow, synovial membrane, synovial fluid and plasma of patients with late phase RA (Alstergren and Kopp, 1997; Bliziotis et al., 2001; Kopp and Alstergren, 2002; Battaglini et al., 2004; Seide et al., 2004).

High plasma level of serotonin has been positively associated with progression of bone tissue resorption indicating an influence by serotonin on local bone tissue resorption. In addition high serum levels of serotonin were associated with regression while low levels were associated with progression in the late phase of TMJ RA (Voog et al., 2003a; Voog et al., 2004). The exact nature of the serotonin contribution to bone tissue resorption and the influence of sex is still unclear.

SYSTEMIC MARKERS OF DISEASE AND INFLAMMATORY ACTIVITY

Recent data from early undiagnosed arthritis show that solely presence of erosions is not predictive for development of RA. Patients who developed RA were more often positive for rheumatoid factor and had higher systemic disease activity than those who did not develop the disease (Thabet et al., 2009). This indicates that early detection of the disease needs to combine several predictors accurately.

C-reactive protein and erythrocyte sedimentation rate, together with rheumatoid factor and anti-citrullinated cyclic peptide, comprise the basis for estimation of systemic inflammatory activity in RA. These markers have all been associated with occurrence, progression and prediction of articular bone tissue resorption in RA (Amos et al., 1977; Larsen et al., 1977; van Leeuwen et al., 1993; Combe et al., 1995; Jansen et al., 2001; Syversen et al., 2008). C-reactive protein was found to be related to radiographic signs as well as progression of TMJ bone tissue resorption in RA (Celiker et al., 1995; Nordahl et al., 2001; Voog et al., 2004). C-reactive protein and erythrocyte sedimentation rate are up regulated as a result of elevated cytokine expression. However, C-

reactive protein is a better marker of inflammation and joint damage while erythrocyte sedimentation rate adds information reflecting disease severity (Bull et al., 1989; Wolfe, 1997; Jansen et al., 2001).

The rheumatoid factor is a well accepted predictor of disease activity in RA and poor prognosis (van der Heijde et al., 1992; van der Heijde, 1995; van Leeuwen et al., 1995; Uhlig et al., 2000; Vittecoq et al., 2003).

Rheumatoid factor has been found to be related to radiographic signs of TMJ bone tissue resorption in the late phase (Celiker et al., 1995). How these systemic markers reflect TMJ articular bone tissue resorption in early RA is not known.

SEX HORMONES

The influence by sex hormones on the disease activity of RA is an interesting topic since the frequency of RA is higher in fertile women and sex hormones have been suggested to be responsible for the sex-related differences in RA (Masi et al., 1999; Dao and LeResche, 2000; Kopp, 2001; Cutolo et al., 2004; Cutolo et al., 2006).

Both estradiol and testosterone are known to have immunomodulating effects but different roles with estradiol enhancing humoral immunity and androgens as natural immune suppressors (Masi et al., 1999; Cutolo and Wilder, 2000; Khosla et al., 2001; Riggs et al., 2002; Cutolo et al., 2004; Cutolo et al., 2006; Cutolo, 2009). Estrogen and testosterone have direct and indirect inhibitory effects on osteoclast formation and bone tissue resorption. Estrogens inhibit the release of TNF and IL-1 β by monocytes and T-cells and IL-6 by stromal cells and stimulate the secretion of OPG by osteoblasts. Testosterone enhances bone formation through its antiapoptotic effect on osteoblasts (Shevde et al., 2000; Riggs, 2002; Michael et al., 2005; Galal et al., 2008). Reduction of estrogens or androgens increases the rate of bone remodelling and also causes a focal imbalance between resorption and formation by prolonging the lifespan of osteoclasts and shortening the lifespan of osteoblasts (Vanderschueren et al., 2004).

Estrogen deficiency in post-menopausal women seems to be responsible for the major part of the differences in prevalence of RA between men and women (Straub, 2007). The increased bone tissue resorption in postmenopausal women is clearly related to estrogen deficiency (Riggs, 2002). In addition, estrogen and inflammation contribute additively to osteoporosis (Jochems et al., 2005). Testosterone levels have been

inversely related to clinical severity in RA (Gordon et al., 1988; Tengstrand et al., 2009). Low systemic and local levels of androgens, including testosterone, have been observed in both women and men with RA (Cutolo et al., 2005; Cutolo, 2009). Surprisingly, men with RA were found to have higher estradiol levels than healthy men and the levels were related to indices of inflammation (Tengstrand et al., 2003).

Aims

GENERAL AIM

The general aim of this thesis was to investigate the occurrence and development of the radiographic sign of TMJ bone tissue resorption in patients with early RA as well as to identify predictors for the development over the first two years of disease.

SPECIFIC AIMS

Cross-sectional part (Study I and II)

The aims of the cross-sectional part were to investigate:

- if the plasma level of glutamate is related to TMJ bone tissue resorption (Study I)
- if the relation between TMJ bone tissue resorption and glutamate is influenced by systemic inflammatory activity as well as levels of the sex hormones estradiol and testosterone (Study I)
- if TMJ bone tissue resorption is related to TMJ crepitus as a clinical sign of articular cartilage and bone damage (Study II)
- if TMJ bone tissue resorption in patients with or without crepitus is related to glutamate, serotonin, cytokines (TNF, IL-1 β , IL-6, VEGF), C-reactive protein, erythrocyte sedimentation rate, rheumatoid factor, estradiol or testosterone (Study II)
- if these relationships are influenced by sex (Study II)

Longitudinal part (Study III and IV)

The aims of the longitudinal part were to investigate

- whether TMJ bone tissue resorption, glutamate, crepitus of the TMJ, circulating inflammatory mediators (serotonin, TNF, IL-1 β , IL-6, VEGF) and markers (C-reactive protein, erythrocyte sedimentation rate) change over time (Study III)
- whether longitudinal changes in TMJ bone tissue resorption are related to changes in glutamate or TMJ crepitus (Study III)
- whether longitudinal changes in TMJ bone tissue resorption are related to change in inflammatory mediators (serotonin, IL- 1 β),

markers (C-reactive protein, erythrocyte sedimentation rate) or estradiol and testosterone (Study III)

- whether progression of TMJ bone tissue resorption can be predicted by glutamate, joint crepitus or serotonin (Study IV)
- whether progression of TMJ bone tissue resorption can be predicted by the circulating inflammatory mediators (IL-1 β , TNF, IL-6, VEGF) or inflammatory markers (C-reactive protein, erythrocyte sedimentation rate, rheumatoid factor) (Study IV)

Materials and methods

PATIENTS

Between May 2004 and May 2006, 160 out of approximately 380 patients recently diagnosed with RA were invited to participate in the present study. They were diagnosed at the Department of Rheumatology, Karolinska University Hospital in Huddinge, Sweden. A total of 47 patients (29 women and 18 men; 28 (60%) seropositive) accepted to participate and 41 (26 women and 15 men) of them fulfilled the two-year longitudinal follow-up. The complete numbers of patients included in the study are presented in Fig 1.

Six patients did not attend any follow-up examination, due to death (1), health reasons (2), unknown reasons (2) and panic in the CT (1). At the one-year follow-up one patient declined to participate because she did not experience any symptoms and another did not accept the CT examination due to pregnancy. At the two-year follow-up three patients dropped out for health or other personal reasons and one CT scan failed even after rescan. Out of 47 patients 38 fulfilled the longitudinal part, 35 of those went through all examinations at all three visits. The baseline characteristics of the 41 patients represented a typical RA patient population sample. A comparison with 379 recently diagnosed RA patients included in a Swedish multicenter survey, Better Anti-




Figure 1. Number of patients and examinations at first visit as well as at the one-year and two-year follow-up.

Table 1. Baseline characteristics of 41 patients in the current TMJ study and a RA patient sample of 379 patients from the BARFOT study.

| | TMJ study | n | BARFOT study | n |
|-------------------------------------------|------------|----|---------------|-----|
| Age (years) | 62 (53-67) | 41 | 55 (45-67) | 379 |
| Sex (female) | 71% | 41 | 65% | 379 |
| Disease duration (months) | 7 (4-10) | 41 | 6 (4-8) | 379 |
| Anti-cyclic citrullinated peptide (IU/mL) | 52% | 33 | 54% | 374 |
| C-reactive protein (mg/L) | 0 (0-18) | 41 | 19 (6-43) | 364 |
| Erythrocyte sedimentation rate (mm/h) | 22 (17-32) | 41 | 29 (14-50) | 378 |
| Rheumatoid factor (IU/mL) | 53% | 40 | 61% | 373 |
| DAS28 (0-10) | 6 (5-6.5) | 38 | 5.1 (4.2-5.8) | 371 |
| General pain (NRS 0-10) | 2 (0-5) | 41 | 4 (3-7) | 367 |
| TMJ pain (NRS 0-10) | 0 (0-0) | 41 | | |

Values are median (25th to 75th percentile) or percentage. NRS = numerical rating scale. The following values were considered positive: Anti-cyclic citrullinated peptide ≥ 25 U/mL, C-reactive protein ≥ 3 mg/L, erythrocyte sedimentation rate > 24 mm/h, rheumatoid factor ≥ 20 IU, and DAS28 > 3.2 . DAS28 = 28 joint disease activity score, n = number of patients .

Rheumatic Farmacotherapy (BARFOT), is shown in Table 1 (Forslind et al., 2004). BARFOT is an observation study with the intention to follow disease status and treatment in newly (maximum duration 12 months) diagnosed RA patients longitudinally over 15 years. Eighty-five percent (35/41) of the patients in the current project were also included in the BARFOT study.

The patients entered this study at a median (25th – 75th percentiles) of 23 days after diagnosis of RA. They were diagnosed with RA according to the revised American College of Rheumatology (ACR) criteria (Arnett et al., 1988), age above 18 and verbal consent. Exclusion criteria were current malignancies, TMJ surgery or trauma within two years, recent intra-articular glucocorticoid injection in the TMJ (within 1 month) as well as diseases other than RA as a cause of craniofacial pain. They were assigned to a brief questionnaire in order to evaluate if they fulfilled the following inclusion criteria; RA diagnosis according to the ACR criteria, duration less than one year and age more than eighteen years. The exclusion criteria were; current malignancies,

Table 2. Clinical and serological data at RA diagnosis and start of this study. 47 patients participated in the cross-sectional part (I and II) and 41 patients in the longitudinal part (III and IV).

| Study I and II (n = 47) | Diagnosis (%) | n | Study start (%) | n |
|-------------------------------------|----------------------|----------|------------------------|----------|
| General Pain | 100 | 45 | 83 | 47 |
| Pain of the Temporomandibular Joint | | | 13 | 47 |
| C-reactive protein | 100 | 45 | 43 | 47 |
| Erythrocyte sedimentation rate | 63 | 46 | 36 | 47 |
| Rheumatoid factor | 60 | 47 | 57 | 47 |
| Study III and IV (n = 41) | Diagnosis (%) | n | Study start (%) | n |
| General Pain | 100 | 41 | 71 | 41 |
| Pain of the Temporomandibular Joint | | | 12 | 41 |
| C-reactive protein | 100 | 39 | 37 | 41 |
| Erythrocyte sedimentation rate | 58 | 40 | 33 | 40 |
| Rheumatoid factor | 59 | 41 | 53 | 40 |

Values in percent abnormal. The following values were considered abnormal: C-reactive protein ≥ 3 mg/L, Erythrocyte sedimentation rate > 24 mm/h and Rheumatoid factor ≥ 20 IU. n = number of patients

TMJ surgery or trauma within two years, recent, less than one month, intra-articular glucocorticoid injection in the TMJ. Table 2 presents the variation in median (25th-75th percentile) values and frequency of clinical and serological data over 23 days after diagnosis.

TREATMENTS

The individual medication was based on the general clinical and serological status. The distribution of the principal treatments given is illustrated in Fig. 2. A majority of the patients started with a single DMARD, where the first choice was methotrexate but also salazopyrine or gold salt injections were given to a minority of the patients. At the three-month follow-up non-responders had changed treatment to either a combination of DMARDs, including methotrexate, sulfasalazine and hydroxychloroquine, or anti-TNF in combination with methotrexate.




Figure 2. Venn-diagrams showing the distribution (%) of major anti-rheumatic therapy as well as the combination with additional anti-inflammatory drugs at: first visit (47 patients), one-year follow-up (41 patients) and two-year follow-up (41 patients). Violet = anti-rheumatic therapy, green = low dose glucocorticoids, blue = non-steroidal anti-inflammatory drugs.

Glucocorticoids, either by oral or intra-articular administration were given as additional treatment to 20-27% of the patients. The need of supplemental NSAIDs, which were administered on a regular basis or in case of need, was reduced from 57% to 34% during the study period.

If a patient showed clinical symptoms of TMJ RA or extended radiographic signs of TMJ bone tissue resorption intra-articular administration with 0.5 mL methylprednisolone and lidocain (Depo-Medrol cum Lidocaine 40 mg/mL + 10 mg/mL) was performed. Three patients received bilateral local glucocorticoids at first visit and seven at the one-year follow-up (3 bilaterally and 4 unilaterally).

ETHICAL CONSIDERATIONS

All invited patients received verbal and written information about the project and time to think before deciding about participation. The study design, methods used and patient selection was approved by the ethical

committee at Karolinska Institutet, Stockholm, Sweden (03-204 & 452/03-204) and by the local radiation committee at Karolinska University Hospital in Huddinge, Sweden (19/03). All participating patients gave their informed consent according to the Declaration of Helsinki before inclusion.

RADIOGRAPHIC EXAMINATION

Radiographic examination of the TMJ region was performed with a craniofacial cone-beam CT (NewTom QR DVT mod 9000; QR s.r.l; Verona, Italy; Fig. 3A). The cone-beam technology obtained all the necessary data for the volumetric region desired by a complete rotation of the X-ray source and detector around a fix axis. The acquisition of the digital volume was carried out with the patient placed on an adjustable bench in a supine position with the TMJ located centrally in the region of interest (Fig. 3B). A reconstruction volume of 130x130x130 mm was performed and the mean scan time 75 seconds.

The acquired volume, the so-called raw data of the exam, were then able to be reconstructed in any plane desired giving planar images for analysis by the NewTom software (NewTom 3G). The reconstructions were processed by the same operator.




Figure 3. A) Cone-beam computerized tomograph, NewTom Model QR-DVT 9000, at the Department of Oral radiology, Karolinska Institutet, Huddinge. B) Patient adjustment, with the TMJ centrally in the region of interest, showing two laser indicators, a vertical line on the facial midline and a point over the TMJ lateral pole, used to precisely define the patient orientation.

Reconstruction procedure

The reconstruction procedure was performed in four steps: 1) Determination of the axial inclination 2) "Primary reconstructions", the reconstruction of axial views 3) Determination of the trans-axial plane 4) "Secondary reconstructions", the reconstruction of the trans-axial views used for the final analyses. The serial axial sections were obtained at 0.3mm (large field) and trans-axial sections at 0.5mm (35mm width) intervals.

Determination of the axial plane

The first step aimed to define a reproducible orientation of serial sections from the raw data. In order to control the orientation between three examinations reference planes were determined on a lateral cephalogram taken at every visit (Planmeca ProMax, Helsinki, Finland; Fig. 4a). The cephalogram was taken of the patient in a reliable natural head position keeping the mandible in exactly the same position as during the cone-beam CT scan with the help of an occlusion index (Lundstrom et al., 1995). As the digital volume was processed separately for each component of the TMJ, one reference plane was used for the temporal part and one for the condylar part. For the temporal component a reference plane parallel to the hard palate was used (Fig. 4A) and for the condylar component the reference plane was perpendicular to the mandibular ramus (Fig. 4B). The exact superior-inferior position of both reference planes was located across a constructed point 1cm inferior to the posterior nasal spine, at a line along the axis of the pterygo palatal fossa. In order to control the magnitude of the linear measurement on the lateral cephalogram a gauge was used.




Figure 4. A) A lateral cephalogram with the patient in the natural head position. B) Showing orientation and extension of the serial axial sections defined and transferred on tracings; temporal component (upper) and condylar component (lower). C) Showing orientation and extension of the serial axial sections from the raw data perspective; temporal component (upper) and condylar component (lower).

Primary reconstructions

The second step aimed to define a reproducible volumetric extension of serial axial sections, from the raw data, which originated from the predetermined reference planes. Therefore the reference planes and anatomic structures of interest were first traced at the lateral cephalogram on a transparent film, transferred and superimposed on the raw data image in the New Tom software. The superimposition was performed after synchronizing the magnification of the raw data view (Fig. 4C). The primary reconstruction could then be processed with the axial section of the volume at first visit through the reference plane and the last through the tangent of the most inferior point of the Turkish saddle. The total amount of section from first examination was saved as a reference for the follow-up exams and reconstructions.

Determination of the trans-axial plane

The fourth step aimed to define a reproducible orientation of serial trans-axial sections from the primary reconstruction. In order to do so a




Figure 5. A) Raw data view showing the extension of the serial axial sections and the level of axial section selected for the trans-axial reconstruction procedure. B) NewTom software tool showing the section level at maximum condylar width and long-axis inclination of the condylar head. C) Axial view showing the orientation of trans-axial sections used for analysis patients right temporal part. D) Axial view showing the orientation of trans-axial sections used for analysis of a patient's right condyle.

reference axial section incorporating the maximum condylar medio-lateral width was selected for each side of the temporal and condylar reconstructions and saved as a reference for the follow-up exams. The registered width (mm) and angulation of condylar axis was also saved for the repeated reconstructions. For every single patient the secondary reconstructions were processed four times; the right and left temporal part as well as on the right and left condylar part. The settings of trans-axial reconstructions were 35mm in width and 0,5mm thickness.

Secondary reconstructions

Allowing identification of the exact localization of the trans-axial sections of the temporal and condylar components the secondary reconstructions was defined in relation to the condylar long axis, the angulation and maximum width (Fig. 5B). That reference for the serial trans-axial sections was used and identical for the condylar part as well as the temporal part.

The trans-axial temporal sections were posteriorly-anteriorly oriented perpendicular to the long condylar axis and initiated two mm medially of the medial pole through all condyle to the lateral pole. Three specific sections were determined labeled with a marker for the final analysis, one in the middle of the long axis and the other two one third of the distances to the medial and lateral poles (Fig. 5A). The section numbers as well as distances were saved as a reference for repeated reconstructions. In order to reconstruct an identical section for all three cone-beam CT exams the reconstructions and measurements were performed from the medial pole to the lateral. The number of the three sections of interest from first examination was saved as a reference for the follow-up reconstructions.

The trans-axial condylar sections were medially-laterally oriented parallel to the condylar long axis and initiated two mm posteriorly, to a tangent through the most posterior condylar border and parallel to the long axis. Two specific sections were determined labeled with a marker for the final analysis, one through the long axis and the other at the middle of the distance between the long axis and the tangent of the anterior border of the condyle (Fig. 5B). The section numbers as well as distances were measured saved as a reference for repeated reconstructions. In order to reconstruct an identical section for all cone-beam CT exams the reconstructions and measurements were performed from the posterior to anterior. The number of the two sections of interest from first examination was saved as a reference for the follow-up reconstructions.

Analyses of radiographs

The cone-beam CT trans-axial sections were finally evaluated for presence or absence of erosions. Erosion was defined as a local area with decreased density of the cortical joint surface sometimes including the adjacent subcortical bone (Akerman, 1987). The planar images were viewed from the frontal aspect of the condyle and from the sagittal aspect of the temporal component. The two corresponding sections of the condyle, one central (the section with the largest condylar width) and one anterior were analyzed in the lateral, central and medial parts. The extension of erosive changes was expressed as a score ranging from 0 to 24 for each patient (Fig. 6). The sum of the score of the right and left sides




Figure 6. Red markers illustrate the orientation and precise position of the transaxial sections from an axial view of the right and left temporomandibular joint. The readings of the temporal component were performed on three lateral sections perpendicular to the condylar axis, a lateral, central and medial, and of the condylar component two anterior sections parallel to the condylar axis, a central and anterior, were used.

for each individual was used in the analysis. The development over time was expressed as the difference between the final score at two-year follow-up and the first visit.

The reading of the erosions was made by one skilled radiologist. When erosions were assessed, their presence was verified in the frontal, sagittal and axial views. Intra observer consistency between repeated readings of erosions was tested by a randomized selection of 20 joints from 10 individuals. The scoring system was tested in a blinded fashion by the same radiologist two to three years apart. On a patient level, the repeated readings showed a difference of one score unit between the two occasions in three patients, whereas the readings of the other seven patients showed no differences. The relative reproducibility of detection of erosive changes varied between 70 and 100% for the different areas where the medial part of the anterior frontal section of the condyle showed the lowest degree of reproducibility and regions on the temporal part the highest.

The longitudinal development of two patients seen in the frontal view of the central sections through the condylar axis is illustrated in Fig. 7. They demonstrated advanced erosions at first visit or at the one-year follow-up.




Figure 7. Frontal views of cone-beam computerized tomographs, showing the two-year development of a temporomandibular joint in two patients with TMJ erosions. Patient A showed TMJ erosion at first visit and received intra articular injection of glucocorticoids. At the one-year follow-up the erosion was absent and the new bone stayed in remission at the two-year follow-up. Patient B developed TMJ erosion during the first year after diagnosis which was treated with intra articular glucocorticoid injection at the one-year follow-up. New bone formation was observed at the two-year follow-up.

CLINICAL EXAMINATION

General clinical variables

A disease-activity score based on 28 joint counts (DAS28, 0-10) was assessed by a rheumatologist when the diagnosis was determined at the Department of Rheumatology, Karolinska University Hospital, Huddinge, Sweden.

Number of involved joints was a summation variable used as a reference for the general clinical status. Patient reported pain, tenderness, stiffness or swollenness from nine joint regions (neck, upper and lower back, shoulders, elbows, hands, hips, knees and feet), giving a maximum score of 9.

Clinical signs of TMJ bone tissue resorption

TMJ crepitus was assessed bilaterally as a clinical sign of cartilage and bone tissue resorption. The degree of crepitus was recorded during mandibular rotational and translational movements using a score (0-4), where 1 = palpable and 2 = audible.

Blood sampling and laboratory procedure

Venous blood samples were collected the same day as the clinical and radiographic examinations. In order to assess the inflammatory activity the inflammatory markers erythrocyte sedimentation rate and C-reactive protein as well as the disease marker rheumatoid factor were analyzed in serum. Blood samples were also taken to determine the serum levels of estradiol and testosterone. All these analyses except for erythrocyte sedimentation rate were performed with standard procedures at the Department of Laboratory Medicine (Clinical Chemistry and Immunology), Karolinska University Hospital, Huddinge, Sweden. Determination of the erythrocyte sedimentation rate was performed according to the Westergren procedure. Venous blood sampled in EDTA tubes was immediately cooled and centrifuged (1500 *g* for 10 min at +4°C) for analysis of glutamate, TNF, IL-1 β and IL-6 in plasma. Other venous blood samples were stored without additives in room temperature for 1 hour to coagulate and thereafter centrifuged (1500 *g* for 30 min at +4°C) for analyses of serotonin serum levels. VEGF was analysed in lysed whole blood. After preparation the supernatants were pipetted into polypropylene tubes except for serum samples that were stored in polystyrene tubes in -80°C until analysis. The concentrations of

all investigated mediators were determined by commercially available assays at the Department of Dental Medicine Laboratory. Glutamate in plasma was analysed by the Amplex® Red Glutamic acid/glutamate oxidase assay (Amplex, Invitrogen, Eugene, USA). This glutamate assay had a detection limit of 10 µmol/L according to the manufacturer. TNF, IL-1β and IL-6 in plasma were analysed with the high-sensitivity human cytokine Lincoplex kit for simultaneous multianalyte detection with Luminex technology and instrumentation (Lincoplex kit, Linco Research, St Charles, USA). The assay minimum detection concentration is 0.05 (TNF), 0.06 (IL-1β) and 0.10 (IL-6) pg/mL, respectively. The intraassay coefficients of variation are 3.5% (TNF), 3.1% (IL-1β) and 3.5% (IL-6) and the interassay coefficient are 3.8% (TNF), 2.2% (IL-1β) and 4.5% (IL-6). VEGF was analysed by a commercially available enzyme-linked immunoassay kit (Quantikine Human Immunoassays, R&D Inc, Minneapolis, USA) in lysed whole blood. Serotonin was analyzed by a commercially available competitive enzyme immunoassay kit (EIA-kit. No 0642, Immunotech International, Marseille, France). The kit has a detection limit and sensitivity of 0.5 nmol/L. The intraassay coefficient of variation is less than 9.4% and the interassay coefficient of variation less than 9.9%, according to the manufacturer.

STATISTICAL ANALYSES

Median, min, max and the 25th and 75th percentiles as well as mean and standard deviation were used for descriptive statistics. Both parametric and nonparametric methods were used when applicable. Normality, constant variance and power were tested for each analysis. A probability level of less than 0.05 was considered significant.

Cross-sectional part (study I and II)

Significance of univariate correlations was calculated with Spearman rank correlation coefficient (r_s). Correlations between the variables of interest were tested in all patients and the significance of the correlations between TMJ erosion score and glutamate was tested in prespecified subgroups according to C-reactive protein, erythrocyte sedimentation rate, crepitus, estradiol and testosterone as well as sex. The significance of the differences between groups was tested with the Mann-Whitney U-test. In order to select the strongest independent variables explaining TMJ erosion score a stepwise linear multiple regression analysis was performed. Independent variables (Sex, serotonin, TNF, IL-1β, IL-6, VEGF,

estradiol, testosterone, C-reactive protein, erythrocyte sedimentation rate and rheumatoid factor) were included in the multivariate regression to test for the dependent (parametric) variable TMJ erosion score in order to investigate their relative importance to explain erosions. Interaction analysis was performed by including an interaction term in addition to the main effects in a multiple linear regression model.

Longitudinal part (study III and IV)

The significance of the differences between visits was tested by the Friedman ANOVA on ranks. Change of independent variables (joint crepitus, glutamate, serotonin, TNF, IL-1 β , IL-6, VEGF, C-reactive protein, erythrocyte sedimentation rate, rheumatoid factor, estradiol and testosterone) between first visit and the two-year follow-up were tested for multivariate regression to the dependent variable change in TMJ erosion score in order to investigate their relative influence. The significance of univariate correlations in change between the variables was tested in all patients as well as in prespecified subgroups regarding sex, TMJ crepitus and glutamate by the Spearman rank correlation coefficient (r_s). In order to select the strongest independent predictors to change in TMJ erosion score a stepwise linear multiple regression analysis was performed. Independent variables (initial TMJ erosion score, sex, joint crepitus, glutamate, serotonin, TNF, IL-6, IL-1 β , C-reactive protein, erythrocyte sedimentation rate and rheumatoid factor) at the start of the study were tested for multivariate regression to the dependent (parametric) variable change in TMJ erosion score in order to investigate their relative importance to predict the changes of erosions. The significance of the differences between groups was calculated with Fisher's exact test for variables on a nominal scale (radiographic progression, regression) and Mann-Whitney U-test for variables on an ordinal scale. Interaction analysis was performed by including an interaction term in addition to the main effects in a multiple linear regression model.

Results and discussion

CROSS-SECTIONAL PART (STUDY I AND II)

TMJ bone tissue resorption is common in early RA

A total of 72% of the patients had TMJ bone tissue resorption with a median TMJ erosion score (25th - 75th percentile) of 2 (0 - 4). There was a great inter-individual variation with a maximum score of 9, which was found in one patient.

A large proportion of the patients, all with a disease duration less than one year, showed signs of TMJ bone tissue resorption. The extension of the resorption was small in general with a median score of 2 on a 0-24 scale. This finding, nevertheless, indicates that the TMJ is commonly and very early involved in RA, which has not been known previously. In two earlier studies, 50-52% of RA patients with a disease duration of 9-14 years had radiographic signs of TMJ bone tissue resorption (Gynther et al., 1996; Voog et al., 2003a). The considerably higher frequency reported in our study can be explained by the use of a different and more sensitive technique with computer tomography.

TMJ bone tissue resorption is related to glutamate and crepitus and inflammatory activity

TMJ erosion score was positively correlated to glutamate ($r_s = 0.30$, $n = 47$, $p = 0.043$) and TMJ crepitus ($r_s = 0.38$, $n = 47$, $p = 0.008$). Multiple stepwise regression showed that TMJ crepitus ($p < 0.001$) together with glutamate ($p = 0.005$) and erythrocyte sedimentation rate ($p = 0.010$) were positively related to the erosion score and explained 41% of its variation ($p < 0.001$). Sex, serotonin, TNF, IL-1 β , IL-6, VEGF, C-reactive protein, erythrocyte sedimentation rate, rheumatoid factor, estradiol, progesterone and testosterone were also included into the stepwise regression but did not significantly add to the ability of the model to predict erosion score. In the patients without TMJ crepitus, bone erosion score was positively correlated to glutamate ($r_s = 0.45$, $n = 35$, $p = 0.007$). The corresponding correlation in the patients with crepitus was negative but did not reach statistical significance ($r_s = -0.35$, $n = 12$, $p = 0.264$). Interaction analysis with multiple linear regression showed that there was a different relation ($p = 0.050$) between bone erosion score and glutamate level in the patients without crepitus compared to those with

crepitus. In a subgroup of patients with estradiol levels ≥ 65 pmol/L there was a positive correlation between erosion score and C-reactive protein ($r_s = 0.68$, $n = 16$, $p = 0.004$). The erosion score was higher in patients with C-reactive protein levels ≥ 3 mg/L and estradiol ≥ 65 pmol/L than those with lower C-reactive protein and estradiol levels ($p = 0.025$, $n = 16$).

These results agree with the hypothesis that glutamate is associated with bone tissue resorption of the TMJ RA as assessed by radiographic erosions. It seems that glutamate is released in the bone microenvironment by osteoblasts, activated osteoclasts and peripheral nerves as part of normal bone remodeling but also by inflammatory cells in the inflamed synovium (Lawand et al., 2000; Genever and Skerry, 2001; Chenu, 2002b; Gravallese, 2002; Staud et al., 2003; Aliprandi et al., 2005; Tremolizzo et al., 2006; Spencer et al., 2007). In turn, activation of NMDA receptors stimulates osteoblast precursor proliferation, NF κ B-mediated osteoclast differentiation and inhibition of osteoclast apoptosis and the resulting net signaling effect modulates changes in bone mass (Spencer et al., 2006). The mature resorbing osteoclast is a target for glutamate by the NMDA receptor, which when activated results in bone tissue resorption in the healthy organism (Saika et al., 2001; Chenu, 2002a; Spencer et al., 2007). Fibroblast-like synoviocytes from patients with active RA express glutamate receptor mRNA for the glutamate NMDA, kainate and AMPA receptors, where activated kainate/AMPA receptors increase the release of proinflammatory cytokine IL-6 and subsequently matrix metalloproteinase 2, an enzyme highly involved in matrix degradation (Flood et al., 2007). The findings indicate that glutamate modulates bone tissue resorption, which is an intriguing pathophysiologic mechanism in RA as previously suggested by Flood et al. (2007). Flood and coworkers also found that glutamate receptor activation may contribute to the synovial hyperplasia and thus pannus development by stimulation of synoviocyte proliferation. Whether glutamate regulates release of cytokines from synoviocytes or other synovial cells remains to be investigated.

Glutamate had a stronger correlation with TMJ bone tissue resorption when crepitus was absent. The correlation could not be found when crepitus was present. One possible explanation might be that the crepitus in this study was mainly palpable (21%) and infrequently audible (2%), which indicates a lower degree of destruction in agreement with the short disease duration. The frequency of crepitus was also lower (26%)

than in a previous study of TMJ RA where 75% of both men and women were reported to have crepitus (Tegelberg and Kopp, 1987). However, the majority of these patients had a disease duration of more than 5 years. The relation between crepitus and TMJ bone tissue resorption was significant. Another study reported a weak but statistically significant relationship between crepitus and radiologic changes of the TMJ in RA patients, where 80% of joints with crepitus showed evidence of bone tissue resorption while only 47% of the joints with bone tissue resorption had crepitus (Akerman et al., 1988). This latter finding is in agreement with ours and indicates that articular cartilage and bone tissue destruction develop with different mechanisms in TMJ RA.

The regression model also included erythrocyte sedimentation rate as a predictor of bone tissue resorption when the other variables were accounted for. Neither C-reactive protein, erythrocyte sedimentation rate nor rheumatoid factor was significantly correlated to the bone erosion score in the non-parametric univariate analysis. The multivariate relationship between bone tissue resorption and erythrocyte sedimentation rate should therefore be interpreted with caution. However, such a relation can be expected for erosions of systemic inflammatory nature and a strong association between serial measurements of erythrocyte sedimentation rate and radiographic joint destruction has been found (Graudal, 2004). TMJ bone tissue resorption was thus not directly related to systemic inflammatory activity in the univariate analysis but there was a correlation found between TMJ bone tissue resorption and C-reactive protein which depended on the level of estradiol. Estradiol has been found to increase the expression of osteoprotegerin via the estrogen receptor α (Yoneda et al., 2004). Osteoprotegerin reduces osteoclastic bone tissue resorption by binding to RANKL. Estradiol can also decrease proinflammatory cytokine release through modulation of CD16 expression in monocytes and monocyte-derived macrophages (Kotake et al., 2001). These results are consistent with the finding of no erosions in the RA patients with estradiol levels \geq 65 pmol/L combined with low systemic inflammatory activity. However, the erosion score correlated to C-reactive protein in the patients with estradiol level \geq 65 pmol/L, i.e. a combination of high levels of C-reactive protein and estradiol resulted in a high score of erosions. In previous studies of progression of radiographic changes in RA, time-averaged C-reactive protein levels were correlated to increase in Larsen score and regarding the TMJ, progression during a period of 25-46 months was

related to C-reactive protein level (Plant et al., 2000; Voog et al., 2004). The influence of estradiol was not investigated in those studies. The short disease duration as well as the cross-sectional character of the present study may also influence the relation between systemic inflammatory activity and erosions.

Glutamate modulates TMJ bone tissue resorption at low systemic inflammation and low levels of sex hormones

Glutamate levels in plasma correlated to TMJ erosion score in the subgroup of patients with C-reactive protein levels of < 3 mg/L ($r_s = 0.46$, $n = 27$, $p = 0.017$) or to testosterone levels ≤ 1.2 nmol/L ($r_s = 0.56$, $n = 21$, $p = 0.008$). There was stronger positive correlations between glutamate and erosions in the patients with C-reactive protein < 3 mg/L and testosterone levels ≤ 1.2 pmol/L ($r_s = 0.84$, $n = 14$, $p < 0.001$) as well as in the patients with C-reactive protein < 3 mg/L and estradiol levels < 50 pmol/L ($r_s = 0.63$, $n = 13$, $p = 0.022$). The results suggest an influence on the relationship between plasma glutamate and TMJ bone tissue resorption by testosterone since this relation was considerably stronger in the patients with testosterone levels ≤ 1.2 nmol/L. In addition, plasma glutamate was associated with TMJ bone tissue resorption particularly in patients with low systemic inflammatory activity, a factor that in combination with estradiol and testosterone seems to influence the glutamate modulation of TMJ bone tissue resorption.

Glutamate may thus be a modulator of connective tissue cell function regarding bone tissue resorption in joints with RA especially in individuals with low systemic inflammatory activity and low levels of sex hormones. Circulating glutamate may activate locally expressed receptors on e.g. osteoclasts or fibroblast like synoviocytes and modulate bone tissue resorption in patients with low systemic inflammation in combination with low estradiol or testosterone and thereby represent a bone resorbing mechanism independent on systemic inflammation. Estrogen deficiency plays a crucial role in acceleration of bone tissue resorption in autoimmune arthritis associated with RANKL-mediated osteoclastogenesis in a murine model for rheumatoid arthritis (Yoneda et al., 2004). This finding may explain the bone tissue resorption in the patients with estradiol levels < 50 pmol/L in combination with high systemic inflammatory activity. RANKL mRNA and protein have been identified in cultured synovial fibroblasts from patients with RA (Khalkhali-Ellis et al., 2000). Glutamate may modulate the bone tissue

resorption via the RANKL/RANK/OPG system under influence of estradiol and cytokines.

The influence by testosterone in bone tissue resorption is more unclear in RA (Cutolo, 2009). However, low testosterone levels may be expected during inflammation since systemic proinflammatory cytokines suppress the hypothalamic-pituitary-gonadal axis (Jones and Bhalla, 1993). In testosterone deficiency the inhibiting influence on osteoclast formation and activity is reduced (Michael et al., 2005). Androgens including testosterone are down-regulated in active RA synovial tissues and systemic levels are reduced in both women and men in RA and inversely related to clinical severity (Gordon et al., 1988; Cutolo et al., 2005; Cutolo, 2009; Tengstrand et al., 2009). As androgens exert anti-inflammatory activities in RA synovium one may hypothesize that low androgen levels may facilitate local inflammation (Cutolo et al., 2007).

Serotonin mediates TMJ bone tissue resorption in men

The TMJ bone erosion score was positively correlated to serotonin level in serum ($r_s = 0.54$, $n = 16$, $p = 0.030$) in men. Interaction analysis showed a significant ($p = 0.014$) difference between the sexes.

The relationship between bone tissue resorption and serum level of serotonin seems to be positive and specific for men since there was a significant sex difference regarding this relationship. Serotonin may modulate osteoblast bone formation via the 5-HT_{2B} receptor and the osteoclast differentiation through its transporter 5-HTT (Trang et al., 1985; Westbroek et al., 2001; Battaglino et al., 2004; Collet et al., 2008). In this study the observed bone tissue resorption cannot be considered as normal remodeling due to the combination with active RA. However, serotonin may play an important role for articular bone tissue remodeling in both healthy individuals and in patients with RA since osteocytes and osteoblasts express receptors for serotonin and 5-HTT (Bliziotis et al., 2006). A longitudinal study of mainly women with late phase RA showed that low serum levels of serotonin were associated with progression of erosions, while high levels were associated with regression during a 25 – 46 month interval (Voog et al., 2004). Whether serotonin in serum is associated with continuing bone tissue resorption in the early stage of RA needs further studies. The exact nature of the serotonin contribution to bone tissue resorption and the influence of sex is, however, still unclear.

LONGITUDINAL PART – DEVELOPMENT (STUDY III)

Change in TMJ bone tissue resorption is inconsistent

The TMJ erosion score for all patients was reduced from a median (25th - 75th percentile) of 2 (0 - 4) to 0(0 - 2). The reduction was not significant through the two first years after diagnosis ($p = 0.488$). The change in erosion score varied inconsistently over time as 29 % (11/37) showed progression of erosion score, 43 % (16/37) showed regression and 28% (10/37) showed the same erosion score at the two-year follow-up as at study start.

The results show that a substantial portion of patients can be expected to show a regression or no change in TMJ bone tissue resorption over the first two years after diagnosis of RA. The progression of TMJ bone tissue resorption was less marked than the regression over the two years after RA diagnosis. The TMJ progression rate was similar to the 19-33% found in patients with late phase and symptomatic TMJ RA in well-controlled disease and general duration of about 9-14 years (Nordahl et al., 2001; Voog et al., 2004). Regression in TMJ erosions was also seen in 31% of the patients in the study by Voog et al. (2004). The progression was less marked than that reported from hand and foot joints on plain radiographs, where erosion score progressed in 63.6% of the patients over three years despite early treatment with conventional DMARDs (Machold et al., 2006). The lower progression rate in the TMJ in the current study may be explained by the early general treatment given. The group of patients who showed progression of TMJ bone tissue resorption may belong to a subgroup who does not respond to the anti-inflammatory treatment or that the bone tissue resorption may include a non-inflammatory osteoporosis-like resorption (Machold et al., 2007; Clarke and Khosla, 2010).

TMJ bone tissue resorption is modulated by inflammatory and non-inflammatory mechanisms

The erythrocyte sedimentation rate decreased ($p = < 0.001$, $n = 33$) but not C-reactive protein ($p = 0.540$, $n = 36$) during the study period. Multiple stepwise regression showed that change of the independent variables IL-1 β ($p = 0.024$) and testosterone ($p = 0.012$) were positively related to the change in erosion score ($p = 0.010$, $n = 21$) and explained 34% of that change over two years. IL-1 β ($p = 0.027$, $n = 36$) as well as testosterone ($p = 0.048$, $n = 32$) increased during the two years period.

Progression of TMJ bone tissue resorption can be expected to occur especially in patients who show an increase of testosterone and IL-1 β in the blood during the first two-year of disease. IL-1 β is involved in the mechanisms behind destruction of bone and cartilage in RA. IL-1 β acts through two different pathways; by differentiation and activation of osteoclasts by upregulating RANKL expression and by triggering production of proteolytic enzymes from synoviocytes and inflammatory cells in the inflamed synovium (Rengel et al., 2007; Schett, 2007). Increase of IL-1 β appeared in this study as the most important inflammatory factor explaining progression of bone tissue resorption. The weak relation between the systemic markers of inflammation and progression of TMJ bone tissue resorption is in contrast to earlier studies where radiographic progression in hands and feet as well as the TMJ was related to C-reactive protein levels (Plant et al., 2000; Voog et al., 2004). One explanation may be diverging treatment strategies or the time between start of treatment and first visit in the current study.

The influence of testosterone on bone tissue resorption is unclear in RA (Cutolo, 2009). Testosterone has immunosuppressive properties (Masi et al., 1999). In addition, the testosterone level increases when the disease activity decreases in men (Gordon et al., 1988; Tengstrand et al., 2009). The testosterone levels increased during the two-year period in our study and the systemic inflammation decreased. Our results showed that the increase of testosterone is positively associated with progression of bone tissue resorption during the first two years of disease, independently of IL-1 β . The positive relationship between the increase of testosterone and TMJ bone tissue resorption is in contrast to earlier studies where testosterone has been related to inhibition of osteoclast formation and activity (Riggs, 2002; Michael et al., 2005). In addition, we have also shown that testosterone was associated with TMJ bone tissue resorption in RA patients shortly after diagnosis (Study I). The influence by testosterone on articular bone tissue resorption needs further evaluation, especially regarding character of bone tissue resorption.

Progression of TMJ cartilage destruction is influenced by glutamate and systemic inflammation

Glutamate increased from a median(25th-75th percentiles) of 3.7(1.0-7.3) mmol/L to 9.1(5.8-17.5) mmol/L during the two-year period ($p < 0.001$). In a subgroup with glutamate increase more than our median value ≥ 3.4 mmol/L between first visit and two-year follow-up progression of

crepitus was positively correlated to increase in erythrocyte sedimentation rate ($r_s = 0.59$, $n = 17$, $p = 0.012$) while patients with less than median increase of glutamate (< 3.4 mmol/L) showed a negative correlation ($r_s = -0.77$, $n = 16$, $p < 0.001$). Interaction analysis with multiple regression showed that the difference between these groups was significant ($p < 0.001$). TMJ crepitus increased ($p = 0.020$) but that change was not correlated to TMJ erosion score. The serum level of serotonin decreased ($p = 0.003$, $n = 36$) over the two-year period. The reduction in serotonin was larger in the patients who had TMJ crepitus after two years (median = -406 , 25th/75th percentile = $-202/-505$) than those without (median = -137 , 25th/75th percentile = $-391/5.3$; $p = 0.045$). The thrombocyte particle count decreased during the two years after RA diagnosis ($p < 0.001$, $n = 34$).

The subgroup with increase of plasma glutamate of more than 3.4 mmol/L showed a positive correlation between change in TMJ crepitus, a clinical sign of cartilage or bone damage and change in the inflammatory marker erythrocyte sedimentation rate. This finding is in agreement with earlier studies by Flood et al. (2007) who showed that elevated glutamate in synovial fluid may influence destruction of cartilage by stimulating synovial fibroblasts to express proMMP-2 in the synovial tissues (Flood et al., 2007). Development of crepitus from cartilage or bone degradation may thus be due to release of glutamate in the inflamed synovium over the first two years after diagnosis. However, there was a positive relationship between glutamate and TMJ erosion score at the first visit in the absence of crepitus, which may be due to not yet sufficiently developed crepitus at this early stage. Glutamate may partly be involved in the cartilage degradation and partly in the bone tissue resorption through different mechanisms where the cartilage degradation is a slower process and dependent on the local inflammation (Study II). TMJ crepitus increased during the two-year period, indicating escalated articular cartilage damage. However, this change was not related to the change in the TMJ erosion score, which further suggests that articular cartilage and bone tissue degradation are mediated by at least partly independent mechanisms (Schett, 2007). Crepitus reflects the inflammatory event which leads to expression of matrix-degrading enzymes and their destruction of the surface cartilage (Pap et al., 2001).

Serotonin was reduced and the reduction was more pronounced in patient with crepitus at the two-year follow-up indicating that the reduction of serotonin was more pronounced in patients with

destructive changes of the articular surface of the TMJ. The osteoblast has receptors for serotonin (Bliziotis et al., 2001; Westbroek et al., 2001; Collet et al., 2008) and the biological effect of these receptors may be reduced by reduction of serotonin and thereby modulate bone turnover.

LONGITUDINAL PART - PREDICTION (STUDY IV)

Regression of TMJ bone tissue resorption is predicted by the early extent of TMJ bone tissue resorption

The change of TMJ erosion score was negatively correlated with erosion score at the first visit ($r_s = -0.64$, $n = 38$, $p < 0.001$) i.e. regression of bone tissue resorption was associated with high initial erosion score. Multiple stepwise regression showed that initial erosion score was the only significant and a negative predictor explaining 48% of the variation of the change in erosion score after two years ($p < 0.001$, $n = 35$). However, univariate analysis showed that patients with joint crepitus at the first visit more frequently had a decrease in erosion score ($7/10 = 70\%$) than the patients without ($9/28 = 32\%$; $p = 0.044$).

The association between presence of early erosions and regression of erosions may be due to the phenomenon called “regression towards the mean”, which means that spontaneous remission of abnormal values, unrelated to treatment, is likely to occur if many patients show abnormal values before treatment. However, in this case the decrease in TMJ bone tissue resorption is likely to be mainly due to the treatment since the RA disease normally progresses regarding articular tissue destruction. Another possibility is that there was an observer change in the reading of the radiographic data between visits, i.e. that a more cautious registration was made of erosions at follow-up. This is a possibility since the time period between registrations was about two years. However, the change in erosion score was associated with changes in e.g. IL-1 β , TNF, VEGF and testosterone, which rather indicate that the change in regression was related to biologic processes (Study III).

Patients with presence of crepitus at the first visit showed a decrease in TMJ erosion score more frequently than patients without crepitus suggesting that patients where the surface cartilage is damaged have a better response to anti-inflammatory treatment. In turn, this indicates that crepitus in RA is a sign of inflammatory cartilage and bone tissue destruction. Crepitus is a clinical sign of cartilage destruction that may

not be present when RA starts but develops gradually and later becomes a common feature (Tegelberg and Kopp, 1987; Akerman et al., 1988; Wiese et al., 2008). However, the mechanisms behind articular cartilage and bone tissue resorption may differ in the early disease process (Appel et al., 2006; Schett, 2007; Appel et al., 2009; Study II; Study III). In RA, cartilage destruction is probably a direct consequence of the synovial inflammation with cytokines and other inflammatory factors that elicit production and release of proteolytic enzymes (Pap et al., 2001). Bone tissue resorption is related to an unbalance in the RANKL/RANK/OPG system most likely caused by synovial inflammation but also by non-inflammatory osteoporosis (Clarke and Khosla, 2010).

Regression of TMJ bone tissue resorption is predicted by high plasma level of glutamate

Patients with plasma glutamate at median level or above (≥ 3.7 mmol/L) at first visit showed a more frequent decrease (11/18 = 61%) in erosion score than patients with a level below the median (< 3.7 mmol/L; 5/19 = 26%; $p = 0.035$, $n = 37$; Table 3) over two years. The group with higher plasma level of glutamate showed a median decrease in TMJ erosion score of -1 compared to no change in the group with lower level.

High plasma level of glutamate (≥ 3.7 mmol/L) early after diagnosis seems to predict decrease of TMJ bone tissue resorption during the first two years of RA. It seems that glutamate modulates synovial inflammation but also non-inflammatory bone tissue resorption (Chenu et al., 1998; Flood et al., 2007). Glutamate was related to bone tissue resorption in patients with low systemic inflammatory activity, which indicates a role of glutamate in non-inflammatory bone tissue resorption (Study I). In addition, glutamate was associated with bone tissue resorption selectively in patients without joint crepitus (Study II). It is therefore possible that glutamate has a dual role in bone turnover by regulating both bone tissue resorption and bone formation giving the possibility of glutamate to promote bone repair in RA (Chenu et al., 1998). The glutamate NMDA receptor on preosteoclasts and osteoclasts stimulates differentiation and activation, i.e. bone tissue resorption, while it also prevents bone tissue resorption by activating osteoblasts (Chenu et al., 1998).

Regression of TMJ bone tissue resorption is predicted by serotonin in men

In the men, the change in erosion score was negatively correlated to serum serotonin level at the first visit ($r_s = -0.57$, $n = 14$, $p = 0.034$), which means that the extent of TMJ bone tissue resorption decreased in men with high initial levels of serotonin. Interaction analysis with multiple linear regression showed a significant ($p = 0.034$) interaction between sex and serotonin. This implies a significant difference in relation between TMJ bone tissue resorption and serotonin in men compared to women.

In a cross-sectional study high serum serotonin was positively associated with TMJ bone tissue resorption in males with early RA (Study II). In the present study, early and high serotonin levels in serum predicted regression of TMJ bone tissue resorption after two years in the male RA patients. During the first two years in RA disease serotonin decreased significantly in men (Study III). Our results are in agreement with a previous study where the serum level of serotonin also predicted regression of TMJ bone tissue resorption in a chronic phase of RA (Voog et al., 2004). High serotonin levels in TMJ synovial fluid have been found to predict a positive treatment response on TMJ pain by intra articular anti-inflammatory treatment in patients with chronic RA, suggesting that serotonin is involved in the inflammatory process (Fredriksson et al., 2005). These results indicate that serotonin contribute to inflammatory bone tissue resorption.

Inflammatory markers do not predict TMJ bone tissue resorption

None of the inflammatory markers C-reactive protein, erythrocyte sedimentation rate or rheumatoid factor did significantly add to the ability of the multiple regression model to predict TMJ erosion score when included in the stepwise regression.

This result differs from previous studies in local as well as general involvement of RA (Celiker et al., 1995; Uhlig et al., 2000; Jansen et al., 2001; Nordahl et al., 2001; Voog et al., 2003a; Syversen et al., 2008; Thabet et al., 2009) where C-reactive protein, erythrocyte sedimentation rate as well as rheumatoid factor were related to articular bone tissue resorption. The reason for why the current study was not able to identify any predictive inflammatory markers for progression of TMJ bone tissue resorption is unclear but may be due to dual character of bone tissue

resorption, i.e. inflammatory versus non-inflammatory. Another explanation may be the general treatment before the first visit.

General discussion

The main finding of this study is that TMJ bone tissue resorption is present early after diagnosis of RA. The high frequency of TMJs with bone tissue resorption indicates that a subclinical process is already present at diagnosis of the general disease. The etiology of the bone tissue resorption may comprise both inflammatory and non-inflammatory components. Another major finding is that a large proportion of the patients showed regression of TMJ bone tissue resorption during the first two years of RA, which may be due to early and aggressive treatment but differs from earlier studies.

Articular tissue destruction in RA comprises both cartilage degradation and bone tissue resorption. In the present study, the early glutamate-dependent bone tissue resorption was found when TMJ crepitus was absent, indicating a process underneath the articular cartilage (Appel et al., 2006; Appel et al., 2009). The cartilage degradation is most probably of inflammatory nature whereas the bone tissue resorption may also be due to non-inflammatory processes like periarticular osteopenia, which in turn may be due to a deficiency in sex hormones (Gough et al., 1994; Sambrook, 2000; Vanderschueren et al., 2004; Appel et al., 2006; Jimenez-Boj et al., 2007; Schett, 2008; Study I; Schett and Sieper, 2009).

The positive relationship between glutamate and TMJ erosion score at the early stage was found in patients without crepitus, i.e. with intact cartilage. Glutamate may also be involved in bone tissue resorption through systemic inflammatory mechanisms, since increased crepitus was associated with increased erythrocyte sedimentation rate only in patients with markedly increased glutamate. There are indications of a dual role of glutamate regarding bone tissue turnover as glutamate regulates both bone tissue resorption and formation but this study was solely concerned with bone resorption. The biologic mechanisms are scarcely known but the glutamate NMDA receptor on preosteoclasts stimulates osteoclastogenesis, i.e. bone tissue resorption (Chenu et al., 1998). The findings in the current project indicate that glutamate modulates bone tissue resorption and probably cartilage degradation but through different mechanisms in early RA.

Regression of articular bone tissue resorption, as assessed by radiography, has not been shown for other RA joints. One explanation may be differences in radiographic methodology. The current study aimed at register erosions and did not include assessment of total bone

tissue loss or other morphological changes. This means that while the erosions decreased, probably as a consequence of the early anti-inflammatory treatment, a progression of the total bone loss might still have occurred.

Regression of TMJ bone tissue resorption was more frequent (43%) in the current study than in a previous study of late phase TMJ RA (24%; Nordahl et al., 2001). In that study, the follow-up interval was one year whereas it was two years in the current project. Another explanation to the high regression frequency in the current study may be the early and aggressive treatment. Patients who failed to respond to methotrexate monotherapy, as judged by disease activity ($DAS \geq 3.2$), were given additional sulfasalazine and hydroxychloroquine or anti-TNF therapy as early as three months after diagnosis. Synovitis in a particular joint is associated with progression of erosions and joint space narrowing but not in patients treated early with infliximab (Klarenbeek et al., 2010). Anti-TNF therapy is generally associated with inhibition of joint destruction in RA (Aletaha et al., 2007; Tanino et al., 2009) and reduction in disease activity (Aletaha et al., 2007; van der Heijde et al., 2007). In the current study 15% of the patients were on treatment with infliximab or etanercept at the 2 year follow-up. Another explanation to the regression found in the current study, although of less importance, may be that 7% of the patients with clinical signs of TMJ arthritis or radiographic erosions at the first visit and 17% at the one-year follow-up received intra-articular injection of glucocorticoid. The individual reduction in erosion score over time may therefore, at least partly, be explained by the response to the early and specific treatment (Quinn and Emery, 2003; Nell et al., 2004; van Vollenhoven et al., 2009).

The high frequency of TMJ bone tissue resorption (72%) found in Study I compared to other studies may be explained by the advanced radiographic technique used. Erosions may be detected earlier by CT because of its ability to clearly delineate cortical bony margins (Perry et al., 2005). CT has been considered valuable for detecting and monitoring articular bone erosions in RA as the sensitivity and reproducibility is higher than plain radiographs for detecting bone erosions in metacarpophalangeal joints as well as in the TMJ (Goupille et al., 1990; Larheim and Kolbenstvedt, 1990; Dohn et al., 2007). Cone-beam CT was chosen because of its low exposure of radiation (Mozzo et al., 1998; Danforth et al., 2003; Hashimoto et al., 2003; Ludlow et al., 2003; Mah et al., 2003; Scarfe and Farman, 2008). Cone-beam CT is at least as reliable

as conventional CT in diagnostics of TMJ bone tissue changes but provide superior reliability and greater accuracy than conventional tomograms in detecting TMJ erosions (Honda et al., 2006; Honey et al., 2007).

The patients included were diagnosed with RA shortly before their first visit in the current project; most of them were examined for this study within less than one month after diagnosis and about 6 months after their report of debut of symptoms. Most patients began treatment with DMARDs promptly after diagnosis, i.e. before the start of this investigation, and many of these also used NSAIDs. A reduced systemic and local inflammatory activity by this early medication is likely. However, the influence of the medication on the results from the first visit regarding radiographic signs of bone tissue resorption can be expected to be minor or even insignificant.

The level of plasma glutamate was lower in the patients at the first visit than in healthy individuals, which may be due to the early treatment. However, higher levels of glutamate with a median 7.0 mmol/L were found in patients with low C-reactive protein and estradiol, in whom glutamate was most strongly associated with TMJ bone tissue resorption. One explanation may be that glutamate participates in local rather than systemic inflammatory mechanisms of bone tissue resorption. However, it has been shown that the glutamate level in TMJ synovial fluid is associated with local inflammation, as expressed by local TNF concentration, despite low systemic inflammatory activity (Alstergren et al., 2007; Clarke and Khosla, 2010).

The patients of our study must be considered as representative for the current RA patient population in Sweden since 85% of the patients were included in BARFOT, a Swedish multicenter registry project aimed to observe clinical, serological, radiographic and treatment data longitudinally. The BARFOT project includes all newly diagnosed RA patients from six rheumatologic centers in Sweden and should therefore be a suitable reference population. However, the systemic inflammatory activity, as assessed by C-reactive protein, was lower in our patient sample. The probable reason for this difference is the time of a median of 23 days from the start of medication to the first visit. The frequency of TMJ pain at rest was only 13% in our study which may partly explain why rather few of the invited patients, 47 out of 380, were willing to participate in this particular study.

Conclusions

Cross-sectional part (I and II)

This part of the study indicated that

- a majority of patients with early RA present with radiographic signs of bone tissue resorption of the TMJ (Study I)
- circulating glutamate is associated with the extent of TMJ bone tissue resorption (Study I)
- the relationship between glutamate and TMJ bone tissue resorption seems to be modulated by systemic inflammatory activity, estradiol or testosterone levels (Study I)
- presence of TMJ bone tissue resorption can be explained by a combination of TMJ crepitus, elevated plasma level of glutamate and systemic inflammatory activity (Study II)
- the relation between glutamate and TMJ bone tissue resorption is different in patients with or without crepitus indicating that glutamate has a particular role in joints without cartilage destruction (Study II)
- bone tissue resorption at this early stage is related to the inflammatory mediator serotonin in men but not in women (Study II)

Longitudinal part (III and IV)

The results of this part of the study showed that during the first two years of RA

- TMJ bone tissue resorption can be expected to regress in most patients, however, progression can be expected in a considerable number (Study III)
- progression of TMJ bone tissue resorption seems to be associated with IL-1 β related inflammatory mechanisms as well as non-inflammatory mechanisms involving testosterone (Study III)
- progression of TMJ crepitus seems to be associated with inflammation (increase of erythrocyte sedimentation rate) under influence of glutamate (in patients with high increase of glutamate levels) (Study III)

- regression of TMJ bone tissue resorption is predicted by early high TMJ erosion score (Study IV)
- regression of TMJ bone tissue resorption is predicted by TMJ crepitus (Study IV)
- regression of TMJ bone tissue resorption is predicted by high glutamate levels in plasma and in the men by high levels of serotonin (Study IV)

Populärvetenskaplig sammanfattning

Ledgångsreumatism är en kronisk, systemisk, inflammatorisk sjukdom som angriper och bryter ner kroppens leder. Sjukdomen förekommer hos ca 0,5-1 % av befolkningen och drabbar oftare kvinnor än män. Orsaken till reumatism är inte helt känd men sjukdomen kan bli väldigt handikappande med trötthet, smärta och nedsatt rörlighet som vanliga kännetecken när den är etablerad. Många faktorer har nämnts som orsak däribland genetiska och miljöfaktorer. Sannolikt är det en kombination av flera. Man tror även att könshormoner inverkar eftersom sjukdomen främst drabbar kvinnor efter klimakteriet.

Diagnosen ledgångsreumatism ställs utifrån kliniskt ledstatus, röntgen och blodprov som visar särskilda sjukdomsmarkörer. Tidiga tecken på bennedbrytning i lederna på röntgenbilder anses vara en indikation på dålig prognos. Därför utgör ledröntgen en viktig del i diagnostiken men även i uppföljningen sjukdomen. Idag vet man att tidigt fastställd diagnos och insatt behandling (både lokal i den enskilda leden och generell) är avgörande för det fortsatta sjukdomsförloppet. Ju tidigare och effektivare behandling desto större är chansen att bromsa och i bästa fall stanna upp ledernas fortsatta nedbrytning.

Ungefär 50 % av reumatikerna drabbas av reumatism i käkleden med smärta och funktionsnedsättning som följd. Käkledsengagemanget är handikappande av många skäl då det påverkar tuggning och tal. Ny kunskap har visat att leddestruktion kan fortgå trots att man inte har några symptom eller kliniska tecken på inflammation. Även om det inte finns några entydiga resultat så har det visat sig från andra leder att flera inflammatoriska ämnen eller markörer kan vara av diagnostisk och prognostisk betydelse för den fortsatta utvecklingen. Vi har studerat glutamat, serotonin, TNF, IL-1 β , IL-6, VEGF, C-reaktivt protein, sänkingsreaktionen, reumatoid faktor samt könshormonerna östradiol och testosteron.

Det övergripande målet med denna avhandling var att ta redan på i vilken utsträckning det förekommer röntgenförändringar i käkleden tidigt i sjukdomsförloppet, hur dessa utvecklas över tid och om de indikerar en sämre prognos.

Avhandlingen består av två delar: en tvärsnitts studie och en longitudinell studie. Målet med tvärsnittsstudien var att relatera röntgenfynden till inflammatoriska ämnen och hormoner i blodet samt

kliniska tecken på nedbrytning i käkled. Målen med den longitudinella studien var att reda ut hur förändringar på röntgen kan relateras till förändringar avseende inflammatoriska ämnen och hormoner samt kliniska tecken.

Den första delen av avhandlingen omfattade 47 patienter och den andra 41, som undersöktes med avseende på röntgenförändringar, käkledsljud, inflammatoriska ämnen i blod och könshormoner. Den första undersökningen genomfördes ca tre veckor efter fastställande av diagnosen ledgångsreumatism. Därefter upprepades denna undersökning efter ett och två år. Analys av röntgenförändringar, s.k. erosioner, utfördes på bilder tagna med dator tomografi. Resultaten från första delen visade att 72 % av patienterna uppvisade erosioner på röntgen kort efter att diagnosen ställts. Erosionerna var relaterade till det inflammatoriska ämnet glutamat och krepitationer men också till markörer för inflammatorisk aktivitet. Sambandet mellan glutamat och erosioner var starkare vid frånvaro av krepitationer, vid låg inflammatorisk aktivitet och låga nivåer av kvinnligt eller manligt könshormon. Resultaten från andra delen visade att 29 % av patienterna hade fortsatt nedbrytning av leden medan 43 % av patienterna uppvisade en minskning. Fortsatt nedbrytning kunde delvis förklaras med ökningen av en inflammations substans som är vanligt förekommande i samband med inflammatoriskt betingad bennedbrytning, interleukin-1 β . Den förklarades också av en ökning av det manliga könshormonet, testosteron. Det tycks vara så att tidigt uppkomna erosioner eller högre glutamat nivåer leder till en minskad nedbrytning. Sannolikt är dessa fynd relaterade till tidigt insatt och effektiv behandling.

Sammanfattningsvis indikerar resultaten från denna avhandling att tidiga tecken på bennedbrytning i käkleden hos reumatiker är vanligt förekommande. Bennedbrytningen tycks dock kunna minska om den upptäcks och behandlas tidigt. Tidiga röntgen tecken tycks vara relaterade till glutamat förekomst särskilt vid låg inflammatorisk aktivitet och i frånvaro av kliniska tecken. Däremot tycks tidiga röntgen tecken eller glutamat vara positivt för en återgång under de två första åren efter diagnosen ledgångsreumatism.

Acknowledgements

To complete a thesis would be impossible without the excellent contribution from a number of people. I would therefore like to express my sincere gratitude to all of you who made my project come true. In particular, I would like to thank the following persons:

Professor **Sigvard Kopp**, my main supervisor, for your excellent scientific guidance and never-ending support in the increasing challenge being a scientist. And most of all your humanity which reflects the non-prestigious atmosphere of your research group where there is plenty of space to grow.

Associate Professor **Per Alstergren**, my co-supervisor, for always being available, willing to share your time and your truly support. I am even more grateful that you kept me on the track and believed in me at times when I did not.

Dr. **Fredrik Bergstrand**, my external mentor, reminding me about the privilege being a clinically acting orthodontist and that there is a life waiting out there when I got lost in the scientific analyses.

My co-authors: **Karin Näsström** for your never-ending enthusiasm and for all hours you spent on analysing the CBCT images with me and **Johan Bratt** for your very kind assistance and support in the recruitment process of the patients.

Associate Professor **Malin Ernberg** for your friendship and support. You have been a true source of inspiration with your positive attitude and solution-oriented way of thinking.

Professor **Lennart Lagerström** for providing me with keys to my post-doctoral life. The doors that you enabled me to open made any effort of my doctoral studies worthy. My hope is to bring back new inspiration to the profession of orthodontics with help of these keys.

Mats Trulsson my department head for providing me with conclusive perspectives.

My dear friends and colleagues in the **research group** at the Division of Clinical Oral Physiology; Nicolaos Christidis, Sofia Louca, Ola Bjärbo, Lars Fredriksson, Mikael Nilsson and Britt Hedenberg-Magnusson. It has been a sincere privilege to work in your company. Thank you for all joyful moments.

The assisting **personnel at the department of rheumatology** providing me with patients. The **staff at the department of oral radiology** for help with the radiographic exams. **Karin Trollsås** and **Agneta Gustafsson** for your skilful laboratory work, the dental assistants **Marie-Louise Bjerdahl** and **Mira Lukenic**, for your excellent assistance and flexibility.

To the **researchers** and **PhD-students** at the Department of Dental Medicine; Janina Strauss, Anders Sjögren, Ann-Sofie Johansen, Sarah Rickardsson, Lena Karlsson, Jari Taghavi and Tülay Lindbeg for being such a great support and company on the way.

All patients participating in the project for offering your time and effort which is a true challenge for everyone with RA.

My parents **Lars** and **Gun-Britt Mauritzson** from who I learnt that nothing in life is for granted. My brave hearted dear brother **Josef** for having you in my life.

My husband and dearest friend **Reza** for your endless faith and love.

My daughter **Alexandra** for the precious young lady you are and for your patience of the never ending work that kept your mother away from you. Sometimes even more than a child deserves.

This is dedicated to the two of you.

Grants

The project was financially supported by the National Institute of Dental and Craniofacial Research, the Department of Dental Medicine at Karolinska Institutet, the Swedish Dental Society, Folk tandvården AB Public Stockholm County Council, American Dental Society of Sweden and Ortodonti Akademin Sverige AB.

References

- Achen MG, Stacker SA. The vascular endothelial growth factor family; proteins which guide the development of the vasculature. *Int J Exp Pathol* 1998;79:255-265.
- Aho K, Heliövaara M, Maatela J, Tuomi T, Palosuo T. Rheumatoid factors antedating clinical rheumatoid arthritis. *J Rheumatol* 1991;18:1282-1284.
- Aho K, von Essen R, Kurki P, Palosuo T, Heliövaara M. Antikeratin antibody and antiperinuclear factor as markers for subclinical rheumatoid disease process. *J Rheumatol* 1993;20:1278-1281.
- Ainola M, Mandelin J, Liljeström M, Konttinen YT, Salo J. Imbalanced expression of RANKL and osteoprotegerin mRNA in pannus tissue of rheumatoid arthritis. *Clin Exp Rheumatol* 2008;26:240-246.
- Akerman S. Morphologic, radiologic and thermometric assessment of degenerative and inflammatory temporomandibular joint disease. an autopsy and clinical study. *Swed Dent J Suppl* 1987;52:1-110.
- Akerman S, Kopp S, Nilner M, Petersson A, Rohlin M. Relationship between clinical and radiologic findings of the temporomandibular joint in rheumatoid arthritis. *Oral Surg Oral Med Oral Pathol* 1988;66:639-643.
- Akerman S, Jonsson K, Kopp S, Petersson A, Rohlin M. Radiologic changes in temporomandibular, hand, and foot joints of patients with rheumatoid arthritis. *Oral Surg Oral Med Oral Pathol* 1991;72:245-250.
- Alamanos Y, Voulgari PV, Drosos AA. Incidence and prevalence of rheumatoid arthritis, based on the 1987 american college of rheumatology criteria: A systematic review. *Semin Arthritis Rheum* 2006;36:182-188.
- Aletaha D, Funovits J, Keystone EC, Smolen JS. Disease activity early in the course of treatment predicts response to therapy after one year in rheumatoid arthritis patients. *Arthritis Rheum* 2007;56:3226-3235.
- Alfredson H, Forsgren S, Thorsen K, Lorentzon R. In vivo microdialysis and immunohistochemical analyses of tendon tissue demonstrated high amounts of free glutamate and glutamate NMDAR1 receptors, but no signs of inflammation, in jumper's knee. *Journal of Orthopaedic Research* 2001;19:881-886.
- Aliprandi A, Longoni M, Stanzani L, Tremolizzo L, Vaccaro M, Begni B, Galimberti G, Garofolo R, Ferrarese C. Increased plasma glutamate in stroke patients might be linked to altered platelet release and uptake. *J Cereb Blood Flow Metab* 2005;25:513-519.
- Alstergren P, Hajati A, Sessle B, Kopp S. Glutamate in Temporomandibular Joint Synovial Fluid and Plasma. 2007; Abstract International Association for Dental Research General Meeting, New Orleans, LO, USA.
- Alstergren P, Kopp S. Pain and synovial fluid concentration of serotonin in arthritic temporomandibular joints. *Pain* 1997;72:137-143.
- Alstergren P, Ernberg M, Kvarnström M, Kopp S. Interleukin-1beta in synovial fluid from the arthritic temporomandibular joint and its relation to pain,

- mobility, and anterior open bite. *J Oral Maxillofac Surg* 1998;56:1059-65; discussion 1066.
- Alstergren P, Ernberg M, Kopp S, Lundeberg T, Theodorsson E. TMJ pain in relation to circulating neuropeptide Y, serotonin, and interleukin-1 beta in rheumatoid arthritis. *J Orofac Pain* 1999;13:49-55.
- Amos RS, Constable TJ, Crockson RA, Crockson AP, McConkey B. Rheumatoid arthritis: Relation of serum C-reactive protein and erythrocyte sedimentation rates to radiographic changes. *Br Med J* 1977;1:195-197.
- Appel H, Kuhne M, Spiekermann S, Kohler D, Zacher J, Stein H, Sieper J, Loddenkemper C. Immunohistochemical analysis of hip arthritis in ankylosing spondylitis: Evaluation of the bone-cartilage interface and subchondral bone marrow. *Arthritis Rheum* 2006;54:1805-1813.
- Appel H, Loddenkemper C, Miossec P. Rheumatoid arthritis and ankylosing spondylitis - pathology of acute inflammation. *Clin Exp Rheumatol* 2009;27:S15-9.
- Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper NS, Healey LA, Kaplan SR, Liang MH, Luthra HS. The american rheumatism association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988;31:315-324.
- Ballara S, Taylor PC, Reusch P, Marme D, Feldmann M, Maini RN, Paleolog EM. Raised serum vascular endothelial growth factor levels are associated with destructive change in inflammatory arthritis. *Arthritis Rheum* 2001;44:2055-2064.
- Battaglino R, Fu J, Spate U, Ersoy U, Joe M, Sedaghat L, Stashenko P. Serotonin regulates osteoclast differentiation through its transporter. *J Bone Miner Res* 2004;19:1420-1431.
- Bellido T, Stahl N, Farruggella TJ, Borba V, Yancopoulos GD, Manolagas SC. Detection of receptors for interleukin-6, interleukin-11, leukemia inhibitory factor, oncostatin M, and ciliary neurotrophic factor in bone marrow stromal/osteoblastic cells. *J Clin Invest* 1996;97:431-437.
- Blizotes M, Eshleman A, Burt-Pichat B, Zhang XW, Hashimoto J, Wiren K, Chenu C. Serotonin transporter and receptor expression in osteocytic MLO-Y4 cells. *Bone* 2006;39:1313-1321.
- Blizotes MM, Eshleman AJ, Zhang XW, Wiren KM. Neurotransmitter action in osteoblasts: Expression of a functional system for serotonin receptor activation and reuptake. *Bone* 2001;29:477-486.
- Boini S, Guillemin F. Radiographic scoring methods as outcome measures in rheumatoid arthritis: Properties and advantages. *Ann Rheum Dis* 2001;60:817-827.
- Boonen A, van der Heijde D. Conventional x-ray in early arthritis. *Rheum Dis Clin North Am* 2005;31:681-698.
- Brown LF, Detmar M, Claffey K, Nagy JA, Feng D, Dvorak AM, Dvorak HF. Vascular permeability factor/vascular endothelial growth factor: A multifunctional angiogenic cytokine. *EXS* 1997;79:233-269.

- Buchan G, Barrett K, Turner M, Chantry D, Maini RN, Feldmann M. Interleukin-1 and tumour necrosis factor mRNA expression in rheumatoid arthritis: Prolonged production of IL-1 alpha. *Clin Exp Immunol* 1988;73:449-455.
- Bull BS, Westengard JC, Farr M, Bacon PA, Meyer PJ, Stuart J. Efficacy of tests used to monitor rheumatoid arthritis. *Lancet* 1989;2:965-967.
- Celiker R, Gokce-Kutsal Y, Eryilmaz M. Temporomandibular joint involvement in rheumatoid arthritis. relationship with disease activity. *Scand J Rheumatol* 1995;24:22-25.
- Chenu C, Serre CM, Raynal C, Burt-Pichat B, Delmas PD. Glutamate receptors are expressed by bone cells and are involved in bone resorption. *Bone* 1998;22:295-299.
- Chenu C. Glutamatergic regulation of bone resorption. *J Musculoskelet Neuronal Interact* 2002a;2:423-431.
- Chenu C. Glutamatergic innervation in bone. *Microsc Res Tech* 2002b;58:70-76.
- Chenu C. Glutamatergic regulation of bone remodeling. *J Musculoskelet Neuronal Interact* 2002c;2:282-284.
- Clarke BL, Khosla S. Physiology of bone loss. *Radiol Clin North Am* 2010;48:483-495.
- Collet C, Schiltz C, Geoffroy V, Maroteaux L, Launay JM, de Vernejoul MC. The serotonin 5-HT_{2B} receptor controls bone mass via osteoblast recruitment and proliferation. *FASEB J* 2008;22:418-427.
- Combe B, Eliaou JF, Dures JP, Meyer O, Clot J, Sany J. Prognostic factors in rheumatoid arthritis. comparative study of two subsets of patients according to severity of articular damage. *Br J Rheumatol* 1995;34:529-534.
- Crotti TN, Smith MD, Weedon H, Ahern MJ, Findlay DM, Kraan M, Tak PP, Haynes DR. Receptor activator NF-kappaB ligand (RANKL) expression in synovial tissue from patients with rheumatoid arthritis, spondyloarthropathy, osteoarthritis, and from normal patients: Semiquantitative and quantitative analysis. *Ann Rheum Dis* 2002;61:1047-1054.
- Cutolo M, Wilder RL. Different roles for androgens and estrogens in the susceptibility to autoimmune rheumatic diseases. *Rheum Dis Clin North Am* 2000;26:825-839.
- Cutolo M, Sulli A, Capellino S, Villaggio B, Montagna P, Seriolo B, Straub RH. Sex hormones influence on the immune system: Basic and clinical aspects in autoimmunity. *Lupus* 2004;13:635-638.
- Cutolo M, Otsa K, Aakre O, Sulli A. Nocturnal hormones and clinical rhythms in rheumatoid arthritis. *Ann N Y Acad Sci* 2005;1051:372-381.
- Cutolo M, Capellino S, Sulli A, Serioli B, Secchi ME, Villaggio B, Straub RH. Estrogens and autoimmune diseases. *Ann N Y Acad Sci* 2006;1089:538-547.
- Cutolo M, Straub RH, Bijlsma JW. Neuroendocrine-immune interactions in synovitis. *Nat Clin Pract Rheumatol* 2007;3:627-634.
- Cutolo M. Androgens in rheumatoid arthritis: When are they effectors? *Arthritis Res Ther* 2009;11:126.

- Danforth RA, Dus I, Mah J. 3-D volume imaging for dentistry: A new dimension. *J Calif Dent Assoc* 2003;31:817-823.
- Danning CL, Illei GG, Hitchon C, Greer MR, Boumpas DT, McInnes IB. Macrophage-derived cytokine and nuclear factor kappaB p65 expression in synovial membrane and skin of patients with psoriatic arthritis. *Arthritis Rheum* 2000;43:1244-1256.
- Dao TT, LeResche L. Gender differences in pain. *J Orofac Pain* 2000;14:169-84; discussion 184-95.
- de Bandt M, Grossin M, Weber AJ, Chopin M, Elbim C, Pla M, Gougerot-Pocidal MA, Gaudry M. Suppression of arthritis and protection from bone destruction by treatment with TNP-470/AGM-1470 in a transgenic mouse model of rheumatoid arthritis. *Arthritis Rheum* 2000;43:2056-2063.
- Dinarello CA. Biologic basis for interleukin-1 in disease. *Blood* 1996;87:2095-2147.
- Dohn UM, Ejbjerg BJ, Hasselquist M, Narvestad E, Court-Payen M, Szkudlarek M, Moller J, Thomsen HS, Ostergaard M. Rheumatoid arthritis bone erosion volumes on CT and MRI: Reliability and correlations with erosion scores on CT, MRI and radiography. *Ann Rheum Dis* 2007;66:1388-1392.
- Duff GW. Cytokines and acute phase proteins in rheumatoid arthritis. *Scand J Rheumatol Suppl* 1994;100:9-19.
- Eastgate JA, Symons JA, Wood NC, Grinlinton FM, di Giovine FS, Duff GW. Correlation of plasma interleukin 1 levels with disease activity in rheumatoid arthritis. *Lancet* 1988;2:706-709.
- Ejbjerg BJ, Vestergaard A, Jacobsen S, Thomsen H, Ostergaard M. Conventional radiography requires a MRI-estimated bone volume loss of 20% to 30% to allow certain detection of bone erosions in rheumatoid arthritis metacarpophalangeal joints. *Arthritis Res Ther* 2006;8:R59.
- Endresen GK. Evidence for activation of platelets in the synovial fluid from patients with rheumatoid arthritis. *Rheumatol Int* 1989;9:19-24.
- Englund M, Joud A, Geborek P, Felson DT, Jacobsson LT, Petersson IF. Prevalence and incidence of rheumatoid arthritis in southern sweden 2008 and their relation to prescribed biologics. *Rheumatology (Oxford)* 2010;49:1563-1569.
- Fava RA, Olsen NJ, Spencer-Green G, Yeo KT, Yeo TK, Berse B, Jackman RW, Senger DR, Dvorak HF, Brown LF. Vascular permeability factor/endothelial growth factor (VPF/VEGF): Accumulation and expression in human synovial fluids and rheumatoid synovial tissue. *J Exp Med* 1994;180:341-346.
- Flood S, Parri R, Williams A, Duance V, Mason D. Modulation of interleukin-6 and matrix metalloproteinase 2 expression in human fibroblast-like synoviocytes by functional ionotropic glutamate receptors. *Arthritis Rheum* 2007;56:2523-2534.
- Forslind K, Ahlmen M, Eberhardt K, Hafstrom I, Svensson B, BARFOT Study Group. Prediction of radiological outcome in early rheumatoid arthritis in clinical practice: Role of antibodies to citrullinated peptides (anti-CCP). *Ann Rheum Dis* 2004;63:1090-1095.

- Forslind K, Boonen A, Albertsson K, Hafstrom I, Svensson B, Barfot Study Group. Hand bone loss measured by digital X-ray radiogrammetry is a predictor of joint damage in early rheumatoid arthritis. *Scand J Rheumatol* 2009;38:431-438.
- Fredriksson L, Alstergren P, Kopp S. Serotonergic mechanisms influence the response to glucocorticoid treatment in TMJ arthritis. *Mediators Inflamm* 2005;2005:194-201.
- Galal N, El Beialy W, Deyama Y, Yoshimura Y, Yoshikawa T, Suzuki K, Totsuka Y. Effect of estrogen on bone resorption and inflammation in the temporomandibular joint cellular elements. *Int J Mol Med* 2008;21:785-790.
- Genever PG, Skerry TM. Regulation of spontaneous glutamate release activity in osteoblastic cells and its role in differentiation and survival: Evidence for intrinsic glutamatergic signaling in bone. *FASEB J* 2001;15:1586-1588.
- Geusens PP, Landewe RB, Garnero P, Chen D, Dunstan CR, Lems WF, Stinissen P, van der Heijde DM, van der Linden S, Boers M. The ratio of circulating osteoprotegerin to RANKL in early rheumatoid arthritis predicts later joint destruction. *Arthritis Rheum* 2006;54:1772-1777.
- Girasole G, Jilka RL, Passeri G, Boswell S, Boder G, Williams DC, Manolagas SC. 17 beta-estradiol inhibits interleukin-6 production by bone marrow-derived stromal cells and osteoblasts in vitro: A potential mechanism for the antiosteoporotic effect of estrogens. *J Clin Invest* 1992;89:883-891.
- Goekoop-Ruiterman YP, de Vries-Bouwstra JK, Allaart CF, van Zeven D, Kerstens PJ, Hazes JM, Zwinderman AH, Ronday HK, Han KH, Westedt ML, Gerards AH, van Groenendael JH, Lems WF, van Krugten MV, Breedveld FC, Dijkmans BA. Clinical and radiographic outcomes of four different treatment strategies in patients with early rheumatoid arthritis (the BeSt study): A randomized, controlled trial. *Arthritis Rheum* 2005;52:3381-3390.
- Gordon D, Beastall GH, Thomson JA, Sturrock RD. Prolonged hypogonadism in male patients with rheumatoid arthritis during flares in disease activity. *Br J Rheumatol* 1988;27:440-444.
- Gough AK, Lilley J, Eyre S, Holder RL, Emery P. Generalised bone loss in patients with early rheumatoid arthritis. *Lancet* 1994;344:23-27.
- Goupille P, Fouquet B, Cotty P, Goga D, Mateu J, Valat JP. The temporomandibular joint in rheumatoid arthritis. correlations between clinical and computed tomography features. *J Rheumatol* 1990;17:1285-1291.
- Goupille P, Fouquet B, Goga D, Cotty P, Valat JP. The temporomandibular joint in rheumatoid arthritis: Correlations between clinical and tomographic features. *J Dent* 1993;21:141-146.
- Graudal N. The natural history and prognosis of rheumatoid arthritis: Association of radiographic outcome with process variables, joint motion and immune proteins. *Scand J Rheumatol Suppl* 2004;(118):1-38.
- Gravallese EM, Manning C, Tsay A, Naito A, Pan C, Amento E, Goldring SR. Synovial tissue in rheumatoid arthritis is a source of osteoclast differentiation factor. *Arthritis Rheum* 2000;43:250-258.
- Gravallese EM. Bone destruction in arthritis. *Ann Rheum Dis* 2002;61:84-6.

- Gu Y, Genever PG, Skerry TM, Publicover SJ. The NMDA type glutamate receptors expressed by primary rat osteoblasts have the same electrophysiological characteristics as neuronal receptors. *Calcif Tissue Int* 2002;70:194-203.
- Gustafsson BI, Thommesen L, Stunes AK, Tommeras K, Westbroek I, Waldum HL, Slordahl K, Tamburstuen MV, Reseland JE, Syversen U. Serotonin and fluoxetine modulate bone cell function in vitro. *J Cell Biochem* 2006;98:139-151.
- Gynther GW, Tronje G, Holmlund AB. Radiographic changes in the temporomandibular joint in patients with generalized osteoarthritis and rheumatoid arthritis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1996;81:613-618.
- Harada M, Mitsuyama K, Yoshida H, Sakisaka S, Taniguchi E, Kawaguchi T, Ariyoshi M, Saiki T, Sakamoto M, Nagata K, Sata M, Matsuo K, Tanikawa K. Vascular endothelial growth factor in patients with rheumatoid arthritis. *Scand J Rheumatol* 1998;27:377-380.
- Hashimoto A, Sato H, Nishibayahi Y, Shiino Y, Kutsuna T, Ishihara Y, Hoshi K, Fujimori J, Tsuboi S, Kondo H, Akizuki M, Moroi Y, Yoshida S. A multicenter cross-sectional study on the health related quality of life of patients with rheumatoid arthritis using a revised japanese version of the arthritis impact measurement scales version 2 (AIMS 2), focusing on physical disability and its associative factors. *Ryumachi* 2001;41:9-24.
- Hashimoto K, Arai Y, Iwai K, Araki M, Kawashima S, Terakado M. A comparison of a new limited cone beam computed tomography machine for dental use with a multidetector row helical CT machine. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2003;95:371-377.
- Hayer S, Redlich K, Korb A, Hermann S, Smolen J, Schett G. Tenosynovitis and osteoclast formation as the initial preclinical changes in a murine model of inflammatory arthritis. *Arthritis Rheum* 2007;56:79-88.
- Haynes DR, Barg E, Crotti TN, Holding C, Weedon H, Atkins GJ, Zannettino A, Ahern MJ, Coleman M, Roberts-Thomson PJ, Kraan M, Tak PP, Smith MD. Osteoprotegerin expression in synovial tissue from patients with rheumatoid arthritis, spondyloarthropathies and osteoarthritis and normal controls. *Rheumatology (Oxford)* 2003;42:123-134.
- Herbert MK, Schmidt RF. Activation of normal and inflamed fine articular afferent units by serotonin. *Pain* 1992;50:79-88.
- Ho ML, Tsai TN, Chang JK, Shao TS, Jeng YR, Hsu C. Down-regulation of N-methyl D-aspartate receptor in rat-modeled disuse osteopenia. *Osteoporos Int* 2005;16:1780-1788.
- Hofbauer LC, Lacey DL, Dunstan CR, Spelsberg TC, Riggs BL, Khosla S. Interleukin-1beta and tumor necrosis factor-alpha, but not interleukin-6, stimulate osteoprotegerin ligand gene expression in human osteoblastic cells. *Bone* 1999;25:255-259.
- Hollmann M, Heinemann S. Cloned glutamate receptors. *Annu Rev Neurosci* 1994;17:31-108.

- Honda K, Larheim TA, Maruhashi K, Matsumoto K, Iwai K. Osseous abnormalities of the mandibular condyle: Diagnostic reliability of cone beam computed tomography compared with helical computed tomography based on an autopsy material. *Dentomaxillofac Radiol* 2006;35:152-157.
- Honey OB, Scarfe WC, Hilgers MJ, Klueber K, Silveira AM, Haskell BS, Farman AG. Accuracy of cone-beam computed tomography imaging of the temporomandibular joint: Comparisons with panoramic radiology and linear tomography. *Am J Orthod Dentofacial Orthop* 2007;132:429-438.
- Itzstein C, Cheynel H, Burt-Pichat B, Merle B, Espinosa L, Delmas PD, Chenu C. Molecular identification of NMDA glutamate receptors expressed in bone cells. *J Cell Biochem* 2001;82:134-144.
- Jackson JR, Bolognese B, Kircher CH, Marshall LA, Winkler JD. Modulation of angiogenesis in a model of chronic inflammation. *Inflamm Res* 1997;46 Suppl 2:S129-30.
- Jansen LM, van der Horst-Bruinsma IE, van Schaardenburg D, Bezemer PD, Dijkmans BA. Predictors of radiographic joint damage in patients with early rheumatoid arthritis. *Ann Rheum Dis* 2001;60:924-927.
- Jimenez-Boj E, Nobauer-Huhmann I, Hanslik-Schnabel B, Dorotka R, Wanivenhaus AH, Kainberger F, Trattinig S, Axmann R, Tsuji W, Hermann S, Smolen J, Schett G. Bone erosions and bone marrow edema as defined by magnetic resonance imaging reflect true bone marrow inflammation in rheumatoid arthritis. *Arthritis Rheum* 2007;56:1118-1124.
- Jimi E, Nakamura I, Duong LT, Ikebe T, Takahashi N, Rodan GA, Suda T. Interleukin 1 induces multinucleation and bone-resorbing activity of osteoclasts in the absence of osteoblasts/stromal cells. *Exp Cell Res* 1999;247:84-93.
- Jochems C, Islander U, Erlandsson M, Verdrengh M, Ohlsson C, Carlsten H. Osteoporosis in experimental postmenopausal polyarthritis: The relative contributions of estrogen deficiency and inflammation. *Arthritis Res Ther* 2005;7:R837-43.
- Jones SM, Bhalla AK. Osteoporosis in rheumatoid arthritis. *Clin Exp Rheumatol* 1993;11:557-562.
- Jung YO, Cho ML, Lee SY, Oh HJ, Park JS, Park MK, Park MJ, Ju JH, Kim SI, Park SH, Kim HY, Min JK. Synergism of toll-like receptor 2 (TLR2), TLR4, and TLR6 ligation on the production of tumor necrosis factor (TNF)-alpha in a spontaneous arthritis animal model of interleukin (IL)-1 receptor antagonist-deficient mice. *Immunol Lett* 2009;123:138-143.
- Khalkhali-Ellis Z, Seftor EA, Nieva DR, Handa RJ, Price RH, Jr, Kirschmann DA, Baragi VM, Sharma RV, Bhalla RC, Moore TL, Hendrix MJ. Estrogen and progesterone regulation of human fibroblast-like synoviocyte function in vitro: Implications in rheumatoid arthritis. *J Rheumatol* 2000;27:1622-1631.
- Khosla S, Melton LJ, 3rd, Atkinson EJ, O'Fallon WM. Relationship of serum sex steroid levels to longitudinal changes in bone density in young versus elderly men. *J Clin Endocrinol Metab* 2001;86:3555-3561.
- Klarenbeek NB, Guler-Yuksel M, van der Heijde DM, Hulsmans HM, Kerstens PJ, Molenaar TH, de Sonnaville PB, Huizinga TW, Dijkmans BA, Allaart CF. Clinical

synovitis in a particular joint is associated with progression of erosions and joint space narrowing in that same joint, but not in patients initially treated with infliximab. *Ann Rheum Dis* 2010;.

- Kobayashi K, Takahashi N, Jimi E, Udagawa N, Takami M, Kotake S, Nakagawa N, Kinoshita M, Yamaguchi K, Shima N, Yasuda H, Morinaga T, Higashio K, Martin TJ, Suda T. Tumor necrosis factor alpha stimulates osteoclast differentiation by a mechanism independent of the ODF/RANKL-RANK interaction. *J Exp Med* 2000;191:275-286.
- Koch AE, Harlow LA, Haines GK, Amento EP, Unemori EN, Wong WL, Pope RM, Ferrara N. Vascular endothelial growth factor. A cytokine modulating endothelial function in rheumatoid arthritis. *J Immunol* 1994;152:4149-4156.
- Kokkonen H, Soderstrom I, Rocklov J, Hallmans G, Lejon K, Rantapaa Dahlqvist S. Up-regulation of cytokines and chemokines predates the onset of rheumatoid arthritis. *Arthritis Rheum* 2010;62:383-391.
- Komine M, Kukita A, Kukita T, Ogata Y, Hotokebuchi T, Kohashi O. Tumor necrosis factor-alpha cooperates with receptor activator of nuclear factor kappaB ligand in generation of osteoclasts in stromal cell-depleted rat bone marrow cell culture. *Bone* 2001;28:474-483.
- Kong YY, Feige U, Sarosi I, Bolon B, Tafuri A, Morony S, Capparelli C, Li J, Elliott R, McCabe S, Wong T, Campagnuolo G, Moran E, Bogoch ER, Van G, Nguyen LT, Ohashi PS, Lacey DL, Fish E, Boyle WJ, Penninger JM. Activated T cells regulate bone loss and joint destruction in adjuvant arthritis through osteoprotegerin ligand. *Nature* 1999;402:304-309.
- Kopp S, Rockler B. Relationship between clinical and radiographic findings in patients with mandibular pain or dysfunction. *Acta Radiol Diagn (Stockh)* 1979;20:465-477.
- Kopp S. Neuroendocrine, immune, and local responses related to temporomandibular disorders. *J Orofac Pain* 2001;15:9-28.
- Kopp S, Alstergren P. Blood serotonin and joint pain in seropositive versus seronegative rheumatoid arthritis. *Mediators Inflamm* 2002;11:211-217.
- Kotake S, Udagawa N, Hakoda M, Mogi M, Yano K, Tsuda E, Takahashi K, Furuya T, Ishiyama S, Kim KJ, Saito S, Nishikawa T, Takahashi N, Togari A, Tomatsu T, Suda T, Kamatani N. Activated human T cells directly induce osteoclastogenesis from human monocytes: Possible role of T cells in bone destruction in rheumatoid arthritis patients. *Arthritis Rheum* 2001;44:1003-1012.
- Kraan MC, Versendaal H, Jonker M, Bresnihan B, Post WJ, Hart BA, Breedveld FC, Tak PP. Asymptomatic synovitis precedes clinically manifest arthritis. *Arthritis Rheum* 1998;41:1481-1488.
- Kwan Tat S, Padrines M, Theoleyre S, Heymann D, Fortun Y. IL-6, RANKL, TNF-alpha/IL-1: Interrelations in bone resorption pathophysiology. *Cytokine Growth Factor Rev* 2004;15:49-60.
- Kvien TK, Glennas A, Knudsen OG, Smedstad LM, Mowinckel P, Forre O. The prevalence and severity of rheumatoid arthritis in oslo. results from a county register and a population survey. *Scand J Rheumatol* 1997;26:412-418.

- Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, Elliott R, Colombero A, Elliott G, Scully S, Hsu H, Sullivan J, Hawkins N, Davy E, Capparelli C, Eli A, Qian YX, Kaufman S, Sarosi I, Shalhoub V, Senaldi G, Guo J, Delaney J, Boyle WJ. Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* 1998;93:165-176.
- Larheim TA, Kolbenstvedt A. Osseous temporomandibular joint abnormalities in rheumatic disease. computed tomography versus hypocyclusoidal tomography. *Acta Radiol* 1990;31:383-387.
- Larsen A, Dale K, Eek M. Radiographic evaluation of rheumatoid arthritis and related conditions by standard reference films. *Acta Radiol Diagn (Stockh)* 1977;18:481-491.
- Lawand NB, Willis WD, Westlund KN. Excitatory amino acid receptor involvement in peripheral nociceptive transmission in rats. *Eur J Pharmacol* 1997;324:169-177.
- Lawand NB, McNearney T, Westlund KN. Amino acid release into the knee joint: Key role in nociception and inflammation. *Pain* 2000;86:69-74.
- Lee SS, Joo YS, Kim WU, Min DJ, Min JK, Park SH, Cho CS, Kim HY. Vascular endothelial growth factor levels in the serum and synovial fluid of patients with rheumatoid arthritis. *Clin Exp Rheumatol* 2001;19:321-324.
- Lin TH, Yang RS, Tang CH, Wu MY, Fu WM. Regulation of the maturation of osteoblasts and osteoclastogenesis by glutamate. *Eur J Pharmacol* 2008;589:37-44.
- Lin YC, Hsu ML, Yang JS, Liang TH, Chou SL, Lin HY. Temporomandibular joint disorders in patients with rheumatoid arthritis. *J Chin Med Assoc* 2007;70:527-534.
- Loppnow H, Bil R, Hirt S, Schonbeck U, Herzberg M, Werdan K, Rietschel ET, Brandt E, Flad HD. Platelet-derived interleukin-1 induces cytokine production, but not proliferation of human vascular smooth muscle cells. *Blood* 1998;91:134-141.
- Ludlow JB, Davies-Ludlow LE, Brooks SL. Dosimetry of two extraoral direct digital imaging devices: NewTom cone beam CT and orthophos plus DS panoramic unit. *Dentomaxillofac Radiol* 2003;32:229-234.
- Lundstrom A, Lundstrom F, Lebrecht LM, Moorrees CF. Natural head position and natural head orientation: Basic considerations in cephalometric analysis and research. *Eur J Orthod* 1995;17:111-120.
- Machold KP, Nell V, Stamm T, Aletaha D, Smolen JS. Early rheumatoid arthritis. *Curr Opin Rheumatol* 2006;18:282-288.
- Machold KP, Stamm TA, Nell VP, Pflugbeil S, Aletaha D, Steiner G, Uffmann M, Smolen JS. Very recent onset rheumatoid arthritis: Clinical and serological patient characteristics associated with radiographic progression over the first years of disease. *Rheumatology (Oxford)* 2007;46:342-349.
- MacNaul KL, Hutchinson NI, Parsons JN, Bayne EK, Tocci MJ. Analysis of IL-1 and TNF-alpha gene expression in human rheumatoid synoviocytes and normal monocytes by in situ hybridization. *J Immunol* 1990;145:4154-4166.

- Mah JK, Danforth RA, Bumann A, Hatcher D. Radiation absorbed in maxillofacial imaging with a new dental computed tomography device. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2003;96:508-513.
- Masi AT, Bijlsma JW, Chikanza IC, Pitzalis C, Cutolo M. Neuroendocrine, immunologic, and microvascular systems interactions in rheumatoid arthritis: Physiopathogenetic and therapeutic perspectives. *Semin Arthritis Rheum* 1999;29:65-81.
- Mason DJ. Glutamate signalling and its potential application to tissue engineering of bone. *Eur Cell Mater* 2004;7:12-25; discussion 25-6.
- McNearney T, Speegle D, Lawand N, Lisse J, Westlund KN. Excitatory amino acid profiles of synovial fluid from patients with arthritis. *J Rheumatol* 2000;27:739-745.
- McNearney T, Baethge BA, Cao S, Alam R, Lisse JR, Westlund KN. Excitatory amino acids, TNF-alpha, and chemokine levels in synovial fluids of patients with active arthropathies. *Clin Exp Immunol* 2004;137:621-627.
- McNicol A, Israels SJ. Platelet dense granules: Structure, function and implications for haemostasis. *Thromb Res* 1999;95:1-18.
- Mentaverri R, Kamel S, Wattel A, Prouillet C, Sevenet N, Petit JP, Tordjmann T, Brazier M. Regulation of bone resorption and osteoclast survival by nitric oxide: Possible involvement of NMDA-receptor. *J Cell Biochem* 2003;88:1145-1156.
- Michael H, Harkonen PL, Vaananen HK, Hentunen TA. Estrogen and testosterone use different cellular pathways to inhibit osteoclastogenesis and bone resorption. *J Bone Miner Res* 2005;20:2224-2232.
- Miotla J, Maciewicz R, Kendrew J, Feldmann M, Paleolog E. Treatment with soluble VEGF receptor reduces disease severity in murine collagen-induced arthritis. *Lab Invest* 2000;80:1195-1205.
- Modder UI, Achenbach SJ, Amin S, Riggs BL, Melton LJ, 3rd, Khosla S. Relation of serum serotonin levels to bone density and structural parameters in women. *J Bone Miner Res* 2010;25:415-422.
- Mozzo P, Procacci C, Tacconi A, Martini PT, Andreis IA. A new volumetric CT machine for dental imaging based on the cone-beam technique: Preliminary results. *Eur Radiol* 1998;8:1558-1564.
- Nagashima M, Yoshino S, Ishiwata T, Asano G. Role of vascular endothelial growth factor in angiogenesis of rheumatoid arthritis. *J Rheumatol* 1995;22:1624-1630.
- Nakagawa M, Kaneda T, Arakawa T, Morita S, Sato T, Yomada T, Hanada K, Kumegawa M, Hakeda Y. Vascular endothelial growth factor (VEGF) directly enhances osteoclastic bone resorption and survival of mature osteoclasts. *FEBS Lett* 2000;473:161-164.
- Nakashima T, Kobayashi Y, Yamasaki S, Kawakami A, Eguchi K, Sasaki H, Sakai H. Protein expression and functional difference of membrane-bound and soluble receptor activator of NF-kappaB ligand: Modulation of the expression by osteotropic factors and cytokines. *Biochem Biophys Res Commun* 2000;275:768-775.

- Nell VP, Machold KP, Eberl G, Stamm TA, Uffmann M, Smolen JS. Benefit of very early referral and very early therapy with disease-modifying anti-rheumatic drugs in patients with early rheumatoid arthritis. *Rheumatology (Oxford)* 2004;43:906-914.
- Nordahl S, Alstergren P, Eliasson S, Kopp S. Interleukin-1beta in plasma and synovial fluid in relation to radiographic changes in arthritic temporomandibular joints. *Eur J Oral Sci* 1998;106:559-563.
- Nordahl S, Alstergren P, Kopp S. Tumor necrosis factor-alpha in synovial fluid and plasma from patients with chronic connective tissue disease and its relation to temporomandibular joint pain. *Journal of Oral and Maxillofacial Surgery* 2000;58:525-530.
- Nordahl S, Alstergren P, Eliasson S, Kopp S. Radiographic signs of bone destruction in the arthritic temporomandibular joint with special reference to markers of disease activity. A longitudinal study. *Rheumatology (Oxford)* 2001;40:691-694.
- Omote K, Kawamata T, Kawamata M, Namiki A. Formalin-induced release of excitatory amino acids in the skin of the rat hindpaw. *Brain Res* 1998;787:161-164.
- Paleolog EM, Miotla JM. Angiogenesis in arthritis: Role in disease pathogenesis and as a potential therapeutic target. *Angiogenesis* 1998;2:295-307.
- Paleolog EM. Angiogenesis in rheumatoid arthritis. *Arthritis Res* 2002;4 Suppl 3:S81-90.
- Pap T, Aupperle KR, Gay S, Firestein GS, Gay RE. Invasiveness of synovial fibroblasts is regulated by p53 in the SCID mouse in vivo model of cartilage invasion. *Arthritis Rheum* 2001;44:676-681.
- Pap T, Distler O. Linking angiogenesis to bone destruction in arthritis. *Arthritis Rheum* 2005;52:1346-1348.
- Peet NM, Grabowski PS, Laketic-Ljubojevic I, Skerry TM. The glutamate receptor antagonist MK801 modulates bone resorption in vitro by a mechanism predominantly involving osteoclast differentiation. *FASEB J* 1999;13:2179-2185.
- Perry D, Stewart N, Benton N, Robinson E, Yeoman S, Crabbe J, McQueen F. Detection of erosions in the rheumatoid hand; a comparative study of multidetector computerized tomography versus magnetic resonance scanning. *J Rheumatol* 2005;32:256-267.
- Pierce PA, Xie GX, Peroutka SJ, Green PG, Levine JD. 5-hydroxytryptamine-induced synovial plasma extravasation is mediated via 5-hydroxytryptamine_{2A} receptors on sympathetic efferent terminals. *J Pharmacol Exp Ther* 1995;275:502-508.
- Plant MJ, Williams AL, O'Sullivan MM, Lewis PA, Coles EC, Jessop JD. Relationship between time-integrated C-reactive protein levels and radiologic progression in patients with rheumatoid arthritis. *Arthritis Rheum* 2000;43:1473-1477.

- Poli V, Balena R, Fattori E, Markatos A, Yamamoto M, Tanaka H, Ciliberto G, Rodan GA, Costantini F. Interleukin-6 deficient mice are protected from bone loss caused by estrogen depletion. *EMBO J* 1994;13:1189-1196.
- Quinn MA, Emery P. Window of opportunity in early rheumatoid arthritis: Possibility of altering the disease process with early intervention. *Clin Exp Rheumatol* 2003;21:S154-7.
- Rantapaa-Dahlqvist S, de Jong BA, Berglin E, Hallmans G, Wadell G, Stenlund H, Sundin U, van Venrooij WJ. Antibodies against cyclic citrullinated peptide and IgA rheumatoid factor predict the development of rheumatoid arthritis. *Arthritis Rheum* 2003;48:2741-2749.
- Raza K, Falciani F, Curnow SJ, Ross EJ, Lee CY, Akbar AN, Lord JM, Gordon C, Buckley CD, Salmon M. Early rheumatoid arthritis is characterized by a distinct and transient synovial fluid cytokine profile of T cell and stromal cell origin. *Arthritis Res Ther* 2005;7:R784-95.
- Rengel Y, Ospelt C, Gay S. Proteinases in the joint: Clinical relevance of proteinases in joint destruction. *Arthritis Res Ther* 2007;9:221.
- Reynolds IJ, Miller RJ. 3H]MK801 binding to the NMDA receptor/ionophore complex is regulated by divalent cations: Evidence for multiple regulatory sites. *Eur J Pharmacol* 1988;151:103-112.
- Riggs BL. Endocrine causes of age-related bone loss and osteoporosis. *Novartis Found Symp* 2002;242:247-59; discussion 260-4.
- Riggs BL, Khosla S, Melton LJ, 3rd. Sex steroids and the construction and conservation of the adult skeleton. *Endocr Rev* 2002;23:279-302.
- Saika M, Inoue D, Kido S, Matsumoto T. 17beta-estradiol stimulates expression of osteoprotegerin by a mouse stromal cell line, ST-2, via estrogen receptor-alpha. *Endocrinology* 2001;142:2205-2212.
- Sambrook PN. The skeleton in rheumatoid arthritis: Common mechanisms for bone erosion and osteoporosis? *J Rheumatol* 2000;27:2541-2542.
- Saxne T, Palladino MA, Jr, Heinegard D, Talal N, Wollheim FA. Detection of tumor necrosis factor alpha but not tumor necrosis factor beta in rheumatoid arthritis synovial fluid and serum. *Arthritis Rheum* 1988;31:1041-1045.
- Scarfe WC, Farman AG. What is cone-beam CT and how does it work? *Dent Clin North Am* 2008;52:707-30, v.
- Schett G. Erosive arthritis. *Arthritis Res Ther* 2007;9 Suppl 1:S2.
- Schett G. Early rheumatoid arthritis. *Internist (Berl)* 2008;49:263-270.
- Schett G, Sieper J. Inflammation and repair mechanisms. *Clin Exp Rheumatol* 2009;27:S33-5.
- Scott DL, Pugnier K, Kaarela K, Doyle DV, Woolf A, Holmes J, Hieke K. The links between joint damage and disability in rheumatoid arthritis. *Rheumatology (Oxford)* 2000;39:122-132.
- Seide MF, Ulrich-Merzenich G, Fiebich B, Candelario-Jalil E, Koch FW, Vetter H. Tropisetron inhibits serotonin-induced PGE2 release from macrophage-like synovial cells in serum-free tissue culture. *Scand J Rheumatol Suppl* 2004;(119):33.

- Shevde NK, Bendixen AC, Dienger KM, Pike JW. Estrogens suppress RANK ligand-induced osteoclast differentiation via a stromal cell independent mechanism involving c-jun repression. *Proc Natl Acad Sci U S A* 2000;97:7829-7834.
- Shweiki D, Itin A, Soffer D, Keshet E. Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis. *Nature* 1992;359:843-845.
- Simonet WS, Lacey DL, Dunstan CR, Kelley M, Chang MS, Luthy R, Nguyen HQ, Wooden S, Bennett L, Boone T, Shimamoto G, DeRose M, Elliott R, Colombero A, Tan HL, Trail G, Sullivan J, Davy E, Bucay N, Renshaw-Gegg L, Hughes TM, Hill D, Pattison W, Campbell P, Sander S, Van G, Tarpley J, Derby P, Lee R, Boyle WJ. Osteoprotegerin: A novel secreted protein involved in the regulation of bone density. *Cell* 1997;89:309-319.
- Skerry TM. Neurotransmitter functions in bone remodeling. *J Musculoskeletal Neuronal Interact* 2002;2:281.
- Sokka T. Work disability in early rheumatoid arthritis. *Clin Exp Rheumatol* 2003;21:S71-4.
- Sone H, Sakauchi M, Takahashi A, Suzuki H, Inoue N, Iida K, Shimano H, Toyoshima H, Kawakami Y, Okuda Y, Matsuo K, Yamada N. Elevated levels of vascular endothelial growth factor in the sera of patients with rheumatoid arthritis correlation with disease activity. *Life Sci* 2001;69:1861-1869.
- Spencer GJ, Genever PG. Long-term potentiation in bone--a role for glutamate in strain-induced cellular memory? *BMC Cell Biol* 2003;4:9.
- Spencer GJ, Utting JC, Etheridge SL, Arnett TR, Genever PG. Wnt signalling in osteoblasts regulates expression of the receptor activator of NFkappaB ligand and inhibits osteoclastogenesis in vitro. *J Cell Sci* 2006;119:1283-1296.
- Spencer GJ, McGrath CJ, Genever PG. Current perspectives on NMDA-type glutamate signalling in bone. *Int J Biochem Cell Biol* 2007;39:1089-1104.
- Staud R, Robinson ME, Vierck CJ, Price DD. Diffuse noxious inhibitory controls (DNIC) attenuate temporal summation of second pain in normal males but not in normal females or fibromyalgia patients. *Pain* 2003;101:167-174.
- Straub RH. The complex role of estrogens in inflammation. *Endocr Rev* 2007;28:521-574.
- Suda T, Kobayashi K, Jimi E, Udagawa N, Takahashi N. The molecular basis of osteoclast differentiation and activation. *Novartis Found Symp* 2001;232:235-47; discussion 247-50.
- Symmons DP. Epidemiology of rheumatoid arthritis: Determinants of onset, persistence and outcome. *Best Pract Res Clin Rheumatol* 2002;16:707-722.
- Syrjanen SM. The temporomandibular joint in rheumatoid arthritis. *Acta Radiol Diagn (Stockh)* 1985;26:235-243.
- Syversen SW, Gaarder PI, Goll GL, Odegard S, Haavardsholm EA, Mowinckel P, van der Heijde D, Landewe R, Kvien TK. High anti-cyclic citrullinated peptide levels and an algorithm of four variables predict radiographic progression in

- patients with rheumatoid arthritis: Results from a 10-year longitudinal study. *Ann Rheum Dis* 2008;67:212-217.
- Takahashi N, Yamana H, Yoshiki S, Roodman GD, Mundy GR, Jones SJ, Boyde A, Suda T. Osteoclast-like cell formation and its regulation by osteotropic hormones in mouse bone marrow cultures. *Endocrinology* 1988;122:1373-1382.
- Takayanagi H, Iizuka H, Juji T, Nakagawa T, Yamamoto A, Miyazaki T, Koshihara Y, Oda H, Nakamura K, Tanaka S. Involvement of receptor activator of nuclear factor kappaB ligand/osteoclast differentiation factor in osteoclastogenesis from synoviocytes in rheumatoid arthritis. *Arthritis Rheum* 2000;43:259-269.
- Tanino M, Matoba R, Nakamura S, Kameda H, Amano K, Okayama T, Nagasawa H, Suzuki K, Matsubara K, Takeuchi T. Prediction of efficacy of anti-TNF biologic agent, infliximab, for rheumatoid arthritis patients using a comprehensive transcriptome analysis of white blood cells. *Biochem Biophys Res Commun* 2009;387:261-265.
- Tegelberg A, Kopp S. Clinical findings in the stomatognathic system for individuals with rheumatoid arthritis and osteoarthritis. *Acta Odontol Scand* 1987;45:65-75.
- Tengstrand B, Carlstrom K, Fellander-Tsai L, Hafstrom I. Abnormal levels of serum dehydroepiandrosterone, estrone, and estradiol in men with rheumatoid arthritis: High correlation between serum estradiol and current degree of inflammation. *J Rheumatol* 2003;30:2338-2343.
- Tengstrand B, Carlstrom K, Hafstrom I. Gonadal hormones in men with rheumatoid arthritis--from onset through 2 years. *J Rheumatol* 2009;36:887-892.
- Thabet MM, Huizinga TW, van der Heijde DM, van der Helm-van Mil AH. The prognostic value of baseline erosions in undifferentiated arthritis. *Arthritis Res Ther* 2009;11:R155.
- Trang LE, Furst P, Odeback AC, Lovgren O. Plasma amino acids in rheumatoid arthritis. *Scand J Rheumatol* 1985;14:393-402.
- Tremolizzo L, DiFrancesco JC, Rodriguez-Menendez V, Sirtori E, Longoni M, Cassetti A, Bossi M, El Mestikawy S, Cavaletti G, Ferrarese C. Human platelets express the synaptic markers VGLUT1 and 2 and release glutamate following aggregation. *Neurosci Lett* 2006;404:262-265.
- Tunyogi-Csapo M, Kis-Toth K, Radacs M, Farkas B, Jacobs JJ, Finnegan A, Mikecz K, Glant TT. Cytokine-controlled RANKL and osteoprotegerin expression by human and mouse synovial fibroblasts: Fibroblast-mediated pathologic bone resorption. *Arthritis Rheum* 2008;58:2397-2408.
- Turner CH, Robling AG, Duncan RL, Burr DB. Do bone cells behave like a neuronal network? *Calcif Tissue Int* 2002;70:435-442.
- Uhlig T, Smedstad LM, Vaglum P, Moum T, Gerard N, Kvien TK. The course of rheumatoid arthritis and predictors of psychological, physical and radiographic outcome after 5 years of follow-up. *Rheumatology (Oxford)* 2000;39:732-741.

- Walsh NC, Gravalles EM. Bone loss in inflammatory arthritis: Mechanisms and treatment strategies. *Curr Opin Rheumatol* 2004;16:419-427.
- van Aken J, Lard LR, le Cessie S, Hazes JM, Breedveld FC, Huizinga TW. Radiological outcome after four years of early versus delayed treatment strategy in patients with recent onset rheumatoid arthritis. *Ann Rheum Dis* 2004;63:274-279.
- van den Berg WB, Bresnihan B. Pathogenesis of joint damage in rheumatoid arthritis: Evidence of a dominant role for interleukin-I. *Baillieres Best Pract Res Clin Rheumatol* 1999;13:577-597.
- van der Heijde D, Dankert T, Nieman F, Rau R, Boers M. Reliability and sensitivity to change of a simplification of the Sharp/van der heijde radiological assessment in rheumatoid arthritis. *Rheumatology (Oxford)* 1999;38:941-947.
- van der Heijde D, Landewe R, Boonen A, Einstein S, Herborn G, Rau R, Wassenberg S, Weissman BN, Winalski CS, Sharp JT. Expert agreement confirms that negative changes in hand and foot radiographs are a surrogate for repair in patients with rheumatoid arthritis. *Arthritis Res Ther* 2007;9:R62.
- van der Heijde DM, van Riel PL, van Leeuwen MA, van 't Hof MA, van Rijswijk MH, van de Putte LB. Prognostic factors for radiographic damage and physical disability in early rheumatoid arthritis. A prospective follow-up study of 147 patients. *Br J Rheumatol* 1992;31:519-525.
- van der Heijde DM. Joint erosions and patients with early rheumatoid arthritis. *Br J Rheumatol* 1995;34 Suppl 2:74-78.
- van Leeuwen MA, van Rijswijk MH, van der Heijde DM, Te Meerman GJ, van Riel PL, Houtman PM, van De Putte LB, Limburg PC. The acute-phase response in relation to radiographic progression in early rheumatoid arthritis: A prospective study during the first three years of the disease. *Br J Rheumatol* 1993;32 Suppl 3:9-13.
- van Leeuwen MA, Westra J, van Riel PL, Limburg PC, van Rijswijk MH. IgM, IgA, and IgG rheumatoid factors in early rheumatoid arthritis predictive of radiological progression? *Scand J Rheumatol* 1995;24:146-153.
- van Vollenhoven RF. In early RA with insufficient response to MTX, the addition of anti-TNF results in less radiological progression over 24 months than the addition of conventional DMARDs: results from the SWEFOT trial. Abstract American College of Rheumatology Annual Meeting, October 2009 in Philadelphia, PN, USA.
- van Vollenhoven RF, Ernestam S, Geborek P, Petersson IF, Coster L, Waltbrand E, Zickert A, Theander J, Thorner A, Hellstrom H, Teleman A, Dackhammar C, Akre F, Forslind K, Ljung L, Oding R, Chatzidionysiou A, Wornert M, Bratt J. Addition of infliximab compared with addition of sulfasalazine and hydroxychloroquine to methotrexate in patients with early rheumatoid arthritis (swefot trial): 1-year results of a randomised trial. *Lancet* 2009;374:459-466.
- Vanderschueren D, Vandenput L, Boonen S, Lindberg MK, Bouillon R, Ohlsson C. Androgens and bone. *Endocr Rev* 2004;25:389-425.

- Warden SJ, Blizotes MM, Wiren KM, Eshleman AJ, Turner CH. Neural regulation of bone and the skeletal effects of serotonin (5-hydroxytryptamine). *Mol Cell Endocrinol* 2005;242:1-9.
- Westbroek I, van der Plas A, de Rooij KE, Klein-Nulend J, Nijweide PJ. Expression of serotonin receptors in bone. *J Biol Chem* 2001;276:28961-28968.
- Wiese M, Wenzel A, Hintze H, Petersson A, Knutsson K, Bakke M, List T, Svensson P. Osseous changes and condyle position in TMJ tomograms: Impact of RDC/TMD clinical diagnoses on agreement between expected and actual findings. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;106:e52-63.
- Vittecoq O, Pouplin S, Krzanowska K, Jouen-Beades F, Menard JF, Gayet A, Daragon A, Tron F, Le Loet X. Rheumatoid factor is the strongest predictor of radiological progression of rheumatoid arthritis in a three-year prospective study in community-recruited patients. *Rheumatology (Oxford)* 2003;42:939-946.
- Wolfe F. Comparative usefulness of C-reactive protein and erythrocyte sedimentation rate in patients with rheumatoid arthritis. *J Rheumatol* 1997;24:1477-1485.
- Voog U, Alstergren P, Eliasson S, Leibur E, Kallikorm R, Kopp S. Inflammatory mediators and radiographic changes in temporomandibular joints of patients with rheumatoid arthritis. *Acta Odontol Scand* 2003a;61:57-64.
- Voog U, Alstergren P, Leibur E, Kallikorm R, Kopp S. Impact of temporomandibular joint pain on activities of daily living in patients with rheumatoid arthritis. *Acta Odontol Scand* 2003b;61:278-282.
- Voog U, Alstergren P, Eliasson S, Leibur E, Kallikorm R, Kopp S. Progression of radiographic changes in the temporomandibular joints of patients with rheumatoid arthritis in relation to inflammatory markers and mediators in the blood. *Acta Odontol Scand* 2004;62:7-13.
- Yadav VK, Ryu JH, Suda N, Tanaka KF, Gingrich JA, Schutz G, Glorieux FH, Chiang CY, Zajac JD, Insogna KL, Mann JJ, Hen R, Ducy P, Karsenty G. Lrp5 controls bone formation by inhibiting serotonin synthesis in the duodenum. *Cell* 2008;135:825-837.
- Yelin E, Lubeck D, Holman H, Epstein W. The impact of rheumatoid arthritis and osteoarthritis: The activities of patients with rheumatoid arthritis and osteoarthritis compared to controls. *J Rheumatol* 1987;14:710-717.
- Yoneda T, Ishimaru N, Arakaki R, Kobayashi M, Izawa T, Moriyama K, Hayashi Y. Estrogen deficiency accelerates murine autoimmune arthritis associated with receptor activator of nuclear factor-kappa B ligand-mediated osteoclastogenesis. *Endocrinology* 2004;145:2384-2391.