



Karolinska
Institutet

Karolinska Institutet

<http://openarchive.ki.se>

This is a Peer Reviewed Accepted version of the following article, accepted for publication in *Clinical & Experimental Allergy*.

2016-04-26

Impact of asthma medication and familial factors on the association between childhood asthma and attention-deficit/hyperactivity disorder : a combined twin- and register-based study

Holmberg, Kirsten; Lundholm, Cecilia; Anckarsäter, Henrik; Larsson, Henrik; Almqvist, Catarina

Clin Exp Allergy. 2015 May;45(5):964-973.

<http://doi.org/10.1111/cea.12529>

<http://hdl.handle.net/10616/45139>

If not otherwise stated by the Publisher's Terms and conditions, the manuscript is deposited under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

Impact of asthma medication and familial factors on the association between childhood asthma and ADHD

A combined twin- and register-based study

Kirsten Holmberg, MD, PhD^{1,2}, Cecilia Lundholm, MSc¹, Henrik Anckarsäter, MD, PhD³, Henrik Larsson, PhD¹, and Catarina Almqvist, MD, PhD^{1,4}

¹Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden.

²Neuropediatric unit, Sachs' Children and Youth Hospital, South General Hospital, Stockholm, Sweden.

³Institute of Neuroscience and Physiology, Forensic Psychiatry, University of Gothenburg, Sweden.

⁴Lung and allergy unit, Astrid Lindgren Children's Hospital, Karolinska University Hospital, Stockholm, Sweden.

Corresponding author: Kirsten Holmberg, Department of Medical Epidemiology and Biostatistics; PO Box 281, Karolinska Institutet, SE-171 77 Stockholm, SWEDEN
T +46 (0)8 616 4071, F +46 (0)8 31 49 75
e-mail: kirsten.holmberg@ki.se

Running head: The association between childhood asthma and ADHD

Abstract

Background

Asthma and attention-deficit/hyperactivity disorder (ADHD) are prevalent in childhood and may cause functional impairment and stress in families. Previous research supports an association between asthma and ADHD in children, but several aspects of this relationship are unclear.

Objective

Our aim was to study if the association between asthma and ADHD is restricted to either the inattentive or the hyperactive/impulsive symptoms of ADHD, to explore the impact of asthma severity and asthma medication and the contribution of shared genetic and environmental risk factors on the asthma-ADHD relationship.

Methods

Data on asthma, ADHD, zygosity, and possible confounders were collected from parental questionnaires at 9 or 12 years on 20 072 twins through the Swedish Twin Register, linked to the Swedish Medical Birth Register, the National Patient Register and the Prescribed Drug Register. The association between asthma and ADHD, the impact of asthma severity and medication, was assessed by generalized estimating equations. Cross-twin-cross-trait correlations (CTCT) were estimated to explore the relative importance of genes and environment for the association.

Results

Asthmatic children had a higher risk of also having ADHD (odds ratio [OR] 1.53, 95% confidence interval [CI]: 1.16–2.02). The association was not restricted to either of the two dimensions of ADHD. The magnitude of the association increased with asthma severity (OR 2.84, 95% CI: 1.86–4.35) for ≥ 4 asthma attacks in the last 12 months and was not affected by

asthma treatment. The CTCTs possibly indicate that the genetic component in overlap of the disorders is weak.

Conclusions and Clinical Relevance

Childhood asthma, especially severe asthma, is associated with ADHD. Asthma medication seems not to increase the risk of ADHD. Clinicians should be aware of the potential of ADHD in asthma. Optimal asthma care needs to be integrated with effective evaluation and treatment of ADHD in children with co-existing disorders.

Introduction

Asthma and attention-deficit/hyperactivity disorder (ADHD) [1] are prevalent chronic health conditions in childhood which may cause functional impairment and stress in families [2, 3]. There is evidence from some cross-sectional studies to suggest an association between asthma and ADHD [4-7], while other studies did not find a relationship [8, 9]. Several aspects of this relationship are still poorly understood [8, 10].

There are several potential explanations to the observed overlap between asthma and ADHD. Asthma may contribute to the subsequent development of ADHD, and a dose-response relationship between asthma severity and rates of behavioural problems has been suggested [11]. In a longitudinal study of 1480 Swedish twin pairs our group recently found that childhood asthma was associated with a linear increase in risk of hyperactivity–impulsivity behaviours later in adolescence [12]. Asthma severity and medication are however tightly linked, and it is possible that medication rather than asthma is the explanation for the association. Treatment with β_2 -agonists and inhaled corticosteroids (ICS) have been reported to cause behavioural side effects [13, 14]. However, we did not find any significant difference in hyperactivity–impulsivity between asthmatic children with or without asthma medication [12].

Alternatively, asthma and ADHD may be indirectly linked via shared genetic and/or environmental risk factors. Several environmental risk factors such as low parental socioeconomic status and parental smoking have been associated with both asthma and ADHD [15-18]. Although it has been proposed that there is a common environmental mechanism underlying the development of asthma and/or ADHD [19], the association seems to remain in other studies after adjustment for these potential confounders [11, 12].

Twin studies have demonstrated that both asthma and ADHD are highly heritable [20, 21]. However, the few previous family and twin studies have produced conflicting results regarding the genetic and shared environmental contribution to the overlap between asthma and ADHD. Two previous family studies did not find any genetic and/or shared environmental component underlying the association [8, 9], whereas our recent twin study reported that the overlap between asthma and hyperactivity–impulsivity was largely due to genetic influences [12]. Clearly, additional research in larger populations is needed to clarify the impact of environmental and genetic factors on the link between asthma–ADHD.

The aim of this study was to i) confirm the association of asthma and ADHD symptoms in childhood and to explore if the association was restricted to the inattentive or the hyperactive/impulsive symptoms of ADHD, ii) investigate the impact of asthma severity and asthma medication (inhaled β_2 -agonists and/or inhaled corticosteroids) on the asthma-ADHD relationship and iii) examine the contribution of shared genetic and environmental risk factors to the overlap between the two conditions. We used a large cross-sectional twin cohort based on the population-based Swedish Twin Register with data on asthma and ADHD-symptoms as well as asthma medication and specialist care diagnoses from the Swedish National Health Registers.

Methods

Study design and population

This population-based cross-sectional study used data from the Child and Adolescent Twin Study in Sweden (CATSS) [22] and three Swedish national registers held by the Swedish National Board of Health and Welfare. CATSS collects information on 9-year-old (born since July 1995) or 12-year-old (born July 1992–June 1995) twins through parental interviews [22]. The population used in this study included all children born between 1992 and 2002 whose parents were contacted for interviews between Aug 2003 and Aug 2011 (n=27 260). Twins were excluded if they were living abroad, had a parent not fluent in Swedish, or they or their parent were disabled (n=860). In total, 26 400 twins were eligible and 20 302 (75%) completed the interview. The final study population (n=20 072) was restricted to twins with complete answers to the main questions on asthma and ADHD.

Data from CATSS was linked to the Medical Birth Register (MBR), National Patient Register (NPR) and Prescribed Drug Register (PDR) using the Personal Identification Number (PIN), a unique identifier for each resident. Further details regarding data collection and linkage to these national registers are described elsewhere [17, 18, 23-25]. The MBR contains maternal and child characteristics data on more than 99% of all births in Sweden since 1973. The NPR covers all inpatient diagnoses since 1987 and about 80% of all outpatient diagnoses since 2001, recorded with *International Classification of Disease (ICD)-9* and ICD-10 codes. The PDR encompasses data on all prescribed drugs dispensed in Swedish pharmacies since July 1, 2005 [26], coded by the Anatomical Therapeutic Chemical (ATC) system. Register data was available for twins born 1992–2001 (94% of the study population). Data from NPR and PDR was collected until Dec 31, 2009.

Measures

Asthma

The main definition of asthma, “asthma reported by questionnaire”, and a secondary variable “doctor's asthma diagnosis reported by questionnaire”, were derived from the International Study of Asthma and Allergies in Childhood (ISAAC) questions in CATSS [18, 27]. Asthma cases were also identified from the NPR “asthma diagnosis based in register”, if a record in the NPR with asthma as the primary or a secondary diagnosis (ICD-9 code 493 or ICD-10 codes J45 and J46) was recorded at least once. The quality of doctor-diagnosed asthma in the NPR is considered high [24]. Finally, a fourth asthma variable “any asthma” was derived from “asthma reported by questionnaire” and “asthma diagnosis based in register”.

Asthma severity

Parents were also asked ISAAC questions about wheezing. If they reported either asthma or wheezing, they were asked follow-up questions on number of attacks in the last 12 months [27]. Answers were categorised into five levels: No; Yes, previous asthma only; Yes, current asthma but no attacks in the last 12 months; 1-3 attacks in the last 12 months, and ≥ 4 attacks in the last 12 months.

Asthma medication

Dispensed asthma medications β_2 -agonists (R03AC), inhaled corticosteroids (ICS) including those combined with long-acting β_2 -agonists (R03BA, R03AK) and leukotriene receptor antagonists (R03DC) were retrieved from the PDR up until the date of interview, for the subset of individuals with any asthma and at least 18 months between start of the PDR and the interview (i.e. individuals with interviews later than Dec 31, 2006). In our analyses, asthma

medication was categorised in three ways: (1) dispensed any asthma drugs or (2) dispensed inhaled β_2 -agonists and/or ICS at least once or (3) dispensed inhaled β_2 -agonists and/or ICS at least three times.

ADHD

The CATSS interviews included an ADHD scale as part of the Autism – Tics, ADHD, and other Comorbidities inventory (A-TAC). This screening instrument asks symptom questions based on the DSM-IV criteria including two modules: Attention and Hyperactivity. Response options for each question are coded as “0 – no”, “0.5 – yes, to some extent” and “1 – yes”. The A-TAC ADHD scale or subscales can be used to identify possibly impairing ADHD symptoms. A score of 6 or more is used as a “screening” ADHD diagnosis, and a score of 12.5 or more selects children with ADHD at a level of clinical concern (“research” ADHD diagnosis) [28].

In this study, the ADHD outcome variable was defined as parental report of either a screening or research-level ADHD diagnosis. The two modules Attention and Hyperactivity (with cut-off score > 6 on each scale) in A-TAC were also analysed separately. Finally, we defined “register-based ADHD diagnosis” as either an ADHD diagnosis in NPR (Hyperkinetic syndrome, ICD-9 code 314; or Hyperkinetic disorders, ICD-10 code: F900 registered, and/or dispensed ADHD-drugs in PDR: centralstimulants (N06BA01, N06BA04) or atomoxetine (N06BA09) registered at least once.

Covariates

Atopic eczema was denoted if parents reported “atopic eczema” or “flexural eczema” according to the ISAAC question in CATSS [29] . Information on a child's sex and zygosity

[30] as well as data on the mother's birth country and level of education (<9 years, 10–12 years or graduate education) and parental cohabitation at time of interview (yes/no) was collected in the CATSS interview. Data on birth weight and gestational age (GA) and maternal smoking habits at the first antenatal care visit (0, 1–9 or >10 cigarettes per day) and age at delivery were retrieved from MBR.

Statistical analysis

The associations between asthma and ADHD were analysed using the different ADHD diagnoses (screening, research, or register-based) as dichotomous outcome variables, and the different asthma measures as independent variables, one by one. We used generalized estimating equations with the logit link and exchangeable covariance structure to address the correlation caused by the clustering of observations within twin pairs [31]. Odds ratios and 95% confidence intervals (CI) were first adjusted for the child's sex and age. Secondly, models were adjusted for the child's sex, age, birth weight, GA; as well as the maternal characteristics age at delivery, smoking, birth country, education, and parental cohabitation. Thirdly, models were adjusted for all the previous covariates and additionally for atopic eczema. Furthermore we tested sex differences in the associations between asthma and ADHD by including an interaction term, and if statistically significant we estimated separate ORs for males and females. The classical twin design was used to explore the relative impact of genetic, shared environmental and non-shared environmental factors for asthma, ADHD and the overlap between the disorders. Briefly, tetrachoric twin correlations for asthma and ADHD were estimated separately for monozygotic (MZ) and dizygotic (DZ) twin pairs. The corresponding cross-twin cross-trait correlations (i.e. correlation between asthma status in one

twin and ADHD status in the co-twin) were estimated to assess the relative importance of genetic and environmental influences for the overlap between asthma and ADHD.

All statistical analyses were carried out using the SAS software package version 9.2 and the significance level was 0.05.

Permission for the study was obtained from the Regional Ethical Review board in Stockholm, Sweden.

Results

The proportion of study subjects with parent reported asthma and screening ADHD diagnosis by child and maternal characteristics is presented in Supplementary Table S1. The overall cumulative incidence of asthma was 14.0%, and 9.9% of the study population had a screening diagnosis of ADHD. Two percent had both disorders at 9 or 12 years of age. The cumulative incidence of research ADHD diagnosis in the study population was 1.9% (375/20,072) and 2.0% (383/18,931) had a register-based diagnosis of ADHD. About 76% of the children with a register-based ADHD diagnosis also had a parent-reported screening ADHD diagnosis in the A-TAC interviews.

Table 1 presents the associations between asthma and the ADHD outcomes. After adjusting for sex, children with asthma were more likely to have ADHD than children with no asthma. Children with parent-reported asthma in questionnaires had an increased odds of a screening diagnosis of ADHD (fully adjusted OR 1.32, 95% CI: 1.14–1.52), a research diagnosis of ADHD (OR 1.86, 95% CI: 1.39–2.49) and a register-based ADHD diagnosis (OR 1.55, 95% CI: 1.16–2.07). The table displays similar findings for the association between doctors' diagnosis of asthma reported by questionnaires and in registers, or with any asthma, and the various ADHD outcomes. There were no sex differences in the association between asthma and ADHD, except that a research diagnosis of ADHD was more strongly associated with a doctor's diagnosis of asthma in females (p for interaction=0.044; OR 2.88, 95% CI: 1.69–4.89 in females; OR 1.48, 95% CI: 1.02–2.13 in males).

Asthma was associated equally with Attention and Hyperactivity. Using the Attention and Hyperactivity subscales of ADHD separately as outcome variables, we obtained similar adjusted ORs for the association with asthma: parental report of at least six inattentive symptoms: OR 1.56, 95% CI: 1.24–1.97 and parental report of at least six hyperactivity–

impulsivity symptoms: OR 1.59, 95% CI: 1.21–2.10. The association between asthma and ADHD did not change with adjustment for eczema (data not shown).

Table 2 displays the relation between the parent-reported frequency of asthma attacks in the last 12 months and the odds of ADHD. After adjusting for all potential confounders, the OR for the association with the screening ADHD diagnosis consistently increased with an increasing number of asthma attacks compared to no asthma (1-3 attacks: OR 1.40, 95% CI: 1.08–1.83 and ≥ 4 attacks: OR 1.55, 95% CI: 1.17–2.04). The association was also statistically significant for the “yes, previous asthma only” category of asthma attacks (OR 1.29, CI 95%: 1.04–1.60). Similar results were obtained for the association between asthma attacks in the last 12 months and research diagnosis of ADHD, but were only significant for ≥ 4 attacks (OR 2.84, 95% CI: 1.86–4.35) and “yes, previous asthma only” (OR 1.87, 95% CI: 1.23–2.84). Furthermore, there was a similar significant association between a register-based ADHD diagnosis in all categories of asthma attacks in the last 12 months except the 1-3 attacks category (data not shown).

Table 3 illustrates the association between different classes of dispensed asthma medications and the risk of ADHD. Both asthma groups, that is with and without dispensed asthma drugs, had increased odds of a screening or research diagnosis of ADHD, but there was no significant difference between the groups (screening diagnosis of ADHD: OR 1.36, 95% CI: 1.03–1.80 for asthma, but no dispensed drugs; OR 1.56, 95% CI: 1.21–2.01 for asthma with any dispensed asthma medication in the fully adjusted model). A higher number of dispensed drugs did not increase the odds of a screening diagnosis of ADHD: OR 1.46, 95% CI: 1.16–1.84 for no dispensed drugs; OR 1.55, 95% CI: 1.08–2.22 for dispensed both inhaled β_2 -agonists and ICS ≥ 3 times in the fully adjusted model). For a research diagnosis of ADHD,

Kirsten 2015-3-18 13:29

Deleted: i.e.

the OR for having dispensed both inhaled β_2 -agonists and ICS ≥ 3 times (OR 2.69, 95% CI: 1.57–4.62) was slightly higher than for children with asthma but without dispensed drugs (OR 2.06, 95% CI: 1.39–3.07).

Table 4 shows the results of the twin analyses. These include intra-class and cross-trait cross-twin correlations (CTCT; the correlation between one twin's status on asthma and the co-twin's status on a screening or research ADHD diagnosis), correlations for parent-reported asthma and the two ADHD diagnoses, by zygosity. MZ intra-class correlations were consistently higher than the corresponding DZ correlations [parent-reported asthma: MZ (OR 0.87, 95% CI: 0.85–0.90) and DZ same sex (OR 0.43, 95% CI: 0.36–0.49); research ADHD diagnosis: MZ (OR 0.85, 95% CI: 0.77–0.93) and DZ same sex (OR 0.37, 95% CI: 0.20–0.54)]. These results support genetic influences on asthma and on ADHD in both sexes. However, the CTCT correlations for MZ twins and DZ twins were similar in strength, or even higher among DZ twins [asthma and research ADHD diagnosis: MZ 0.05 (-0.08, 0.17) vs DZ same sex 0.13 (0.03, 0.23)], possibly suggesting that any genetic component in the overlap in the liability of the phenotypes is weak.

Discussion

In this cohort analysis of 20,072 Swedish twins aged 9 or 12 we found that children with asthma had an almost twofold increased risk of ADHD compared with those without asthma. The association was not restricted to the inattentive or the hyperactive/impulsive symptoms of ADHD. The magnitude of the association increased with asthma severity defined as frequency of asthma attacks, but was not affected by asthma treatment. The results were robust after adjusting for potential confounders including atopic eczema [32] and independent of whether asthma or ADHD was defined as parent-reported symptoms on a questionnaire or as a register-based diagnosis. To our knowledge, the effect of asthma severity and asthma drug treatment on associated ADHD symptoms in childhood has not been studied previously. The study design is novel, and combines parent reported symptoms with national registers in a large population of twins. Furthermore, this is the largest population-based twin- and register-based study on childhood asthma and ADHD to date, and supports the hypothesis that asthma is associated with ADHD. However, due to the cross-sectional study design, we could not determine the causal direction in the association between asthma and ADHD.

Previous large epidemiological studies have shown a cross-sectional association between asthma and ADHD [4, 6, 33, 34] in contrast to studies with small sample sizes ($N \sim 1,000$) [8, 9]. Recent analyses from nationwide population-based studies with diagnosis based on parental report [6], register-based diagnoses [34] or prescription data [33] reported similar ORs for the association between asthma and ADHD as we found in our study. In a previous study we distinguished between the two DSM-IV ADHD symptom dimensions and reported that childhood asthma predicts hyperactivity–impulsivity, but not inattention, in adolescence [12]. In the present study, we found that the ORs for the association between asthma and ADHD were similar regardless of defining ADHD symptoms as hyperactivity–impulsivity or

inattention, or the combination of those. This divergent result may depend on the difference in study designs or age. In the present cross-sectional study information on asthma and ADHD were collected at the same age (9 or 12 years). In contrast, the previous study was longitudinal between 7 to 14 years of age. Although recent studies have shown a relationship between atopic eczema and ADHD [32], our results did not change when adjusting for eczema which is in line with our previous study [12].

The prevalence of research ADHD diagnosis was rather low and in line with the prevalence of register-diagnosed ADHD, indicating that this group consisted of children with severe ADHD in need of health care. Another explanation is that parents of children with high A-TAC scores might tend to underreport or not participate in the study.

Current asthma severity was linearly associated with ADHD which confirms results from previous studies [10-12]. ADHD causes psychological distress and may exacerbate the severity of asthma [11, 35]. On the other hand, childhood asthma, especially severe disease, may be associated with family distress and behavioural problems and interpreted as ADHD or worsen functional impairment caused by ADHD [2, 16, 35]. The stress of treating severe asthma may also have psychological impact on children and family relationships. In addition, we found an increased prevalence of ADHD in children who only had reported a history of previous asthma. The explanation may be that asthma at a young age influences the risk of developing ADHD symptoms by school-age. This result is in line with our longitudinal study of asthma and ADHD, indicating that the correct causal pathway is that asthma precedes ADHD [12]. Although the present study is cross-sectional, there may be other explanations for the overlap. Gene expression resulting in immune system dysfunction may be a common pathophysiological mechanism underlying both disorders [36].

We found no evidence to support the notion that the association between asthma and ADHD was modified by asthma medication. Even children who had dispensed both inhaled β_2 -agonists and ICS > 3 times had similar odds of screening ADHD diagnosis as those without dispensed medication. This may be due to severe asthma in spite of intensive medication or severe cases being undertreated. Possibly the ADHD symptoms are being ignored, or misinterpreted as asthma or as side-effects caused by asthma medication. Another possibility is that the restlessness caused by β_2 -agonist may be a transient effect [37]. Our findings of no significant effects by medication on the association based on information from the PDR replicate the result from our previous longitudinal study of Swedish twins where use of inhaled β_2 -agonists and/or ICS was assessed from parental questionnaires [12].

The association between asthma and ADHD was stable when controlled for child and maternal characteristics, including sex, birth weight, maternal level of education and smoking habits. Previous research has not adjusted for all these covariates [11, 34, 38]. Results from the twin analyses confirm previous research suggesting a strong genetic component for asthma and ADHD in both sexes [11, 20, 21]. However, the estimated cross-twin cross-trait correlations were similar in MZ and DZ twins, or even higher in DZ twins. This contradicts the notion of a genetic component in the co-occurrence of asthma and ADHD. Consequently, we could not fit a model for estimating the size of such a genetic component.

There are several strengths of our study. Firstly, we studied a large population. Secondly, we obtained robust results independent of whether validated questionnaire or register-based variables on asthma and ADHD were used [24, 28], and we investigated the two subscale measures of ADHD – Attention or Hyperactivity – separately. Thirdly, we adjusted for several socioeconomic confounders which have been limited in previous studies [10, 34].

Fourthly, we were able to evaluate the severity of both asthma and ADHD symptoms in contrast to a recent study which was register-based only [34].

This study should be interpreted in the context of its limitations. Firstly, the study is cross-sectional, which constrains possibilities to study the direction of effect between asthma and ADHD. Secondly, we have no external validation of parent-reported hyperactivity or inattention, such as teacher reports. Teachers tend to report more ADHD symptoms both in children with or without asthma than parents do [2, 39]. However, there were no substantial differences between the results in the parent-reported ADHD groups of screening- or research-based diagnoses of ADHD and the group with the register-based clinical diagnosis of ADHD, which usually includes assessment of impairment in school [1]. Thirdly, asthma severity was based on number of asthma attacks in the last 12 months. We did not have additional ways to assess severity such as children's lung function or other clinical data. Fourthly, we have no access to detailed clinical information on treatment with asthma drugs, doses or duration, or treatment compliance. Finally, we did not control for comorbid psychiatric disorders, [38] or sleeping problems which have been reported to increase the risk of ADHD symptoms [40]. Generalizability of our findings in twins to the general population may be a concern, although previous studies indicate that twins do not differ from singletons concerning their risk of asthma [41] or ADHD [42]. Moreover, our estimated ORs for the association between asthma and ADHD are of similar magnitude as those found in studies in non-twin populations [6, 33, 34], suggesting a similar etiology in twins and singletons.

In conclusion, this large population-based cross-sectional study indicates that asthma and asthma severity but not asthma medication is associated with ADHD symptoms in childhood. These findings highlight the need for optimal clinical care of children with asthma to be

integrated with effective management strategies for ADHD symptoms [34]. Clinicians need to be aware of the potential for ADHD especially in children with severe asthma and optimize drug treatment also for children with both conditions. Our results indicate that it would be valuable to further elucidate the relationship between parental perception of child's behaviour and illness to establish services for early identification and intervention for behavioural difficulties in children with asthma – or asthma intervention for children with ADHD – to minimize the impact of physical illness and psychological disturbance [2, 10].

Acknowledgments

We are grateful to the children and parents of the CATSS twin cohort, without whose participation this study would have been impossible to perform.

Funding: Financial support was provided from the Swedish Research Council (grant number 2011-3060) and through the Swedish Initiative for research on Microdata in the Social And Medical sciences (SIMSAM) framework grant no 340-2013-5867, Swedish Asthma and Allergy Association's Research Foundation, the regional agreement on medical training and clinical research (ALF) between Stockholm County Council and Karolinska Institutet and the Strategic Research Program in Epidemiology at Karolinska Institutet.

SLUTVERSION enligt CA 150318 **Funding:** Financial support was provided from the Swedish Research Council (grant number 2011-3060) and through the Swedish Initiative for research on Microdata in the Social And Medical sciences (SIMSAM) framework grant no 340-2013-5867, the Swedish Asthma and Allergy Association's Research Foundation, the Swedish Heart-Lung Foundation, grants provided by the Stockholm County Council regional agreement on medical training and clinical research (ALF

project) between Stockholm County Council and Karolinska Institutet and the Strategic Research Program in Epidemiology at Karolinska Institutet.

Conflict of Interest: The authors have no potential conflicts of interest to disclose.

Authors' contributions

Dr Holmberg conceptualized and designed the study, had full access to all of the data in the study, conducted the initial analysis and interpretation, drafted the manuscript and revised it critically. Dr Holmberg had final responsibility for the decision to submit for publication.

MSc Lundholm assisted in acquisition of data, took active part in the analysis and interpretation, drafting and critical revision of the manuscript for important intellectual content.

Dr Larsson participated in the conceptualization and design of the study, contributed to the analysis and interpretation of data and revision of the manuscript.

Professor Anckarsäter assisted in acquisition of data and critically reviewed the manuscript.

Professor Almqvist obtained funding for the study and participated in study design, analysis and interpretation, drafting and critical revision of the manuscript. Together with Dr Holmberg, Professor Almqvist had final responsibility for the decision to submit for publication. All authors approved the final manuscript as submitted.

Supplementary Material

Table S1. Study population characteristics, parent-reported asthma in questionnaire and screening ADHD diagnosis in a cohort of 20,072 Swedish twins born 1992 – 2002 with completed questionnaires at 9 or 12 years of age.

References

1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5 ed. Arlington, VA, USA: American Psychiatric Association; 2013.
2. Calam R, Gregg L, Goodman R. Psychological adjustment and asthma in children and adolescents: the UK Nationwide Mental Health Survey. *Psychosom Med*. 2005 Jan-Feb;67(1):105-10.
3. Polanczyk G, de Lima MS, Horta BL, Biederman J, Rohde LA. The worldwide prevalence of ADHD: a systematic review and meta-regression analysis. *Am J Psychiatry*. 2007 Jun;164(6):942-8.
4. Kwon HJ, Lee MY, Ha M, Yoo SJ, Paik KC, Lim JH, et al. The associations between ADHD and asthma in Korean children. *BMC Psychiatry*. 2014;14:70.
5. McQuaid EL, Weiss-Laxer N, Kopel SJ, Mitchell DK, Nassau JH, Wamboldt MZ, et al. Pediatric Asthma and Problems in Attention, Concentration, and Impulsivity: Disruption of the Family Management System. *Fam Syst Health*. 2008 Mar;26(1):16-29.
6. Schieve LA, Gonzalez V, Boulet SL, Visser SN, Rice CE, Van Naarden Braun K, et al. Concurrent medical conditions and health care use and needs among children with learning and behavioral developmental disabilities, National Health Interview Survey, 2006-2010. *Res Dev Disabil*. 2012 Mar-Apr;33(2):467-76.
7. Yuksel H, Sogut A, Yilmaz O. Attention deficit and hyperactivity symptoms in children with asthma. *J Asthma*. 2008 Sep;45(7):545-7.
8. Biederman J, Milberger S, Faraone SV, Guite J, Warburton R. Associations between childhood asthma and ADHD: issues of psychiatric comorbidity and familiarity. *J Am Acad Child Adolesc Psychiatry*. 1994 Jul-Aug;33(6):842-8.
9. Hammerness P, Monuteaux MC, Faraone SV, Gallo L, Murphy H, Biederman J. Reexamining the familial association between asthma and ADHD in girls. *J Atten Disord*. 2005 Feb;8(3):136-43.
10. McQuaid EL, Kopel SJ, Nassau JH. Behavioral adjustment in children with asthma: a meta-analysis. *J Dev Behav Pediatr*. 2001 Dec;22(6):430-9.
11. Blackman JA, Gurka MJ. Developmental and behavioral comorbidities of asthma in children. *J Dev Behav Pediatr*. 2007 Apr;28(2):92-9.
12. Mogensen N, Larsson H, Lundholm C, Almqvist C. Association between childhood asthma and ADHD symptoms in adolescence - a prospective population-based twin study. *Allergy*. 2011 Sep;66(9):1224-30.
13. White MV, Sander N. Asthma from the perspective of the patient. *J Allergy Clin Immunol*. 1999 Aug;104(2 Pt 2):S47-52.
14. Saricoban HE, Ozen A, Harmanci K, Razi C, Zahmacioglu O, Cengizlier MR. Common behavioral problems among children with asthma: is there a role of asthma treatment? *Ann Allergy Asthma Immunol*. 2011 Mar;106(3):200-4.
15. Almqvist C, Pershagen G, Wickman M. Low socioeconomic status as a risk factor for asthma, rhinitis and sensitization at 4 years in a birth cohort. *Clin Exp Allergy*. 2005 May;35(5):612-8.
16. Biederman J, Faraone SV. Attention-deficit hyperactivity disorder. *Lancet*. 2005 Jul 16-22;366(9481):237-48.
17. Gong T, Lundholm C, Rejnö G, Mood C, Långström N, Almqvist C. Parental socioeconomic status, childhood asthma and medication use - a population-based study. *PLoS One*. 2014;9(9):e106579.

18. Örtqvist AK, Lundholm C, Carlström E, Lichtenstein P, Cnattingius S, Almqvist C. Familial factors do not confound the association between birth weight and childhood asthma. *Pediatrics*. 2009 Oct;124(4):e737-43.
19. Pelsler LM, Buitelaar JK, Savelkoul HF. ADHD as a (non) allergic hypersensitivity disorder: a hypothesis. *Pediatr Allergy Immunol*. 2009 Mar;20(2):107-12.
20. Lichtenstein P, Svartengren M. Genes, environments, and sex: factors of importance in atopic diseases in 7-9-year-old Swedish twins. *Allergy*. 1997 Nov;52(11):1079-86.
21. Larsson H, Lichtenstein P, Larsson JO. Genetic contributions to the development of ADHD subtypes from childhood to adolescence. *J Am Acad Child Adolesc Psychiatry*. 2006 Aug;45(8):973-81.
22. Lichtenstein P, Sullivan PF, Cnattingius S, Gatz M, Johansson S, Carlström E, et al. The Swedish Twin Registry in the third millennium: an update. *Twin research and human genetics : the official journal of the International Society for Twin Studies*. 2006 Dec;9(6):875-82.
23. Gong T, Almqvist C, Bölte S, Lichtenstein P, Anckarsäter H, Lind T, et al. Exposure to Air Pollution From Traffic and Neurodevelopmental Disorders in Swedish Twins. *Twin research and human genetics : the official journal of the International Society for Twin Studies*. 2014 Sep 17:1-10.
24. Örtqvist AK, Lundholm C, Wettermark B, Ludvigsson JF, Ye W, Almqvist C. Validation of asthma and eczema in population-based Swedish drug and patient registers. *Pharmacoepidemiol Drug Saf*. 2013 Aug;22(8):850-60.
25. Rejnö G, Lundholm C, Gong T, Larsson K, Saltvedt S, Almqvist C. Asthma during Pregnancy in a Population-Based Study - Pregnancy Complications and Adverse Perinatal Outcomes. *PLoS One*. 2014;9(8):e104755.
26. Wettermark B, Hammar N, Fored CM, Leimanis A, Otterblad Olausson P, Bergman U, et al. The new Swedish Prescribed Drug Register-opportunities for pharmacoepidemiological research and experience from the first six months. *Pharmacoepidemiol Drug Saf*. 2007 Jul;16(7):726-35.
27. Asher MI, Keil U, Anderson HR, Beasley R, Crane J, Martinez F, et al. International Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods. *Eur Respir J*. 1995 Mar;8(3):483-91.
28. Larson T, Lundström S, Nilsson T, Selinus EN, Råstam M, Lichtenstein P, et al. Predictive properties of the A-TAC inventory when screening for childhood-onset neurodevelopmental problems in a population-based sample. *BMC Psychiatry*. 2013;13:233.
29. Lundholm C, Örtqvist AK, Lichtenstein P, Cnattingius S, Almqvist C. Impaired fetal growth decreases the risk of childhood atopic eczema: a Swedish twin study. *Clin Exp Allergy*. 2010 Jul;40(7):1044-53.
30. Lichtenstein P, De Faire U, Floderus B, Svartengren M, Svedberg P, Pedersen NL. The Swedish Twin Registry: a unique resource for clinical, epidemiological and genetic studies. *J Intern Med*. 2002 Sep;252(3):184-205.
31. Laird N. Analysis of longitudinal and cluster-correlated data. Alexandria, Virginia, US: Beachwood, Ohio: Institute of mathematical statistics: American statistical association; 2004.
32. Schmitt J, Buske-Kirschbaum A, Roessner V. Is atopic disease a risk factor for attention-deficit/hyperactivity disorder? A systematic review. *Allergy*. 2010 Dec;65(12):1506-24.
33. Fasmer OB, Riise T, Eagan TM, Lund A, Dilsaver SC, Hundal O, et al. Comorbidity of asthma with ADHD. *J Atten Disord*. 2011 Oct;15(7):564-71.

34. Tsai JD, Chang SN, Mou CH, Sung FC, Lue KH. Association between atopic diseases and attention-deficit/hyperactivity disorder in childhood: a population-based case-control study. *Ann Epidemiol*. 2013 Apr;23(4):185-8.
35. Halterman JS, Yoos HL, Conn KM, Callahan PM, Montes G, Neely TL, et al. The impact of childhood asthma on parental quality of life. *J Asthma*. 2004 Sep;41(6):645-53.
36. Verlaet AA, Noriega DB, Hermans N, Savelkoul HF. Nutrition, immunological mechanisms and dietary immunomodulation in ADHD. *Eur Child Adolesc Psychiatry*. 2014 Jul;23(7):519-29.
37. Hadjikhouri I, Loader P, Bracken M, Milner AD. Bronchodilator therapy and hyperactivity in preschool children. *Arch Dis Child*. 2002 Mar;86(3):202-3.
38. Chen MH, Su TP, Chen YS, Hsu JW, Huang KL, Chang WH, et al. Comorbidity of Allergic and Autoimmune Diseases Among Patients With ADHD: A Nationwide Population-Based Study. *J Atten Disord*. 2013 Feb 11 [Epub ahead of print].
39. Holmberg K, Sundelin C, Hjern A. Screening for attention-deficit/hyperactivity disorder (ADHD): can high-risk children be identified in first grade? *Child Care Health Dev*. 2013 Mar;39(2):268-76.
40. Sung V, Hiscock H, Sciberras E, Efron D. Sleep problems in children with attention-deficit/hyperactivity disorder: prevalence and the effect on the child and family. *Arch Pediatr Adolesc Med*. 2008 Apr;162(4):336-42.
41. Thomsen SF, Kyvik KO, Backer V. A note on twin-singleton differences in asthma. *Twin research and human genetics : the official journal of the International Society for Twin Studies*. 2008 Apr;11(2):161-4.
42. Robbers SC, Bartels M, van Oort FV, van Beijsterveldt CE, van der Ende J, Verhulst FC, et al. A twin-singleton comparison of developmental trajectories of externalizing and internalizing problems in 6- to 12-year-old children. *Twin research and human genetics : the official journal of the International Society for Twin Studies*. 2010 Feb;13(1):79-87.