



**Karolinska  
Institutet**

Karolinska Institutet

<http://openarchive.ki.se>

---

This is a Non Peer Reviewed Manuscript version of the following article, accepted for publication in *Clinical and experimental allergy*.

2017-04-24

# Depression or anxiety in adult twins is associated with asthma diagnosis but not with offspring asthma

Tedner, S.G.; Lundholm, C.; Olsson, H.; Almqvist, C.

---

<http://doi.org/10.1111/cea.12714>

<http://hdl.handle.net/10616/45889>

*If not otherwise stated by the Publisher's Terms and conditions, the manuscript is deposited under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.*



**Karolinska  
Institutet**

This is the peer reviewed version of the following article:

**Depression or anxiety in adult twins is associated with asthma diagnosis but not with offspring asthma**, which has been published in final form at

**DOI: <https://dx.doi.org/10.1111/cea.12714>**

This article may be used for non-commercial purposes in accordance with **Wiley Terms and Conditions for Self-Archiving**.

# Depression or anxiety in adult twins is associated with asthma diagnosis but not with offspring asthma

## Authors

Sandra G Tedner, MD<sup>1</sup>

Cecilia Lundholm, Msc<sup>1</sup>

Henrik Olsson<sup>1</sup>

Catarina Almqvist, MD PhD<sup>1,2</sup>

## Affiliations

<sup>1</sup>Dept of Medical Epidemiology and Biostatistics, Karolinska Institutet

<sup>2</sup>Lung and Allergy unit, Astrid Lindgren Children's Hospital, Karolinska University Hospital  
Stockholm, Sweden

## Corresponding author

Sandra G. Tedner, MD

Department of Medical Epidemiology and Biostatistics, Karolinska Institutet

PO Box 281, SE-171 77 Stockholm, SWEDEN

T. +46 70 543 65 88

F. +46 8 314975

E. [sandra.ganrud-tedner@hotmail.com](mailto:sandra.ganrud-tedner@hotmail.com)

## **Abstract**

### **Background:**

Asthma is common in both children and adults in the Western world, just like anxiety and depression. While some research has revealed that these diseases might share important environmental and pathophysiological aspects, the exact mechanisms still remain unclear.

### **Objective:**

To study the correlation firstly between depression or anxiety and asthma diagnosis in adult twins, and secondly the association between parental depression or anxiety and offspring asthma in children of twins.

### **Methods:**

In total, 24,685 adult twins aged 20-47 years were interviewed or completed a web-based questionnaire and their children were identified through the Multi-Generation Register.

Asthma diagnosis was obtained from the Patient Register and the Prescribed Drug Register.

Assessment of depression and anxiety was obtained from questionnaires using Center for Epidemiologic Studies Depression Scale (CES-D), Major Depression and Generalized

Anxiety Disorder (GAD) from DSM-IV. The association between depression or anxiety and

asthma was analyzed with logistic regression adjusting for confounders in twins and

offspring. To address genetic and familial environmental confounding we performed a co-twin analysis using disease-discordant twin pairs.

### **Results:**

We found an association between asthma and CES-D, major depression and GAD, e.g.

adjusted OR for major depression and register-based asthma 1.56(1.36-1.79). Most of the

point estimates remained in the co-twin control analysis, indicating that the association was

likely not due to genetic or familial environmental factors. There was no association between parental depression and/or anxiety and asthma diagnosis in the offspring which implies lack of genetic confounding.

### **Conclusions:**

We found an association between own asthma diagnosis and anxiety or depression, but not with offspring asthma. Our results indicate that the associations were not due to confounding from genes or environment shared by the twins.

### **Keywords**

CESD, Childhood asthma, Generalized anxiety, Major depression, Offspring asthma, Twin study

### **Abbreviations**

BMI        body mass index

CES-D      Center for Epidemiologic Studies Depression Scale

CI         confidence interval

DZ         dizygotic

GAD        Generalized anxiety disorder

HR         hazard ratio

LISA        The Longitudinal integration database for health insurance and labor market studies

MBR        Medical Birth Register

MD	Major Depression
MGR	Multi Generation Register
MZ	monozygotic
OR	odds ratio
PAR	Patient Register
SPDR	Swedish Prescribed Drug Register
STAGE	The Study of Twin Adults: Genes and Environment
STR	Swedish Twin Registry
WHO	World health organization

## Introduction

The prevalence of asthma among both adults and children is high in many countries around the world [1, 2]. Depression is also a common disease [3] and several studies have found a correlation between asthma and depression [4-7] [8]. Anxiety disorders with generalized anxiety (GAD) defined as an excessive worry and anxiety [9] is common as well and as reviewed by ten Thoren et al there is a higher incidence of anxiety disorders in asthmatic patients [10]. The association between asthma and depression/anxiety is not clearly understood even though several studies have been performed. As reviewed by Di Marco emerging evidence that the diseases share similar environmental and pathophysiological mechanisms suggest interactions between behavioral, neural, endocrine and immune processes [11].

Previous studies on the association between parental depression or anxiety and asthma in offspring have found higher risk of asthma in children in relation to prenatal maternal stress [12-14], maternal stress reported during the child's first year of life [15, 16] and parental depression [17, 18]. Moreover, children exposed to maternal stress during pregnancy or postpartum had an increased risk for wheezing, but children exposed during both periods showed a threefold increase in odds of infant wheeze, suggesting that a more prolonged period of maternal distress is even more likely to increase the risk of childhood wheeze [19]. We have recently reported from a nationwide population-based study that boys born to mothers with loss of an older child during the second trimester had 50% higher risk of asthma event at both 1-4 years and 7-12 years [20].

Genetic and environmental effects may play an important role in the association between depression or anxiety and offspring asthma, and have not yet been thoroughly

studied. The understanding of the influence of genes and familial environment for an association can be improved by using twin design. In a co-twin control design you use the co-twin of an affected twin as a control. Thereby you can adjust your estimates for confounding factors that the twins share, including factors that are unmeasured or even unmeasurable, e.g. genes and childhood environment. Stratification by zygosity may give further insight with regard to the relative importance of confounding by genes and family environment, since dizygotic twins share 50% of their segregating genes and monozygotic 100%.

We have previously studied adult twins and their spouses and adolescent children, and found some association between self-reported maternal anxiety and subjectively reported offspring asthma or breathlessness which may be due to familial effects [21]. This can be further studied in a population of adult twins and their children where a co-twin control design may help differentiate between environmental and genetic confounding of an association between depression or anxiety and offspring asthma.

To better understand if depression and/or anxiety could be causally associated with asthma in adults, we aimed to study the association between anxiety or depression and asthma diagnosis with adjustment for identified confounders. Further we also aimed to study the association between parental depression and/or anxiety and current or subsequent childhood asthma in the offspring and address genetic and shared environmental confounding using co-twin controls.

## Methods

### Study design and study population

The Study of Twin Adults: Genes and Environment (STAGE) study population was derived from the Swedish Twin Registry (STR). During 2005-2006 twins born 1959-1985 were invited to participate in an extensive telephone interview or web-based questionnaire on habits, diseases, diet, living conditions, work etc. to assess the impact of gene and environment on future development of different diseases [22]. The National Board of Health and Welfare holds several registers with information about demography and health. In the Swedish Prescribed Drug register (SPDR) all prescribed drugs dispensed at Swedish pharmacies since July 1, 2005 are included. The Patient Register (PAR) covers all public, in-patient care in Sweden from 1987 and onwards. In 2001 an out-patient part was included which covers approximate 75 % of the out-patients visits. The majority of Swedish children with asthma are followed by pediatricians in out-patient clinics. The Medical Birth Register (MBR) covers data on >98 % of all births in Sweden. The Personal Identity Number (PIN), unique for each resident, allows unambiguous linkage between these registers and registers held by Statistic Sweden such as the Multi-Generation Register (MGR) with links between parent and child and the Longitudinal integration database for health insurance and labor market studies (LISA by Swedish acronym) with data on integrating labor market, social sector, as well as educational data for all inhabitants 16 years or older with annual updates.

To derive our study population, firstly twins included in STAGE were defined as our twin cohort. Secondly they were linked to the MGR to identify those that were parents. Thirdly their children were identified from the MBR and defined as our offspring cohort. Additional data on the twins and their children was collected from the MBR and LISA and

subsequent development of childhood asthma obtained from the PAR (asthma diagnosis) and SPDR (asthma medication).

Adopted children were excluded. Twins who moved out of the country after the questionnaire was answered were also excluded. Twins that could not be linked between the registers were also excluded from further analyses.

## **Variables**

Parental depression or anxiety was assessed using self-reported scales for depression and anxiety included in the STAGE questionnaire.

Depression was defined with two different scales, firstly through the Center for Epidemiologic Studies Depression Scale (CES-D), an index of self-reported depressive symptoms in the last week. Originally developed by Radloff in 1977, the 20 item scale was considered to be too long and therefore short versions are more frequently used. We used the eleven-item Iowa short version, validated by Kohout [23]. Each item can give a score of 0-3 points, maximum total 33 points. A score larger than 8 was classified as a possible mild to moderate depression and a score equal or more than 12 was regarded as a possible major depression. We allowed two missing items, and the score of these two items were calculated using imputation of the mean of the individual's response to the non-missing items of the scale. If an individual's response had more than two missing items the score was set to missing. Secondly with the DSM-IV criteria for Major Depression where the specific symptoms of depression (such as depressed mood (feels empty, sad, hopeless, tearful), sleep disturbance (insomnia/hypersomnia), markedly diminished interest or pleasure from almost every activity, significant weight change (up/down) or change of appetite (increased/decreased), feeling inhibited or agitated, feeling of lack of energy, feeling of guilt

or worthlessness, diminished ability to concentrate, recurrent thoughts about death or suicide) were evaluated [24, 25]. To get a positive score at least 5 of the symptoms (at least one of depressed mood or loss of interest or pleasure) must have been present during at least two weeks, during most of the day every day and caused a significant impact of the patient's daily life. The symptoms should not be attributable to grief, abuse of drugs, medication or other somatic disease or injury and caused clinically significant distress or impairment in social, occupational or other important areas of functioning.

General Anxiety Disorder (GAD) according to DSM-IV criteria [24, 25] was categorized as positive when the symptoms of excessive anxiety and worry for specific situations or activities, which were difficult to control, were present during most of the day, almost every day in the last 6 months. In addition to this the patient had three or more of the following symptoms: feeling restless or on edge, easily get fatigued, difficulty concentrating or mind going blank, irritability, muscle tension, and sleep disturbance (difficulty falling or staying asleep or restless unsatisfying sleep). The anxiety, worry or physical symptoms caused clinically significant distress or impairment in social, occupational or other important areas of functioning and were not attributable to grief, abuse of drugs, medication or other somatic disease or injury.

## **Outcomes**

Register-based asthma in the adult twins was derived from the PAR and defined if ICD code: J45 and J46 as a primary or secondary diagnosis) and/or prescription of asthma drugs in the SPDR, defined as either: 1) any asthma medication except  $\beta_2$ -agonist dispensed at least twice, or 2) any asthma medication dispensed at least three times during one calendar year 2006-2010, identified with the Anatomical Therapeutic Chemical (ATC) codes R03BA (inhaled corticosteroids), R03AK (fixed combinations of  $\beta_2$ -agonist and corticosteroids),

R03DC (leukotriene receptor antagonists) or R03AC ( $\beta_2$ - agonist). Dispensed asthma medication from the SPDR and register based asthma diagnoses in PAR are suitable proxies for asthma diagnosis, according to a recently published validation study by Örtqvist et al [26].

Self-reported asthma was obtained from the STAGE questionnaire, defined if the question “do you have asthma?” was yes, and clarified further with current or previous symptoms, thus as in “asthma ever”. By the usage of two different variables we created an objective diagnosis that occurred after the STAGE questionnaire was answered as well as a more subjective one that occurred previously. We were also able to define a larger number of asthma cases, including those with a mild asthma.

The ICD codes are applied related to a diagnosis given either at hospital or emergency facilities but also in out-patient specialist clinics.

In the offspring younger than 4.5 years of age, childhood asthma was derived from the PAR and defined if ICD code: J45 and J46 as a primary or secondary diagnosis and prescription of asthma drugs in the SPDR, defined as either: 1) any asthma medication except  $\beta_2$ - agonist dispensed at least twice with at least two weeks between distributions, or 2) any asthma medication dispensed at least three times during one calendar year 2006-2010, identified with the Anatomical Therapeutic Chemical (ATC) codes R03BA, R03AK, R03DC and R03AC. We used this definition to avoid misclassification of asthma in younger children with obstructive bronchiolitis due to respiratory infections.

For children > 4.5 years childhood asthma was defined according to the adult definition above.

### **Covariates**

Covariates were collected from STR (twin age, gender, zygosity), the STAGE questionnaire (smoking and BMI), the MBR (child birth weight, age and gender), and from

LISA (educational level) and categorized as follows: parental age at the time of questionnaire (19-25, 26-30, 31-35, 36-40, 41-45, >46), gender (Male/Female), BMI (<18,5, 18.5-24,9, 25-29,9, >30 kg/m<sup>2</sup> or missing), smoking (yes, no or missing) and educational level; middle school 9 years or shorter, high school 3 years or shorter, college studies shorter than 3 years, college graduates or higher education or missing), and child age at the time of questionnaire (0-4.5, >4.5-12, >12-18, >18).

### **Statistical analysis**

Firstly, logistic regression was used to estimate odds ratios (OR) with 95% confidence intervals (CIs) for the association between anxiety/depression and own asthma diagnosis. Crude estimates are reported together with estimates adjusted for age, gender, bmi and educational level while using robust standard errors to account for clustering of observations within twin pairs.

Secondly, for the association between parental anxiety or depression and offspring asthma we estimated ORs adjusted for age of child, parental educational level, parental smoking, parental asthma and parental gender while using robust standard errors to account for clustering of observations within families.

We studied possible effect modification by parental gender by testing for interaction with parental depression or anxiety and offspring asthma using likelihood ratio test.

Co-twin controls was used to estimate the association between depression or anxiety and asthma when controlling for genetic (100% for MZ and 50% for DZ) and family environmental effects. We used conditional logistic regression to estimate ORs among twins

discordant for both exposure and outcome and we adjusted for BMI and educational level.

Twins with unknown zygosity were excluded from these analyses.

Statistical analyses were conducted using Stata release 12.1 (Stata Corp, College Station, TX, USA).

Permission for the study was obtained from the Regional Ethical Review board in Stockholm, Sweden.

## Results

In total, 25,383 twins (59.6%) responded to the questionnaire. Register data was available for 24,685 twins, 44.2 % of them were male and 15,720 had 32,561 biological children, according to the registers.

Table 1 displays background characteristics of the twins. The prevalence of BMI >30 kg/m<sup>2</sup> was 5.1 % and 22.0 % were overweight with BMI 25-29, 9. Those with the lowest level of education had the highest incidence of asthma. The prevalence of mild to moderate CESD was 13.4% and major CESD reported by 16.6%. In total, 3964 twins (16.1%) filled the criteria for Major Depression, for GAD the number was 950 (3.8%). The prevalence of register-based asthma defined from asthma medication in the Prescribed Drug Register and/or diagnose in the Patient Registry was 5.9% in twins. The prevalence of self-reported asthma in the questionnaire was 9.8%.

Table 2 displays general characteristics of the children of the twins. The prevalence of register based childhood asthma was 6.7 %.

Table 3 displays the associations between depression or anxiety and asthma diagnosis in the adult twins. There was a significant association between most measures of depression or anxiety and register-based or self-reported asthma outcomes. For example, there was an association between major CESD and register-based asthma that remained after adjusting for potentially confounding factors, adjusted OR 1.42(95% CI 1.23-1.64) and similarly for self-reported asthma. There was also an increased risk of register-based asthma diagnosis in twins with major depression, adjusted OR 1.56 (95% CI 1.36-1.79) and was stable also for self-reported asthma. In the group of twins with GAD, we found an even higher

risk of register based asthma compared to twins without GAD, adjusted OR 1.73 (95 % CI 1.37-2.19), and this was also similar in the group with self-reported asthma.

Table 4 displays the association between parental depression or anxiety in the twins and asthma in their children. There was no association between any measures of depression or anxiety in the adult twins and register-based asthma in their children. For example, children of parents with mild to moderate CES-D did not have an increased risk of asthma, adjusted OR 1.03 (95% CI 0.90-1.18). The results were similar for parents with major depression as well for those with GAD diagnosis. We also performed a sub analysis in this group with children age 0-6 and age 11-22 years old, with similar estimates. For the studied associations, there was no effect modification by parental sex.

The result of the co-twin control analysis for the association between asthma diagnosis and depression or anxiety in the adult twins is displayed in table 5. In general, point estimates found in cohort analyses remained in the co-twin analyses although with broad confidence intervals and mostly not statistically significant.

For register-based asthma, (n=530 twin pairs) the association between major CESD and asthma after adjustment for confounders was OR 1.20 (0.87-1.65). There was an association between major depression and asthma that remained after adjusting for all confounders, adjusted OR 1.97 (95% CI 1.41-2.76). When stratifying by zygosity there was no significant difference in ORs between MZ and DZ. There was a significant association between parental GAD and register-based asthma, crude OR 1.71 (1.00-2.93) but this result did not remain significant after adjusting for measured confounders.

For self-reported asthma (n=666 twin pairs), there was an association between major CESD and asthma in MZ + DZ twins, which remained significant after adjusting for confounders, adjusted OR 1.44( 95% CI 1.10-1.89). When stratifying on zygosity the association only remained in the group with DZ twins, with an adjusted OR 2.01 (95% CI

1.21-3.35). However, the difference between MZ and DZ pairs was not statistically significant. There was an association between major depression and asthma outcomes, crude OR 1.28 (95 % CI 1.00-1.62) but this did not remain significant after adjusting for confounders, and for GAD we found no statistically significant association with self-reported asthma.

## Discussion

We found a clear association between depression or anxiety and asthma in twins, and this result remained significant after adjusting for confounding factors. The point estimates remained in the co-twin analyses, which indicates that the association was not due to confounding from genes or shared environmental factors. There was no association between parental depression and/or anxiety and asthma diagnosis in the offspring which along with the co-twin analysis implies lack of genetic confounding.

Our finding that there is an association between depression and asthma and between anxiety and asthma is in line with previous studies [4-7],[10] [27, 28]. As reviewed by Di Marco et al [11] this correlation has been confirmed in a large cross-sectional worldwide study and present research is focusing on the interaction between emotional stress, the immune system and the hypothalamic–pituitary–adrenocortical (HPA) and sympathetic–adrenal–medullary (SAM) axes.

We did not find a correlation between parental depression and/or anxiety and offspring asthma. Since previous studies have shown conflicting results it's possible that it depends on the selection of cohort, age groups [15-19, 29] as well as a difference in how offspring asthma diagnosis was determined. Kozyrkyj et al followed a birth cohort of children until the age of 7, and found that continued exposure to maternal distress was associated with higher incidence of childhood asthma OR 1.25 (95 % CI 1.01-1.55) [15]. They defined childhood asthma as two physician visits for asthma, one hospitalization, or two prescriptions for any asthma medication (incl. b-2 agonist, inhaled corticosteroids, leukotrienes or chromons) at the year of 7. Maternal distress was defined as physicians' visits, hospitalization or prescribed medication for anxiety or depression. Some cohorts only followed children during infancy (until 2 and 3 years of age) which is when transient wheeze is more common than asthma diagnose [16, 17, 19]. Some studies used questionnaires about

asthma diagnose filled by parents themselves [16, 19] which might lead to a subjective diagnose rather than objective.

We have previously shown an association between maternal anxiety and subjectively reported offspring asthma but not with register-based offspring asthma diagnosis or medication [21], which is in line with our results in this study. Some previous studies are made solely on women or on pregnant women [12-14]. Like our study most studies have used self-reported questionnaires to estimate level of parental stress/distress/anxiety and this could lead to problems with classification of exposure, both in terms of over- and underestimating the presence of anxiety or depression. Different scales were used in different cohort which also makes it hard to compare.

More women than men suffer from major depression, as previous research also have found [3, 30-32]. More women also suffer from general anxiety disorder GAD, which also confirms previous results [30]. One of few studies taking both parents in account was Lange et al who studied if maternal and paternal psychosocial stress respectively was associated with childhood asthma in a cohort of Puerto Rican twins and found some evidence of a correlation with both maternal and paternal stress [17] at age 1. However at age 3 years this correlation disappeared for the fathers but remained for maternal stress. In this study the offspring asthma diagnosis was not register based, only subjectively reported by parents.

Our association between depression or anxiety and asthma in twins remained in the co-twin analyses, which indicates that it most likely is causal or due to non-shared environmental factors such as medication, smoking habits, exposure to high levels of air pollution or other diseases. There was no association between parental depression and/or anxiety and asthma diagnosis in the offspring which along with the co-twin analysis implies absence of genetic confounding.

## **Strengths and limitations**

### *Strengths*

Firstly, we used a large cohort which results in enough power in the main analysis and well validated scales helping minimizing the level of misclassification of exposure.

Secondly, due to the usage of register data from PAR and SPDR, we defined a diagnosis of asthma that was obtained after the questionnaire was filled, and we got an objective asthma diagnosis since dispensed asthma medication from the SPDR and register based asthma diagnoses in PAR have high positive predicted values as proxies for asthma diagnosis [26]. We could however miss some of those who were sick but didn't seek care, those who got a diagnosis but got the wrong one, those who had a mild asthma that did not need referral to specialist care or did not need any medication or those who were prescribed medications but did not collect them at the pharmacy or not often enough. The self-reported asthma from STAGE answers to asthma ever and is likely to include at least some of those not captured in the registers; a phenomenon which could be reflected by the higher prevalence of self-reported asthma from STAGE was compared to register-based asthma.

Thirdly we accounted for several important confounding factors such as parental asthma, age, gender as well as smoking, leading to a good level of internal validity. Register-based information about educational level minimizes recall bias.

Fourthly, we performed a twin study including a co-twin control analysis and also studied offspring data. Studying several generations help to provide pieces to the puzzle on disease development.

### *Limitations*

Not all invited twins responded in this cohort, possibly due to the fact that the STAGE questionnaire was extensive, and also contained many questions of sensitive character, which might lead to less answers that are true or loss of answers. There is also a risk of differential ascertainment such that a twin with depression/anxiety could be more likely to receive asthma diagnosis than a twin without depression. Such misclassification could lead to overestimation of the association between asthma and depression.

Non-responders were more often male, from a lower educational level, had at least one parent born abroad or had been admitted for psychiatric disorders to a higher degree than responders. We lack information on asthma status in the non-responders. However, unless the differences between responders and non-responders are large with regard to both diseases, we would not expect large biases in the associations [33].

In the co-twin analysis, numbers were small which gave us low power, in particular for detecting differences in associations between MZ and DZ twins.

Self-reported data about depression and/ or anxiety might lead to misclassification of exposure, even though the scales in the questionnaire had high validity. Physician diagnosis of depression or GAD would of course be optimal. In PAR data from primary health care facilities are not included, which is where many patients with mild/moderate asthma normally would seek help. They would however be found in the SPDR if they used enough medication so this should not affect the result [26].

Asthma is a chronic disease causing breathlessness and one could argue that depression/anxiety could be a response to that. Depressive symptoms are also described as a possible side effect of inhaled corticosteroid treatment, however we found stable estimates related to all asthma outcomes regardless of medication. Furthermore, we could not establish the temporal sequence of the phenotypes asthma and depression/anxiety. Nor did we have information about atopic asthma. We could not thoroughly account for both parents since we

had information only about one of them and, as always, residual confounding can not be ruled out.

### **Generalizability**

Generalizability regarding twin studies can be argued, but as indicated in recent studies twins [34] [35] [36] higher risk of asthma compared to singletons appears to be due to lower gestational age and birth weight.

### **Future studies**

A new study with information about the mother before and during pregnancy, followed by annual questionnaires for both parents as well as bio samples like saliva-cortisol from both parents and child, use of registers for information about birth data, siblings, socioeconomic status, asthma diagnose as well as depression/ anxiety could lead to new insights.

### **Conclusions**

We found a significant correlation between own asthma and GAD and major depression, and most point estimates remained when using co-twin controls. This indicates that the association could not be explained by familial environment and genes shared by twins. We did not find a correlation between parental depression and/or anxiety and offspring asthma.

## **Acknowledgements**

We would like to acknowledge Christina Norrby and Rozita Broumandi who contributed with excellent database management and Professor Nancy Pedersen for discussions on study design and data collection. This work was supported by grants provided from Stockholm County Council (ALF projects), the Swedish Research Council (grant number 2011-3060) and through the Swedish Initiative for Research on Microdata in the Social And Medical Sciences (SIMSAM) framework (grant number 340-2013-5867), FORTE, the Swedish Heart-Lung foundation, and the Strategic Research Program in Epidemiology at Karolinska Institutet.

The authors have no conflict of interest to declare.

## References


1. Asher MI, Montefort S, Bjorksten B, Lai CK, Strachan DP, Weiland SK, Williams H, Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet* 2006;368: 733-43.
2. Masoli M, Fabian D, Holt S, Beasley R, The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 2004;59: 469-78.
3. Bromet E, Andrade LH, Hwang I, Sampson NA, Alonso J, de Girolamo G, de Graaf R, Demyttenaere K, Hu C, Iwata N, Karam AN, Kaur J, Kostyuchenko S, Lepine JP, Levinson D, Matschinger H, Mora ME, Browne MO, Posada-Villa J, Viana MC, Williams DR, Kessler RC, Cross-national epidemiology of DSM-IV major depressive episode. *BMC medicine* 2011;9: 90.
4. Delmas MC, Guignon N, Chee CC, Fuhrman C, Herbet JB, Gonzalez L, Asthma and major depressive episode in adolescents in France. *The Journal of asthma : official journal of the Association for the Care of Asthma* 2011;48: 640-6.
5. Scott KM, Von Korff M, Ormel J, Zhang MY, Bruffaerts R, Alonso J, Kessler RC, Tachimori H, Karam E, Levinson D, Bromet EJ, Posada-Villa J, Gasquet I, Angermeyer MC, Borges G, de Girolamo G, Herman A, Haro JM, Mental disorders among adults with asthma: results from the World Mental Health Survey. *Gen Hosp Psychiatry* 2007;29: 123-33.
6. Walters P, Schofield P, Howard L, Ashworth M, Tylee A, The relationship between asthma and depression in primary care patients: a historical cohort and nested case control study. *PloS one* 2011;6: e20750.
7. Wamboldt MZ, Hewitt JK, Schmitz S, Wamboldt FS, Rasanen M, Koskenvuo M, Romanov K, Varjonen J, Kaprio J, Familial association between allergic disorders and depression in adult Finnish twins. *Am J Med Genet* 2000;96: 146-53.
8. Goodwin RD, Fergusson DM, Horwood LJ, Asthma and depressive and anxiety disorders among young persons in the community. *Psychol Med* 2004;34: 1465-74.
9. American Psychiatric Association. *Psychiatric Press I, Washington, DC: 2000, Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition Revised. Text revision. Fourth Edition Revised Edn: Psychiatric Press, Inc., Washington, DC, 2000.*
10. ten Thoren C, Petermann F, Reviewing asthma and anxiety. *Respir Med* 2000;94: 409-15.
11. Di Marco F, Santus P, Centanni S, Anxiety and depression in asthma. *Current opinion in pulmonary medicine* 2011;17: 39-44.
12. Beijers R, Jansen J, Riksen-Walraven M, de Weerth C, Maternal prenatal anxiety and stress predict infant illnesses and health complaints. *Pediatrics* 2010;126: e401-9.
13. Cookson H, Granell R, Joinson C, Ben-Shlomo Y, Henderson AJ, Mothers' anxiety during pregnancy is associated with asthma in their children. *J Allergy Clin Immunol* 2009;123: 847-53 e11.
14. de Marco R, Pesce G, Girardi P, Marchetti P, Rava M, Ricci P, Marcon A, Foetal exposure to maternal stressful events increases the risk of having asthma and atopic diseases in childhood. *Pediatr Allergy Immunol* 2012;23: 724-9. doi: 10.1111/j.399-3038.2012.01346.x. Epub 2012 Sep 9.
15. Kozyrskyj AL, Mai XM, McGrath P, Hayglass KT, Becker AB, Macneil B, Continued exposure to maternal distress in early life is associated with an increased risk of childhood asthma. *American journal of respiratory and critical care medicine* 2008;177: 142-7.
16. Wright RJ, Cohen S, Carey V, Weiss ST, Gold DR, Parental stress as a predictor of wheezing in infancy: a prospective birth-cohort study. *Am J Respir Crit Care Med* 2002;165: 358-65.
17. Lange NE, Bunyavanich S, Silberg JL, Canino G, Rosner BA, Celedon JC, Parental psychosocial stress and asthma morbidity in Puerto Rican twins. *The Journal of allergy and clinical immunology* 2011;127: 734-40 e1-7.

18. Brown ES, Gan V, Jeffress J, Mullen-Gingrich K, Khan DA, Wood BL, Miller BD, Gruchalla R, Rush AJ, Psychiatric symptomatology and disorders in caregivers of children with asthma. *Pediatrics* 2006;118: e1715-20.
19. Mathilda Chiu YH, Coull BA, Cohen S, Wooley A, Wright RJ, Prenatal and postnatal maternal stress and wheeze in urban children: effect of maternal sensitization. *Am J Respir Crit Care Med* 2012;186: 147-54. doi: 10.1164/rccm.201201-0162OC. Epub 2012 May 10.
20. Fang F, Olgart Hoglund C, Arck P, Lundholm C, Langstrom N, Lichtenstein P, Lekander M, Almqvist C, Maternal bereavement and childhood asthma-analyses in two large samples of Swedish children. *PLoS One* 2011;6: e27202.
21. Havland I, Lundholm C, Lichtenstein P, Neiderhiser JM, Ganiban JM, Spotts EL, Walum H, Reiss D, Almqvist C, The observed association between maternal anxiety and adolescent asthma: children of twin design suggest familial effects. *PLoS One* 2013;8: e66040. doi: 10.1371/journal.pone.0066040. Print 2013.
22. Lichtenstein P, Sullivan PF, Cnattingius S, Gatz M, Johansson S, Carlstrom E, Bjork C, Svartengren M, Wolk A, Klareskog L, de Faire U, Schalling M, Palmgren J, Pedersen NL, The Swedish Twin Registry in the third millennium: an update. *Twin Res Hum Genet* 2006;9: 875-82.
23. Kohout FJ, Berkman LF, Evans DA, Cornoni-Huntley J, Two shorter forms of the CES-D (Center for Epidemiological Studies Depression) depression symptoms index. *Journal of aging and health* 1993;5: 179-93.
24. Lux V, Kendler KS, Deconstructing major depression: a validation study of the DSM-IV symptomatic criteria. *Psychol Med* 2010;40: 1679-90.
25. Diagnostic criteria from DSM-IV. Washington, D.C.: The Association, 1994.
26. Ortqvist AK, Lundholm C, Wettermark B, Ludvigsson JF, Ye W, Almqvist C, Validation of asthma and eczema in population-based Swedish drug and patient registers. *Pharmacoepidemiology and drug safety* 2013;22: 850-60.
27. Goodwin RD, Jacobi F, Thefeld W, Mental disorders and asthma in the community. *Arch Gen Psychiatry* 2003;60: 1125-30.
28. Cooper CL, Parry GD, Saul C, Morice AH, Hutchcroft BJ, Moore J, Esmonde L, Anxiety and panic fear in adults with asthma: prevalence in primary care. *BMC Fam Pract* 2007;8: 62.
29. Goodwin RD, Canino G, Ortega AN, Bird HR, Maternal mental health and childhood asthma among Puerto Rican youth: the role of prenatal smoking. *The Journal of asthma : official journal of the Association for the Care of Asthma* 2009;46: 726-30.
30. McLean CP, Asnaani A, Litz BT, Hofmann SG, Gender differences in anxiety disorders: prevalence, course of illness, comorbidity and burden of illness. *Journal of psychiatric research* 2011;45: 1027-35.
31. Naninck EF, Lucassen PJ, Bakker J, Sex differences in adolescent depression: do sex hormones determine vulnerability? *Journal of neuroendocrinology* 2011;23: 383-92.
32. Takkinen S, Gold C, Pedersen NL, Malmberg B, Nilsson S, Rovine M, Gender differences in depression: a study of older unlike-sex twins. *Aging & mental health* 2004;8: 187-95.
33. Furberg H, Lichtenstein P, Pedersen NL, Thornton L, Bulik CM, Lerman C, Sullivan PF, The STAGE cohort: a prospective study of tobacco use among Swedish twins. *Nicotine Tob Res* 2008;10: 1727-35.
34. Hemminki K, Li X, Sundquist K, Sundquist J, Familial risks for asthma among twins and other siblings based on hospitalizations in Sweden. *Clin Exp Allergy* 2007;37: 1320-5.
35. Thomsen SF, Kyvik KO, Backer V, A note on twin-singleton differences in asthma. *Twin Res Hum Genet* 2008;11: 161-4. doi: 10.1375/twin.11.2.161.
36. Ullemar V, Lundholm C, Almqvist C, Twins' Risk of Childhood Asthma Mediated By Gestational Age and Birth Weight. *Clin Exp Allergy* 2015;2: 12547.



**Figure 1** Flow chart of the cohort

Figure 1.



	Total, N (%)	Register-based Asthma, N (%)		Self-reported Asthma, N (%)		
		No	Yes	No	Yes	Missing
<b>Age</b>						
19-25	3171 (12.8)	3019 (13.0)	152 (10.4)	2305 (12.3)	361 (15.0)	505 (15.0)
26-30	4383 (17.8)	4136 (17.8)	247 (16.7)	3194 (17.1)	504 (20.9)	685 (220.4)
31-35	4396 (17.8)	4157 (17.9)	239 (16.3)	3358 (17.9)	413 (17.1)	625 (18.6)
36-40	4672 (18.9)	4370 (18.8)	302 (20.6)	3576 (19.1)	440 (18.3)	656 (19.5)
41-45	5263 (21.3)	4942 (21.3)	321 (21.9)	4097 (21.9)	462 (19.2)	704(21.0)
>46	2800 (11.3)	2596 (11.2)	204 (13.9)	2192 (11.7)	229 (9.5)	379 (11.3)
<b>Gender</b>						
Male	10905 (44.2)	10410 (44.8)	495 (31.0)	8239 (44.0)	974 (40.4)	1692 (47.6)
Female	13780 (55.8)	12810 (33.8)	970 (66.2)	10483 (56.0)	1435 (59.6)	1862 (52.4)
<b>BMI</b>						
<18.5	577 (2.3)	543 (2.3)	34 (2.3)	477 (2.5)	44 (1.8)	56 (1.6)
18.5-24.9	15164 (61.4)	14367 (61.9)	797 (54.4)	12229 (65.3)	1464 (60.8)	1471 (41.4)
25-29.9	5433 (22.0)	5097 (21.9)	336 (22.9)	4387 (23.4)	584 (24.2)	462 (13.0)
>30	1272 (5.1)	1111 (4.8)	161 (11.0)	954 (5.1)	199 (8.3)	119 (3.3)
Missing	2239 (9.1)	2102 (9.0)	137 (9.3)	675 (3.6)	118 (4.9)	1446 (40.7)
<b>Smoking</b>						
Non smokers	18435 (74.7)	17409 (75.0)	1026 (70.0)	14254 (76.1)	1780 (73.9)	2401 (67.5)
Regular smokers <sup>1</sup>	5321 (21.6)	4941 (21.3)	380 (25.9)	4028 (21.5)	561 (23.3)	732 (20.6)
Missing	929 (3.8)	870 (3.7)	59 (4.0)	440 (2.3)	68 (2.8)	421 (11.8)
<b>Education Sun 2000 Level</b>						
Middle school	1495 (6.1)	1386 (6.0)	109 (7.4)	1086 (5.8)	168 (7.0)	241 (7.2)
High school	11964 (48.5)	11249 (48.4)	715 (48.8)	9012 (48.1)	1202 (49.9)	1750 (52.2)
College < 3 years	4469 (18.1)	4210 (18.1)	259 (17.7)	3416 (18.2)	435 (18.0)	618 (18.4)
College graduates/higher education > 3 years <sup>2</sup>	6487 (26.3)	6115 (26.3)	372 (25.4)	5015 (26.8)	576 (23.9)	896 (26.7)
Missing	270 (1.1)	260 (1.1)	10 (0.7)	193 (1.0)	28 (1.2)	49 (1.5)
<b>CES-D</b>						
No	14181 (57.4)	13423 (57.8)	758 (51.7)	12312 (65.8)	1406 (58.4)	463 (13.0)
Mild to moderate	3317 (13.4)	3103 (13.3)	214 (14.6)	2852 (15.2)	387 (16.1)	78 (2.2)
Major	4109 (16.6)	3772 (16.2)	337 (23.0)	3387 (18.1)	593 (24.6)	129(3.6)
Missing	3078 (12.5)	2922 (12.6)	156 (10.6)	171 (0.9)	23 (0.9)	2884 (81.1)
<b>Major Depression</b>						
No	17373 (70.4)	16455 (70.9)	918 (62.6)	15108 (80.7)	1789 (74.3)	476 (13.4)
Yes	3964 (16.1)	3597 (15.5)	367 (25.0)	3210 (17.1)	577 (23.9)	177 (5.0)
Missing	3348 (13.6)	3168 (13.6)	180 (12.3)	404 (2.1)	43 (1.8)	2901 (81.6)
<b>GAD</b>						
No	20714 (83.9)	19610 (84.4)	1104 (75.3)	15517 (82.9)	1868 (77.5)	3329 (93.7)
Yes	950 (3.8)	849 (3.6)	101 (6.9)	768 (4.1)	148 (6.1)	34 (0.9)
Missing	3021(12.2)	2761 (11.9)	260 (17.7)	2437 (13.0)	393 (16.3)	191(5.4)

<sup>1</sup> Smoke daily and who have smoked at least 28 cigarettes in a month

<sup>2</sup> Including doctoral and licentiate degree

**Table 2:** General information of children of STAGE twins

	<u>Total, N (%)</u>	<u>Asthma in children, N (%)</u>	
		No	Yes
<b>Age</b>			
0-4.5	6100 (9.2)	5822 (19.1)	278 (12.8)
>4.5-12	10875 (14.0)	10073 (33.1)	802 (36.9)
>12-18	8176 (25.1)	7507 (24.7)	669 (30.8)
>18	7410 (22.8)	6987 (23.0)	423 (19.5)
<b>Gender</b>			
Boys	15730 (48.3)	14790 (48.7)	940 (43.3)
Girls	16831 (51.7)	15599 (51.3)	1232 (56.7)
<b>Exposed to parental</b>			
<b>CES-D</b>			
No	19759 (60.7)	18423 (60.6)	1336 (61.5)
Mild to moderate	4154 (12.7)	3859 (12.7)	295 (13.6)
Major	4758 (14.6)	4444 (14.6)	314 (14.4)
Missing	3890 (11.9)	3663 (12.0)	227 (10.4)
<b>Major Depression</b>			
No	23437 (72.0)	21861 (71.9)	1576 (72.5)
Yes	4887 (15.0)	4528 (14.9)	352 (16.2)
Missing	4244 (13.0)	4000 (13.2)	244 (11.2)
<b>GAD</b>			
No	27615 (84.8)	25775 (84.8)	1840 (84.7)
Yes	1111 (3.4)	1033 (3.4)	78 (3.6)
Missing	3835 (11.8)	3581 (11.8)	254 (11.7)

**Table 3:** Odds ratios and 95 % confidence intervals for the association between parental anxiety/depression and asthma in STAGE-twins

	<b>Register-based Asthma</b>			<b>Self-reported Asthma</b>		
	Crude OR (95% CI)	Adj * OR (95 % CI)	Adj ^^ OR (95 % CI)	Crude OR (95% CI)	Adj * OR (95 % CI)	Adj ^^ OR (95 % CI)
<b>CES-D</b>						
No	1.00	1.00	1.00	1.00	1.00	1.00
Mild to moderate	1.22 (1.04-1.42)	1.25 (1.07-1.47)	1.24 (1.05-1.46)	1.19 (1.05-1.34)	1.17 (1.04-1.32)	1.17 (1.03-1.33)
Major	1.58 (1.38-1.81)	1.56 (1.36-1.79)	1.42 (1.23-1.64)	1.53 (1.38-1.70)	1.48 (1.33-1.64)	1.42 (1.27-1.58)
<b>Major Depression</b>						
No	1.00	1.00	1.00	1.00	1.00	1.00
Yes	1.83 (1.61-2.07)	1.70 (1.49-1.93)	1.56 (1.36-1.79)	1.52 (1.37-1.68)	1.49 (1.35-1.66)	1.44 (1.29-1.61)
<b>GAD</b>						
No	1.00	1.00	1.00	1.00	1.00	1.00
Yes	2.11 (1.70-2.62)	1.98 (1.60-2.46)	1.73 (1.37-2.19)	1.60 (1.33-1.92)	1.56 (1.29-1.87)	1.44 (1.18-1.75)

\* adjusted for age & gender

^^ adjusted for age, gender, bmi, educational level, smoking

**Table 4:** Odds ratios and 95 % confidence intervals for the association between parental anxiety/depression and asthma in children of STAGE-twins

	<b>Asthma in children</b>		
	Crude OR (95 % CI)	Adj OR † (95 % CI)	Adj OR ☐ (95 % CI)
<b>CES-D</b>			
No	1.00	1.00	1.00
Mild to moderate	1.05 (0.92-1.20)	1.05 (0.92-1.20)	1.03 (0.90-1.18)
Major	0.97 (0.86-1.11)	0.97 (0.86-1.11)	0.93 (0.81-1.06)
<b>Major Depression</b>			
No	1.00	1.00	1.00
Yes	1.08 (0.96-1.21)	1.09 (0.96-1.23)	1.06 (0.94-1.20)
<b>GAD</b>			
No	1.00	1.00	1.00
Yes	1.06 (0.84-1.34)	1.07 (0.84-1.35)	0.99 (0.78-1.27)

† adjusted for child age

☐ adjusted for parental (smoking, gender, educational level, parents own asthma) + child age

**Table 5: Co-twin analysis of asthma in STAGE-twins exposed to anxiety and depression**

Outcome	Register-based asthma (n=530 twin pairs)				Self-reported asthma (n= 666 twin pairs)			
	MZ+DZ	Exposure asthma twin	Crude OR (95% CI)	Exposure asthma twin Adj OR * (95 % CI)	Exposure asthma twin	Crude OR (95% CI)	Exposure asthma twin	Adj OR * (95 % CI)
<b>CES-D</b>		221		190	290		252	
No		83	1.00	74	105	1.00	91	1.00
Mild /moderate		60	1.17 (0.85-1.62)	52	81	1.09 (0.84-1.42)	70	1.11 (0.84-1.48)
Major		78	1.32 (0.63-1.58)	64	104	1.43 (1.12-1.82)	91	1.44 (1.10-1.89)
<b>Major Depression</b>		129		111	187		164	
No		44	1.00	38	85	1.00	77	1.00
Yes		85	2.10 (1.54-2.85)	73	102	1.28(1.00-1.62)	87	1.20 (0.92-1.56)
<b>GAD</b>		36		26	44		36	
No		14	1.00	9	19	1.00	15	1.00
Yes		22	1.71 (1.00-2.93)	17	25	1.53 (0.95-2.47)	21	1.46 (0.85-2.48)
MZ	Exposure asthma twin	Crude OR (95% CI)	Exposure asthma twin	Adj OR* (95 % CI)	Exposure asthma twin	Crude OR (95% CI)	Exposure asthma twin	Adj OR* (95 % CI)
<b>CES-D</b>	117		100		146		129	
No	44	1.00	38	1.00	54	1.00	48	1.00
Mild /moderate	30	1.14 (0.66-1.96)	26	1.05 (0.58-1.88)	44	1.27 (0.81-2.01)	38	1.21 (0.75-1.96)
Major	43	1.17 (0.74-1.86)	36	1.14 (0.67-1.94)	48	1.28 (0.83-1.98)	43	1.46 (0.90-2.38)
<b>Major Depression</b>	65		54		91		43	
No	23	1.00	19	1.00	45	1.00	21	1.00
Yes	42	1.83 (1.10-3.04)	35	1.83 (1.02-3.29)	46	1.02 (0.68-1.54)	22	1.01 (0.64-1.60)
DZ	Exposure asthma twin	Crude OR (95% CI)	Exposure asthma twin	Adj OR * (95 % CI)	Exposure asthma twin	Crude OR (95% CI)	Exposure asthma twin	Adj OR * (95 % CI)
<b>CES-D</b>	104		90		144		123	
No	39	1.00	36	1.00	51	1.00	43	1.00
Mild /moderate	30	1.15 (0.67-1.97)	26	1.19 (0.64-2.20)	37	1.06 (0.67-1.68)	32	1.14 (0.68-1.91)
Major	35	1.25 (0.75-2.10)	28	1.10 (0.61-1.97)	56	1.89 (1.20-2.99)	48	2.01 (1.21-3.35)
<b>Major Depression</b>	64		57		96		87	
No	21	1.00	19	1.00	40	1.00	38	1.00
Yes	43	2.05 (1.21-3.45)	38	1.91 (1.09-3.37)	56	1.40 (0.93-2.10)	49	1.23 (0.80-1.90)

\* adjusted for bmi, educational level, smoking