Errata

AKADEMISK AVHANDLING

Stable High-Sensitivity Cardiac Troponin T Levels and Outcomes

should be:

Andreas Roos

Thesis:

FIGURE 10: For study II, the box showing exclusion of 11,731 patients due to hs-cTnT levels ≤14

ng/l.

Section: Selection bias, p. 70: "In study II-IV, as previously

described..."

TABLE 5: Rates for

cardiovascular and non-

cardiovascular mortality presented

as %.

TABLE 5 & 6: Adjusted HRs for

all outcomes

TABLE 9: eGFR (mL/min/1.73m2) 45-60

TABLE 13: "ACEi/ARB = angiotensin-converting-enzyme inhibitor/angiotensinogenreceptor-blocker; eGFR =

estimated glomerular filtration rate; MI = myocardial infarction."

Paper 1:

Data sources: "≤12 ng/l"

Suppl. Material, p. 6: ">2 ng/l and an admission troponin level ≤12

ng/l"

Paper 2:

Table 3: Odds ratios (OR)

Paper 3:

Methods: "and/or one

measurement of hs-cTnT level

≤12 ng/l"

Paper 4:

Section 2.3. Data resources: "and deaths from the Cause of Death

register."

Suppl. Material, p. 4: "<5 ng/L to 12 ng/L"

should be:

should be:

The box should be placed beneath, and not above, the box showing

the number of 14,246 patients with

a measured hs-cTnT level.

"In study I, III and IV, as previously described..."

No "%" throughout the table,

analogously with all the other outcomes (presented as numbers

per 100 person-years).

HRs should be adjusted also for hypertension. For correct HRs, see

Table 2 in paper 1.

eGFR (mL/min/1.73m2) 15-60

"ACEi/ARB = angiotensin

converting enzyme inhibitor/angiotensinogenreceptor blocker; eGFR = estimated glomerular filtration rate; MI = myocardial infarction."

"<12 ng/l"

">2 ng/l and an admission troponin level <12 ng/l"

Risk ratios (RR)

"and/or one measurement of hs-

cTnT level <12 ng/l"

"and deaths from the Cause of

Death register".

"<5 ng/L to 11 ng/L"